

A derived isometry theorem for sheaves

Nicolas Berkouk, Grégory Ginot

January 9, 2023

Abstract

Persistent homology has been recently studied with the tools of sheaf theory in the derived setting by Kashiwara and Schapira [?] after J. Curry has made the first link between persistent homology and sheaves.

We prove the isometry theorem in this derived setting, thus expressing the convolution distance of sheaves as a matching distance between combinatorial objects associated to them that we call graded barcodes. This allows to consider sheaf-theoretical constructions as combinatorial, stable topological descriptors of data, and generalizes the situation of persistence with one parameter. To achieve so, we explicitly compute all morphisms in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, which enables us to compute distances between indecomposable objects. Then we adapt Bjerkevik's stability proof to this derived setting.

As a byproduct of our isometry theorem, we prove that the convolution distance is closed, give a precise description of connected components of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ and provide some explicit examples of computation of the convolution distance.

Contents

1	Introduction	1
2	Preliminaries	4
2.1	The isometry theorem problem for metric Krull-Schmidt categories	4
2.2	Notations for sheaves and complexes	6
2.3	Constructible sheaves over \mathbb{R}	6
2.4	Metric for sheaves	7
2.5	The isometry theorem problem for $(D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}}), d_C)$	10
3	Computations in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$	10
3.1	Duality and morphisms in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$	10
3.2	The functor $- \star K_{\varepsilon}$	13
4	Structure of ε-interleavings	14
4.1	Characterization of ε -interleavings between indecomposable sheaves	14
4.2	CLR Decomposition	15
5	Isometry theorem and graded barcodes	17
5.1	The easy inequality	17
5.2	The cases $F_R \leftrightarrow G_R$ and $F_L \leftrightarrow G_L$	17
5.2.1	Construction of Ψ_R^j	18
5.2.2	Matching of the central parts	19
5.3	The case $F_C \leftrightarrow G_C$	20
5.3.1	Ordering graded-intervals of type C	20

5.3.2	Induced matching	21
5.4	Isometry theorem	23
6	Applications	24
6.1	Example : projection from the circle	24
6.2	About the closedness of d_C	25
6.3	Description of the connected components of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$	28
6.4	Algorithmic remarks on computing one best matching	31

1 Introduction

Persistence theory appeared in the early 2000’s as an attempt to make some constructions inspired by Morse theory computable in practice. For instance, in the context of studying the underlying topology of a data set. It has since been widely developed and applied in many ways. We refer the reader to [?, ?] for extended expositions of the theory and of its applications. One promising expansion of the theory, initiated by Curry in his Ph.D. thesis [?], is to combine the powerful theory of sheaves with ideas coming from persistence, which are driven by applications in machine learning. However, sheaf theory takes its full strength in the derived setting and Kashiwara and Schapira developed persistent homology in this new framework in [?]. In this paper, we show that the main theorems of one-parameter persistence theory (which we recall below) admit an analogue in the context of derived sheaves on the real line, equipped with the convolution distance.

To our knowledge, this is the first result allowing to actually use sheaves as topological descriptors for noisy data sets.

One-parameter persistence

The theory of one parameter persistence mainly relies on one construction and two theorems that we now explain. Given a real-valued function f on a topological space X and $i \in \mathbb{Z}$, consider $\mathcal{S}_i(f)(s) := H_i(f^{-1}(-\infty; s))$ the i -th singular homology group of the sublevel set of f with coefficient in the fixed field \mathbf{k} . Then for $s \leq t$, the inclusion $f^{-1}(-\infty; s) \subset f^{-1}(-\infty; t)$ induces a linear map $\mathcal{S}_i(f)(s) \rightarrow \mathcal{S}_i(f)(t)$, and the functorial nature of singular homology gives $\mathcal{S}_i(f)$ the structure of a functor from the poset category (\mathbb{R}, \leq) to the category $\text{Mod}(\mathbf{k})$ of \mathbf{k} -vector spaces. This functor, that we still write as $\mathcal{S}_i(f)$, is usually referred to as the i -th sublevel-sets persistence module associated to f . More generally, the category $\text{Pers}(\mathbf{k}^{\mathbb{R}})$ of *persistence modules over* \mathbb{R} is precisely the category of functors $(\mathbb{R}, \leq) \rightarrow \text{Mod}(\mathbf{k})$.

Observe that the category $\text{Pers}(\mathbf{k}^{\mathbb{R}})$ of persistence modules over \mathbb{R} is a Grothendieck category, so that in particular it admits small limits and colimits. In [?], Crawley-Boevey proved that under some finiteness assumptions on f , satisfied for instance when $\mathcal{S}_i(f)(s)$ is finite-dimensional at every $s \in \mathbb{R}$, $\mathcal{S}_i(f)$ decomposes as a locally finite direct sum of persistence modules which are constant, with values \mathbf{k} , and supported on a given list of intervals of \mathbb{R} . This list of intervals entirely characterizes the isomorphism class of $\mathcal{S}_i(f)$ and is referred to as the i -th *barcode* of f , written $\mathbb{B}_i(f)$. One fundamental property that implies that the whole theory can be handled by a computer is that $\mathbb{B}_i(f)$ is a complete, discrete invariant of $\mathcal{S}_i(f)$.

On the other hand, for $\mathbb{B}_i(f)$ to be a meaningful descriptor of real-word –hence noisy– data it must satisfies some form of stability with respect to f . More precisely, it is important to understand under which distance $\mathbb{B}_i(f)$ and $\mathbb{B}_i(g)$ are close, for f and g two functions ε -close in L_{∞} -norm, that is in uniform convergence distance. An answer was first given in

2005 by D. Cohen-Steiner, H. Edelsbrunner, and J. Harer in [?] and is now referred to as the stability theorem. It states that if $f, g : X \rightarrow \mathbb{R}$ are ε -close in the L_∞ -norm, then there exists a one-to-one pairing between the intervals of $\mathbb{B}_i(f)$ and $\mathbb{B}_i(g)$, such that the right (resp. left) endpoints of each interval within a pair are closer than ε , and intervals can be paired to 0 if they have length less than 2ε . Such a pairing is called an ε -matching between $\mathbb{B}_i(f)$ and $\mathbb{B}_i(g)$, and we can define the bottleneck distance between $\mathbb{B}_i(f)$ and $\mathbb{B}_i(g)$ to be the infimum of the values of ε for which there exists an ε -matching between $\mathbb{B}_i(f)$ and $\mathbb{B}_i(g)$. The stability theorem can now then be restated as follows : the bottleneck distance between $\mathbb{B}_i(f)$ and $\mathbb{B}_i(g)$ is less or equal than the L_∞ -norm of $f - g$.

In 2009, Chazal, Cohen-Steiner, Glisse, Guibas, and Oudot [?] expressed the stability theorem algebraically, introducing the interleaving distance between one-parameter persistence modules and proving that an ε -interleaving (a kind of approximate isomorphism) induces an ε -matching between their associated barcodes. This statement is usually referred to as either the algebraic stability theorem or the *isometry theorem*, and is the cornerstone of persistence techniques in the one-parameter case together with Crawley-Boevey's theorem [?].

Persistence and sheaves

The need for studying persistence modules obtained from functions valued in higher-dimensional vector spaces naturally arises from the context of data analysis, see for example [?, ?]. However, as shown in [?], the category $\text{Pers}(\mathbf{k}^{\mathbb{R}^n})$ of functors $(\mathbb{R}^n, \leq) \rightarrow \text{Mod}(\mathbf{k})$ seems to be too general for $n \geq 2$. Indeed, it contains a full sub-category equivalent to the one of finitely generated \mathbb{Z}^n -graded $\mathbf{k}[x_1, \dots, x_n]$ -modules, which implies that there is no hope for a barcode like decomposition when $n \geq 2$. There are mainly two directions undertaken to handle this issue.

The first one, initiated by Magnus Botnan and Michael Lesnick [?, ?], then pursued by Bjerkevik in [?], Cochoy and Oudot in [?], consists in restricting the study of $\text{Pers}(\mathbf{k}^{\mathbb{R}^n})$ to simpler sub-categories, for example, the one of persistence modules that admit a decomposition into interval modules as in the one parameter case. In [?], Bjerkevik proves that the bottleneck distance of two interval decomposable modules is bounded by a multiple (depending on the number of parameters) of the interleaving distance, and in [?], Cochoy and Oudot prove that a certain kind of persistence modules over \mathbb{R}^2 , namely the pointwise-finite dimensional exact bi-modules, actually have an interval decomposition. Nevertheless, it remains unclear how likely is the interval decomposable case to appear in practice.

On the other hand, it seems natural to treat persistence modules as sheaves, which is precisely what Justin Curry initiated in his Ph. D. thesis [?] by expressing persistence ideas in the formalism of (co-)sheaves of vector spaces on topological spaces. In particular, he defined a distance on the category of sheaves inspired by the interleaving distance, based on convolution. He also asked whether in the case of sheaves over \mathbb{R} , this distance could be expressed as a bottleneck distance. In 2018, Kashiwara and Schapira [?] introduced independently a derived version of the constructions of persistence theory in the category of sheaves on euclidean vector spaces, defining the convolution distance on its derived category, proving a stability theorem and introducing a promising notion of higher-dimensional barcodes for a large category : the γ -piecewise linear sheaves [?].

Content of the paper

In this paper, we provide answers to the question asked by Justin Curry at the end of his thesis, in the setting of Kashiwara and Schapira. We will explain later on our choice

and motivations to work in this derived setting.

It follows from general theorems that $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ – the bounded derived category of constructible sheaves on \mathbb{R} – is a Krull-Schmidt category, whose indecomposable objects are constant sheaves over some real interval, concentrated in one degree. Recording all of the interval appearing in this decomposition, together with their degree, lead to the notion of graded barcode. Since Kashiwara and Schapira have equipped this category with the convolution distance, which is inspired by the interleaving distance, a natural question is whether the convolution distance between two sheaves in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ can be computed as a matching between graded intervals, in the same fashion as the well-known isometry theorem for one parameter persistence. We provide a positive answer to this question, proving a derived isometry theorem in this setting. One important feature of our work is that the matching distance associated to the convolution distance allows to match intervals across degree, this being due to the fundamentally derived nature of the convolution distance. We hope that this result will open the door to considering sheaf-theoretical constructions as tools for applications in machine learning. In particular, one future direction of research is to elucidate the implications of this derived isometry theorem with level-sets persistence, which we have undertaken in [?]. Our result might also be of independent interest for mathematical fields where barcodes techniques have allowed to obtain new results, such as symplectic topology [?, ?].

The paper is structured as follows :

1. Section 2 aims at introducing the mathematical context of the paper. We give a general definition of the isometry theorem problem for metric Krull-Schmidt categories. We also explain the convolution distance and the decomposition result of constructible sheaves over \mathbb{R} obtained by Kashiwara and Schapira in [?].
2. Section 3 is dedicated to the complete description of the morphisms in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, the derived category of constructible sheaves on \mathbb{R} , and to compute the action of the convolution functor $-\star K_{\varepsilon}$. Note that the computations of this section (propositions 3.7 and 3.5), may be of independent interest.
3. Section 4 describes the conditions for two indecomposables sheaves to be ε -close, and introduce the notion of CLR decomposition for any sheaf $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, which is adapted to the convolution distance in the following sense : two sheaves are ε -close with respect to d_C if and only if their central (resp. left, resp. right) parts are.
4. In Section 5, we prove that given an ε -interleaving between the central (resp. left, resp. right) parts of two sheaves, it induces an ε -matching between the graded-barcodes of their central (resp. left, resp. right) parts. We reduce the proof for left and right parts to the well-known case of one-parameter persistence modules by introducing a family of fully faithful functors from sheaves supported on half-open intervals to persistence modules. The construction of the ε -matching between the central parts is far less direct. We adapt the proof of Bjerkevik [?] to our setting, introducing a similar pre-order \leq_{α} on central parts, enabling us to “trigonalize” the interleaving morphisms. By a rank argument, this allows us to apply Hall’s marriage theorem and to deduce the existence of a ε -matching. Note that our definition of \leq_{α} differs in nature from Bjerkevik’s, for it enables us to compare elements of the graded-barcodes in *different degrees*. We conclude the section by proving the isometry theorem 5.10, which states that “ $d_C = d_B$ ”.
5. Section 6 provides some applications of the isometry theorem. We start by an example with explicit computations brought to our knowledge by Justin Curry and that motivated our work. Then, we prove that the convolution distance is closed

(two sheaves are ε -close if and only if they are ε -interleaved) thus answering an open question of [?] in dimension one. We provide a counter-example of two non constructible sheaves F and G such that $d_C(F, G) = 0$ but $F \not\simeq G$. We also prove that the open balls of the metric space $(D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}}), d_C)$ are path-connected, hence leading to a characterization of connected components of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ (Theorem 6.12). To do so, we introduce an explicit skeleton for $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ called the category **Barcodes** of barcodes (Definition 6.11). The latter inherits an extended metric space structure.

Acknowledgement The authors would like to thank Justin Curry, Steve Oudot, Pierre Schapira and Magnus Botnan for many enlightening discussions with them.

The second author was partially supported by ANR grants Catag and Chrok.

2 Preliminaries

This section aims at introducing the notation that we will use in this paper, presenting the theoretical framework of [?] and explaining precisely the problem underlying the isometry theorem.

2.1 The isometry theorem problem for metric Krull-Schmidt categories

Let \mathcal{C} be an additive category. Recall that an object $M \in \mathcal{C}$ is indecomposable if $M \neq 0$, and for any isomorphism $M \simeq M_1 \oplus M_2$, either M_1 or M_2 is equal to 0.

DEFINITION 2.1 A category \mathcal{C} is a Krull-Schmidt category if it satisfies the following axioms.

(KS-1) \mathcal{C} is an additive category.

(KS-2) For any object X of \mathcal{C} , there exists a family of indecomposable objects $\mathbb{B}(X)$ of \mathcal{C} such that $X \simeq \bigoplus_{I \in \mathbb{B}(X)} I$ which is essentially unique. That is, for any other family of indecomposable objects $\mathbb{B}'(X)$ with the same property, there exists a bijection $\sigma : \mathbb{B}(X) \rightarrow \mathbb{B}'(X)$ such that $I \simeq \sigma(I)$, for all I in $\mathbb{B}(X)$.

(KS-3) For any object X of \mathcal{C} such that $X \simeq \bigoplus_{I \in \mathbb{B}(X)} I$ with $\mathbb{B}(X)$ a collection of indecomposable objects of \mathcal{C} , then $\prod_{I \in \mathbb{B}(X)} I$ exists in \mathcal{C} and the canonical morphism :

$$\bigoplus_{I \in \mathbb{B}(X)} I \longrightarrow \prod_{I \in \mathbb{B}(X)} I$$

is an isomorphism.

REMARK 2.2 Note that the usual definition of a Krull-Schmidt category asks, with notations of definition 2.1, that $\mathbb{B}(X)$ is finite (see *eg.* [?]). This will not be sufficient for our study of constructible sheaves over \mathbb{R} since they are potentially infinite direct sum of sheaves constant on a real interval (theorem 2.9). However, one important behaviour of these direct sums is that they satisfy axiom (KS-3), since they are locally finite.

DEFINITION 2.3 Let \mathcal{C} be any category. An extended pseudo-distance on \mathcal{C} is a map d defined on $\text{Obj}(\mathcal{C}) \times \text{Obj}(\mathcal{C})$ satisfying, for all X, Y, Z in \mathcal{C} :

(M1) $d(X, Y) \in \mathbb{R}_{\geq 0} \cup \{+\infty\}$,

(M2) $d(X, Y) = d(Y, X)$,

(M3) $d(X, Z) \leq d(X, Y) + d(Y, Z)$,

(M4) if $X \simeq Y$, then $d(X, Y) = 0$.

In this situation, (\mathcal{C}, d) will be called a *metric category*.

Let (\mathcal{C}, d) be a metric category, such that \mathcal{C} is a Krull-Schmidt category. For any object X of \mathcal{C} , one denotes by $\mathbb{B}(X)$ a collection of indecomposables objects of \mathcal{C} such that $X \simeq \bigoplus_{I \in \mathbb{B}(X)} I$.

DEFINITION 2.4 Let $X, Y \in \mathcal{C}$ and $\varepsilon \geq 0$. An ε -matching between $\mathbb{B}(X)$ and $\mathbb{B}(Y)$ is the following data: two subcollections $\mathcal{X} \subset \mathbb{B}(X)$ and $\mathcal{Y} \subset \mathbb{B}(Y)$, and a bijection $\sigma : \mathcal{X} \rightarrow \mathcal{Y}$ satisfying:

1. $d(I, \sigma(I)) \leq \varepsilon$ for all I in \mathcal{X} ;
2. $d(I, 0) \leq \varepsilon$, for all I in $\mathbb{B}(X) \setminus \mathcal{X}$ or in $\mathbb{B}(Y) \setminus \mathcal{Y}$.

Since d satisfies (M4), the existence of a ε -matching does not depend on the choice of representatives in $\mathbb{B}(X)$ and $\mathbb{B}(Y)$.

DEFINITION 2.5 Let X and Y be two objects of \mathcal{C} . One defines the bottleneck distance associated to d between X and Y as the possibly infinite following quantity:

$$d_B(X, Y) = \inf\{\varepsilon \geq 0 \mid \text{there exists an } \varepsilon\text{-matching between } \mathbb{B}(X) \text{ and } \mathbb{B}(Y)\}.$$

PROPOSITION 2.6 The map d_B is an extended pseudo-metric on \mathcal{C} .

PROOF The fact that d_B takes values in $\mathbb{R}_{\geq 0} \cup \{+\infty\}$ and Properties (M2) and (M4) are directly inherited from d . For the triangle inequality, observe that since d satisfies (M3), we can compose a ε -matching between $\mathbb{B}(X)$ and $\mathbb{B}(Z)$, with a ε' -matching between $\mathbb{B}(Z)$ and $\mathbb{B}(Y)$ to obtain a $\varepsilon + \varepsilon'$ matching between $\mathbb{B}(X)$ and $\mathbb{B}(Y)$.

Terminology. The isometry theorem problem associated to the metric Krull-Schmidt category (\mathcal{C}, d) is to determine whether $d = d_B$.

2.2 Notations for sheaves and complexes

Throughout the paper and except when stated otherwise, we will follow the notations introduced in [?] and [?]. We will also freely refer to some of their proofs.

In the paper, \mathbf{k} will denote a field, $\text{Mod}(\mathbf{k})$ the category of vector spaces over \mathbf{k} , $\text{Mod}_f(\mathbf{k})$ the category of finite dimensional vector spaces over \mathbf{k} . Let X be a topological space. Then we will note $\text{Mod}(\mathbf{k}_X)$ the category of sheaves of \mathbf{k} -vector spaces on X . For shortness, we will also write Hom for $\text{Hom}_{\text{Mod}(\mathbf{k}_{\mathbb{R}})}$.

For \mathcal{C} an abelian category, denote $C^b(\mathcal{C})$ its category of bounded complexes, $K^b(\mathcal{C})$ its bounded homotopy category and $D^b(\mathcal{C})$ its bounded derived category. For simplicity, we shall write $D^b(\mathbf{k})$ instead of $D^b(\text{Mod}(\mathbf{k}))$ and $D^b(\mathbf{k}_X)$ instead of $D^b(\text{Mod}(\mathbf{k}_X))$. When the context is clear, we will simply call sheaves the objects of $D^b(\mathbf{k}_X)$. For a complex $F \in C^b(\mathcal{C})$ and an integer k , define the k -th shift of F by: for $n \in \mathbb{Z}$, $F[k]^n = F^{k+n}$ and $d_{F[k]}^n = (-1)^k d_F^{n+k}$.

We will use the classical notations of [?] for the Grothendieck operations on sheaves. Moreover, we recall the following : for X_1 and X_2 two topological spaces, we denote $p_i : X_1 \times X_2 \rightarrow X_i$, $i = 1, 2$ the canonical projections. Let $F \in \text{Mod}(\mathbf{k}_{X_1})$ and $G \in \text{Mod}(\mathbf{k}_{X_2})$, define their external tensor product $F \boxtimes G \in \text{Mod}(\mathbf{k}_{X_1 \times X_2})$ by the formula :

$$F \boxtimes G := p_1^{-1}F \otimes p_2^{-1}G.$$

Observe that since we are working over a field, this operation is exact, hence need not to be derived.

DEFINITION 2.7 For M a real analytic manifold, and $F \in \text{Mod}(\mathbf{k}_M)$, F is said to be weakly \mathbb{R} -constructible if there exists a locally finite sub-analytic stratification of $M = \sqcup_{\alpha} M_{\alpha}$, such that for each stratum M_{α} , the restriction $F|_{M_{\alpha}}$ is locally constant. If in addition, if the stalks F_x are of finite dimension for every $x \in M$, we say that F is \mathbb{R} -constructible. We might often say constructible instead of \mathbb{R} -constructible, since, in this paper, it is the only notion of constructibility we use.

We will write $\text{Mod}_{\mathbb{R}c}(\mathbf{k}_M)$ for the abelian category of \mathbb{R} -constructible sheaves on M , and $D_{\mathbb{R}c}^b(\mathbf{k}_M)$ the full triangulated subcategory of $D^b(\mathbf{k}_M)$ consisting of complexes of sheaves whose cohomology objects lies in $\text{Mod}_{\mathbb{R}c}(\mathbf{k}_M)$. Note that Theorem 8.4.5 in [?] asserts that the natural functor $D^b(\text{Mod}_{\mathbb{R}c}(\mathbf{k}_M)) \rightarrow D_{\mathbb{R}c}^b(\mathbf{k}_M)$ is an equivalence of triangulated categories.

2.3 Constructible sheaves over \mathbb{R}

Theorem 2.9 below is proved in [?] and generalizes Crawley-Boevey's theorem [?] to the context of constructible sheaves on the real line. Together with Theorem 2.12, they will be the cornerstone to prove that $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ is a Krull-Schmidt category, hence leading to the definition of graded-barcodes Section 5.

DEFINITION 2.8 Let $\mathcal{I} = \{I_{\alpha}\}_{\alpha \in A}$ be a multi-set of intervals of \mathbb{R} , that is, a list of interval where one interval can appear several times. Then \mathcal{I} is said to be locally finite if and only if for every compact set $K \subset \mathbb{R}$, the set $\{\alpha \in A \mid K \cap I_{\alpha} \neq \emptyset\}$ is finite.

THEOREM 2.9 (THEOREM 1.17 - [?]) Let $F \in \text{Mod}_{\mathbb{R}c}(\mathbf{k}_{\mathbb{R}})$, then there exists a unique locally finite multi-set of intervals $\mathbb{B}(F)$ such that

$$F \simeq \bigoplus_{I \in \mathbb{B}(F)} \mathbf{k}_I.$$

Moreover, this decomposition is unique up to isomorphism.

DEFINITION 2.10 The multi-set $\mathbb{B}(F)$ is the called the barcode of F .

COROLLARY 2.11 Let $F, G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, and $j \geq 2$, then : $\text{Ext}^j(F, G) = 0$.

A classical consequence of such a statement is the following :

THEOREM 2.12 Let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$. Then there exists an isomorphism in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$:

$$F \simeq \bigoplus_{j \in \mathbb{Z}} H^j(F)[-j]$$

where $H^j(F)$ is seen as a complex concentrated in degree 0. Not that this isomorphism is not functorial.

DEFINITION 2.13 Let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, we define its *graded-barcode* $\mathbb{B}(F)$ as the collection $(\mathbb{B}^j(F))_{j \in \mathbb{Z}}$ where $\mathbb{B}^j(F) := \mathbb{B}(H^j(F))$. Furthermore, to indicate that an interval $I \subset \mathbb{R}$ appears in degree $j \in \mathbb{Z}$ in the graded-barcode of F , we will write $I^j \in \mathbb{B}(F)$. The element I^j is called a graded-interval.

Note that Theorems 2.9 and 2.12 imply that

$$F \simeq \bigoplus_{I^j \in \mathbb{B}(F)} \mathbf{k}_I[-j].$$

Therefore, $\mathbb{B}(F)$ is a complete discrete invariant of the isomorphism class of F in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$.

PROPOSITION 2.14 The category $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ is Krull-Schmidt.

PROOF The category $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ is additive by construction. Moreover, for $I \subset \mathbb{R}$ an interval and $j \in \mathbb{Z}$, the sheaf $\mathbf{k}_I[-j]$ is indecomposable in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$. Therefore, we deduce from Theorems 2.9 and 2.12 that $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ satisfies (KS2). Let us now prove that $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ satisfies (KS3). Let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, then there exists $N \geq 0$ such that $\mathbb{B}^j(F) = \emptyset$ for $|j| \geq N$. As a consequence, given $x \in \mathbb{R}$, the set $\{I \in \sqcup_j \mathbb{B}^j(F) \mid (x-1, x+1) \cap I \neq \emptyset\}$ is finite. We deduce that the natural morphism:

$$F \simeq \bigoplus_{I^j \in \mathbb{B}(F)} \mathbf{k}_I[-j] \longrightarrow \prod_{I^j \in \mathbb{B}(F)} \mathbf{k}_I[-j]$$

induces an isomorphism when taking the stalk at x . This being true for all $x \in \mathbb{R}$, it is an isomorphism. □

2.4 Metric for sheaves

In [?], Curry defined an interleaving-like distance on $\text{Mod}(\mathbf{k}_X)$, for (X, d) a metric space. It is based on what he calls the smoothing of opens. For $F \in \text{Mod}(\mathbf{k}_X)$, define $F^\varepsilon \in \text{Mod}(\mathbf{k}_X)$ as the sheafification of the mapping $U \rightarrow F(U^\varepsilon)$, with $U^\varepsilon = \{x \in X \mid \exists u \in U, d(x, u) \leq \varepsilon\}$. This yields a functor $[\varepsilon] : \text{Mod}(\mathbf{k}_X) \rightarrow \text{Mod}(\mathbf{k}_X)$ together with a natural transformation $[\varepsilon] \Rightarrow \text{id}_{\text{Mod}(\mathbf{k}_X)}$. Although this seems to mimic the construction of interleaving distance for persistence modules, one must pay attention to the fact that $[\varepsilon]$ is only left-exact. Since topological informations are obtained from sheaves by considering sheaf-cohomology, one needs to derive the functor $[\varepsilon]$ in order to keep track of cohomological informations while smoothing a sheaf. This is precisely the sense of the construction of Kashiwara and Schapira using convolution of sheaves, which has the advantage to have a nice expression in term of Grothendieck operations (that precisely allows appropriate operations for sheaf cohomology).

In this section, we make a short review of the concepts introduced in [?]. The framework is the study of sheaves on a real vector space \mathbb{V} of finite dimension n equipped with a norm $\|\cdot\|$. For two such sheaves, one can define their convolution, which, as the name suggests, will be at the core of the definition of the convolution distance (definition 2.20).

The construction of the convolution of sheaves is as follows. Consider the following maps (addition and the canonical projections):

$$\begin{aligned} s : \mathbb{V} \times \mathbb{V} &\rightarrow \mathbb{V}, & s(x, y) &= x + y, \\ q_i : \mathbb{V} \times \mathbb{V} &\rightarrow \mathbb{V} \quad (i = 1, 2) & q_1(x, y) &= x, \quad q_2(x, y) = y. \end{aligned}$$

DEFINITION 2.15 For $F, G \in D^b(\mathbf{k}_{\mathbb{V}})$, we define the convolution of F and G by the formula :

$$F \star G = \text{Rs}_1(F \boxtimes G).$$

This defines a bi-functor : $D^b(\mathbf{k}_{\mathbb{V}}) \times D^b(\mathbf{k}_{\mathbb{V}}) \rightarrow D^b(\mathbf{k}_{\mathbb{V}})$.

In the following, we will be interested in a more specific case : the convolution will be considered with one of the sheaves being the constant sheaf supported on a ball centered at 0.

For $r > 0$, we denote $B_r := \{x \in \mathbb{V} \mid \|x\| \leq r\}$ the closed ball of radius r centered at 0, and $\overset{\circ}{B}_r$ its interior, that is, the open ball of radius r centered at 0. For $\varepsilon \in \mathbb{R}$ we define :

$$K_\varepsilon := \begin{cases} \mathbf{k}_{B_\varepsilon} & \text{if } \varepsilon \geq 0 \\ \mathbf{k}_{\overset{\circ}{B}_{-\varepsilon}}[\dim(\mathbb{V})] & \text{if } \varepsilon < 0 \end{cases} \quad (1)$$

with $\mathbf{k}_{\overset{\circ}{B}_{-\varepsilon}}[\dim(\mathbb{V})]$, seen as a complex concentrated in degree $-\dim(\mathbb{V})$. We have the following properties:

PROPOSITION 2.16 (SECTION 2.1 - [?]) Let $\varepsilon, \varepsilon' \in \mathbb{R}$ and $F \in D^b(\mathbf{k}_{\mathbb{V}})$.

1. One has functorial isomorphisms $(F \star K_\varepsilon) \star K_{\varepsilon'} \simeq F \star K_{\varepsilon+\varepsilon'}$ and $F \star K_0 \simeq F$.
2. If $\varepsilon' \geq \varepsilon$, there is a canonical morphism $K_{\varepsilon'} \rightarrow K_\varepsilon$ in $D^b(\mathbf{k}_{\mathbb{V}})$ inducing a natural transformation $F \star K_{\varepsilon'} \rightarrow F \star K_\varepsilon$. In the special case where $\varepsilon = 0$, we shall write $\phi_{F, \varepsilon'}$ for this natural transformation.
3. The canonical morphism $F \star K_{\varepsilon'} \rightarrow F \star K_\varepsilon$ induces an isomorphism

$$\mathrm{R}\Gamma(\mathbb{V}; F \star K_{\varepsilon'}) \xrightarrow{\sim} \mathrm{R}\Gamma(\mathbb{V}; F \star K_\varepsilon)$$

and hence an isomorphism in cohomology.

DEFINITION 2.17 With the same notations as in the previous proposition, the morphism $\phi_{F, \varepsilon'}$ is called the ε' -smoothing morphism of F .

In particular, Proposition 2.16 implies that any map $f : F \star K_\varepsilon \rightarrow G$ induces canonical maps

$$f \star K_\tau : F \star K_{\varepsilon+\tau} \simeq F \star K_\varepsilon \star K_\tau \rightarrow G \star K_\tau. \quad (2)$$

The following definition is central.

DEFINITION 2.18 (DEFINITION 2.2 - [?]) For $F, G \in D^b(\mathbf{k}_{\mathbb{V}})$ and $\varepsilon \geq 0$, one says that F and G are ε -interleaved if there exists two morphisms $f : F \star K_\varepsilon \rightarrow G$ and $g : G \star K_\varepsilon \rightarrow F$ (in $D^b(\mathbf{k}_{\mathbb{V}})$) such that the compositions $F \star K_{2\varepsilon} \xrightarrow{f \star K_\varepsilon} G \star K_\varepsilon \xrightarrow{g} F$ and $G \star K_{2\varepsilon} \xrightarrow{g \star K_\varepsilon} F \star K_\varepsilon \xrightarrow{f} G$ are the natural morphisms $F \star K_{2\varepsilon} \xrightarrow{\phi_{F, 2\varepsilon}} F$ and $G \star K_{2\varepsilon} \xrightarrow{\phi_{G, 2\varepsilon}} G$, that is, we have a commutative diagram in $D^b(\mathbf{k}_{\mathbb{V}})$:

$$\begin{array}{ccccc} & & \phi_{F, 2\varepsilon} & & \\ & & \curvearrowright & & \\ & & & & \\ F \star K_{2\varepsilon} & \xrightarrow{f \star K_\varepsilon} & G \star K_\varepsilon & \xrightarrow{g} & F \\ & \searrow & \nearrow & \searrow & \nearrow \\ G \star K_{2\varepsilon} & \xrightarrow{g \star K_\varepsilon} & F \star K_\varepsilon & \xrightarrow{f} & G \\ & \nearrow & \searrow & \nearrow & \searrow \\ & & \phi_{G, 2\varepsilon} & & \end{array}$$

In this case, we write $F \sim_\varepsilon G$.

Observe that F and G are 0-interleaved if and only if $F \simeq G$.

REMARK 2.19 One must be aware that in [?], the authors call this data an ε -isomorphism. Here, we choose to follow the usual terminology of persistence theory.

Since 0-interleaving are isomorphism, the existence of an ε -interleaving between two sheaves expresses a notion of closeness. This leads the authors of [?] to define the convolution distance as follows:

DEFINITION 2.20 (DEFINITION 2.2 - [?]) For $F, G \in D^b(\mathbf{k}_{\mathbb{V}})$, we define their convolution distance as:

$$d_C(F, G) := \inf(\{+\infty\} \cup \{a \in \mathbb{R}_{\geq 0} \mid F \text{ and } G \text{ are } a\text{-interleaved}\})$$

PROPOSITION 2.21 (SECTION 2.2 - [?]) The convolution distance is an extended pseudo-distance on $D^b(\mathbf{k}_{\mathbb{V}})$ that is, it satisfies (M1)-(M4) from definition 2.3.

The following proposition expresses that the functors $\mathrm{R}\Gamma(\mathbb{V}; -)$ and $\mathrm{R}\Gamma_c(\mathbb{V}; -)$ define some necessary conditions for two sheaves to be at finite convolution distance. This is similar to the case of interleaving distance for persistence modules $M : (\mathbb{R}^n, \leq) \rightarrow \mathrm{Mod}(\mathbf{k})$, where the role of $\mathrm{R}\Gamma(\mathbb{V}; -)$ is played by the colimit functor over \mathbb{R}^n .

PROPOSITION 2.22 (REMARK 2.5 - [?]) Let $F, G \in D^b(\mathbf{k}_{\mathbb{V}})$.

1. If $d_C(F, G) < +\infty$ then $\mathrm{R}\Gamma(\mathbb{V}, G) \simeq \mathrm{R}\Gamma(\mathbb{V}, F)$ and $\mathrm{R}\Gamma_c(\mathbb{V}; G) \simeq \mathrm{R}\Gamma_c(\mathbb{V}; F)$.
2. If $\mathrm{supp}(F), \mathrm{supp}(G) \subset B_a$ then $d_C(F, G) \leq 2a$ if and only if $\mathrm{R}\Gamma(\mathbb{V}, G) \simeq \mathrm{R}\Gamma(\mathbb{V}; F)$.

There is a fundamental example to keep in mind in the context of sheaves. This example is the one mimicking the persistence modules $\mathcal{S}_i(f)$ of a continuous map: given X a topological space and $u : X \rightarrow \mathbb{V}$ a continuous map, one can consider the sheaves $Ru_*\mathbf{k}_X$ and $Ru!\mathbf{k}_X$. Roughly speaking and under some smoothness assumptions on X and f , they contain the information on how the cohomologies of the fibers of u evolve when moving on \mathbb{V} . For this information to be meaningful for applications in machine learning, it has to be stable when we perturb u , that is, $Ru_*\mathbf{k}_X$ must stay in a neighborhood in the sense of the convolution distance, controlled by the size of the perturbation of u . This is what expresses the following theorem, which is the analogous of the stability theorem in the context of persistence theory.

THEOREM 2.23 (THEOREM 2.7 - [?]) Let X a locally compact topological set, and $u, v : X \rightarrow \mathbb{V}$ two continuous functions. Then for any $F \in D^b(\mathbf{k}_X)$ one has :

$$d_C(Ru_*F, Rv_*F) \leq \|u - v\| \quad \text{and} \quad d_C(Ru!F, Rv!F) \leq \|u - v\|$$

where we define $\|u - v\| = \sup_{x \in X} \|u(x) - v(x)\|$.

2.5 The isometry theorem problem for $(D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}}), d_C)$

In the previous sections, we have shown that $(D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}}), d_C)$ is a metric Krull-Schmidt category, see Propositions 2.14 and 2.21. We will prove later (see 5.10) that the isometry theorem holds in this context, that is, the convolution distance equals its associated bottleneck distance. In other words, we can compute the convolution distance between two sheaves $F, G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ as a matching distance between the multiset of graded-intervals appearing in their decomposition, which we will later call their graded-barcodes (see Definition 2.10). This matching between barcodes will be similar to the usual matching between barcodes (that is in terms of comparing end points of bars and overall length) but will take into account shift of degrees and types of bars as well.

3 Computations in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$

This section aims at making explicit all the computations of morphisms in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, and determine the action of the functor $- \star K_\varepsilon$.

Combining Theorems 2.9 and 2.12, we see that any object of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ is isomorphic to a direct sum of sheaves constant on an interval seen as a complex concentrated in one degree. Hence, to give a full description of the morphisms, it is enough to compute $\mathrm{RHom}_{\mathrm{Mod}(\mathbf{k}_{\mathbb{R}})}(\mathbf{k}_I, \mathbf{k}_J)$ for I, J two intervals – in the sequel, we shall write for short Hom for $\mathrm{Hom}_{\mathrm{Mod}(\mathbf{k}_{\mathbb{R}})}$. Indeed, for $F, G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ we have by local finiteness of the barcodes of F and G :

$$\mathrm{RHom}(F, G) \simeq \bigoplus_{\substack{I^i \in \mathbb{B}(F) \\ J^j \in \mathbb{B}(G)}} \mathrm{RHom}(\mathbf{k}_I, \mathbf{k}_J)[i - j],$$

and we recall the classical formula $\mathrm{Hom}_{D^b(\mathbf{k}_{\mathbb{R}})}(F, G) \simeq \mathrm{R}^0\mathrm{Hom}(F, G)$.

Our approach¹ will rely on the use of the duality functor D introduced, for instance, in [?, 3.1.16], that we will quickly review.

Our computations for the convolution with K_ε will make use of a classical lemma for convolution of sheaves, that we will also review.

3.1 Duality and morphisms in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$

Since \mathbb{R} is a smooth manifold, it has finite c -soft dimension, and the dualizing complex $\omega_{\mathbb{R}} := a^! \mathbf{k}_{\{pt\}} \simeq \mathbf{k}_{\mathbb{R}}[1]$ is well defined, where $a : \mathbb{R} \rightarrow \{pt\}$. One defines the triangulated functor $D : D^b(\mathbf{k}_{\mathbb{R}})^{\mathrm{op}} \rightarrow D^b(\mathbf{k}_{\mathbb{R}})$, by $D(F) = \mathrm{R}\mathcal{H}om(F, \omega_{\mathbb{R}})$. We recall the following proposition, which is a weaker version of [?, 3.4.3 and 3.4.6] adapted to our setting, noting that constructible sheaves are in particular cohomologically constructible.

PROPOSITION 3.1 Let $F, G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, then:

1. the canonical map $F \xrightarrow{\simeq} D \circ D(F)$ is an isomorphism,
2. $\mathrm{R}\mathcal{H}om(F, G) \simeq D(D(G) \otimes F)$.

For an open subset $U \subset \mathbb{R}$, one has $D(\mathbf{k}_U) \simeq \mathbf{k}_{\overline{U}}[1]$, where \overline{U} is the adherence of U , from which we deduce the following computations:

PROPOSITION 3.2 Let $a < b \in \mathbb{R}$, then:

1. $D(\mathbf{k}_{\{a\}}) \simeq \mathbf{k}_{\{a\}}$,
2. $D(\mathbf{k}_{[a,b]}) \simeq \mathbf{k}_{(a,b)}[1]$.

PROOF To prove these results, one simply applies successively the contravariant triangulated functor D to the distinguished triangles

$$\mathbf{k}_{\mathbb{R} \setminus \{a\}} \rightarrow \mathbf{k}_{\mathbb{R}} \rightarrow \mathbf{k}_{\{a\}} \xrightarrow{+1} \quad \text{and} \quad \mathbf{k}_{(a,b)} \rightarrow \mathbf{k}_{[a,b]} \rightarrow \mathbf{k}_{\{a\}} \xrightarrow{+1}.$$

□

1. which was suggested by the referee, that we wish to thank.

By definition, given $F, G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, one has $\mathrm{RHom}(F, G) \simeq \mathrm{R}\Gamma(\mathbb{R}; \mathrm{R}\mathcal{H}om(F, G))$. We now review the computations of derived global sections of indecomposable constructible sheaves on \mathbb{R} .

PROPOSITION 3.3 Let $a < b \in \mathbb{R}$, then:

1. $\mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{[a, +\infty)}) \simeq \mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{(-\infty, b]}) \simeq \mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{[a, b]}) \simeq \mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{\{a\}}) \simeq \mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{\mathbb{R}}) \simeq \mathbf{k}$,
2. $\mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{(a, +\infty)}) \simeq \mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{(-\infty, b)}) \simeq 0$,
3. $\mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{(a, b)}) \simeq \mathbf{k}[-1]$,
4. $\mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{[a, b]}) \simeq \mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{(a, b]}) \simeq 0$.

PROOF 1. Let I be any of the interval $[a, +\infty)$, $(-\infty, b]$, $[a, b]$, $\{a\}$ and $i : I \rightarrow \mathbb{R}$ be the inclusion. Then by definition $\mathbf{k}_I = i_* i^{-1} \mathbf{k}_{\mathbb{R}}$, and we have the following isomorphisms:

$$\begin{aligned} \mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_I) &\simeq \mathrm{RHom}(\mathbf{k}_{\mathbb{R}}; i_* i^{-1} \mathbf{k}_{\mathbb{R}}) \\ &\simeq \mathrm{RHom}(i^{-1} \mathbf{k}_{\mathbb{R}}; i^{-1} \mathbf{k}_{\mathbb{R}}) \\ &\simeq \mathrm{R}\Gamma(I; \mathbf{k}_I) \\ &\simeq \mathbf{k}. \end{aligned}$$

2. This is a consequence of 1. by applying the triangulated functor $\mathrm{R}\Gamma(\mathbb{R}; -)$ to the distinguished triangles $\mathbf{k}_{(a, +\infty)} \rightarrow \mathbf{k}_{\mathbb{R}} \rightarrow \mathbf{k}_{(-\infty, a]} \xrightarrow{+1}$ and $\mathbf{k}_{(-\infty, b]} \rightarrow \mathbf{k}_{\mathbb{R}} \rightarrow \mathbf{k}_{[b, +\infty)} \xrightarrow{+1}$.
3. This is a consequence of 2. by applying the triangulated functor $\mathrm{R}\Gamma(\mathbb{R}; -)$ to the distinguished triangle $\mathbf{k}_{(a, b)} \rightarrow \mathbf{k}_{\mathbb{R}} \rightarrow \mathbf{k}_{(-\infty, a]} \oplus \mathbf{k}_{[b, +\infty)} \xrightarrow{+1}$.
4. This is a consequence of 3. by applying the triangulated functor $\mathrm{R}\Gamma(\mathbb{R}; -)$ to the distinguished triangles $\mathbf{k}_{[a, b]} \rightarrow \mathbf{k}_{[a, b]} \rightarrow \mathbf{k}_{\{b\}} \xrightarrow{+1}$ and $\mathbf{k}_{(a, b]} \rightarrow \mathbf{k}_{[a, b]} \rightarrow \mathbf{k}_{\{a\}} \xrightarrow{+1}$.

□

REMARK 3.4 Let $a \leq b \in \mathbb{R}$. For simplicity, we adopt the following convention. When stating results about the interval (a, b) , we will always implicitly assume that $a < b$.

PROPOSITION 3.5 Let $a \leq b$ and $c \leq d$ in \mathbb{R} . We have the following derived morphism groups, where the first column defines the support of the left-side object (i.e. the source) in $\mathrm{RHom}(-, -)$ and the first line the right-side one :

PROOF Since all computations work similarly, we shall only treat the case where $I = [a, b]$ and $J = (c, d) \neq \emptyset$. By proposition 3.1:

$$\mathrm{RHom}(\mathbf{k}_I, \mathbf{k}_J) \simeq \mathrm{R}\Gamma(\mathbb{R}; \mathrm{D}(\mathrm{D}(\mathbf{k}_J) \otimes \mathbf{k}_I)).$$

We have $\mathrm{D}(\mathbf{k}_J) \simeq \mathbf{k}_{[c,d]}[1]$. Therefore, $\mathrm{D}(\mathrm{D}(\mathbf{k}_J) \otimes \mathbf{k}_I) \simeq \mathrm{D}(\mathbf{k}_{[c,d] \cap [a,b]})[-1]$. If $[c, d] \cap [a, b] = \emptyset$, then $\mathrm{RHom}(\mathbf{k}_I, \mathbf{k}_J) = 0$. Otherwise, let $S = [c, d] \cap [a, b]$. Then S is a closed interval.

Let us first assume that S has non empty interior $\mathrm{Int}(S)$. Then $\mathrm{D}(\mathbf{k}_{[c,d] \cap [a,b]})[-1] \simeq \mathbf{k}_{\mathrm{Int}(S)}$, and from proposition 3.3:

$$\mathrm{RHom}(\mathbf{k}_I, \mathbf{k}_J) \simeq \mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{\mathrm{Int}(S)}) \simeq \mathbf{k}[-1].$$

If S has empty interior, since S is a closed interval, there exists $t \in \mathbb{R}$ such that $S = \{t\}$. Therefore, from proposition 3.1, we have $\mathrm{D}(\mathbf{k}_{[c,d] \cap [a,b]})[-1] \simeq \mathbf{k}_{\{t\}}[-1]$. Consequently, we obtain from proposition 3.3:

$$\mathrm{RHom}(\mathbf{k}_I, \mathbf{k}_J) \simeq \mathrm{R}\Gamma(\mathbb{R}; \mathbf{k}_{\{t\}}[-1]) \simeq \mathbf{k}[-1].$$

To sum up, we have proved that:

$$\mathrm{RHom}(\mathbf{k}_I, \mathbf{k}_J) \simeq \begin{cases} \mathbf{k}[-1] & \text{if } [a, b] \cap [c, d] \neq \emptyset \\ 0 & \text{else} \end{cases}.$$

3.2 The functor $- \star \mathbf{K}_\varepsilon$

We start by recalling the following classical lemma about convolution of sheaves:

LEMMA 3.6 (EXERCICE II.20 - [?]) Let $A, B \subset \mathbb{V}$ two closed subsets of the finite dimensional real vector space \mathbb{V} (endowed with the topology inherited from any norm) satisfying:

1. the map $s|_{A \times B} : A \times B \rightarrow \mathbb{V}$ is proper,
2. for any $x \in \mathbb{V}$, $s|_{A \times B}^{-1}(x)$ is contractible.

Then $\mathbf{k}_A \star \mathbf{k}_B \simeq \mathbf{k}_{A+B}$ with $A+B = \{a+b \mid a \in A, b \in B\}$.

Given $\varepsilon \geq 0$ and $I \subset \mathbb{R}$ an interval, we compute the convolution (Definition 2.15) $\mathbf{k}_I \star \mathbf{K}_\varepsilon$ (see Equation 1, Section 2). The case where I is closed is a direct consequence of lemma 3.6. We then deduce the other cases using distinguished triangles.

PROPOSITION 3.7 Let $\varepsilon \geq 0$, and $a \leq b$ in $\mathbb{R} \cup \{\pm\infty\}$, then :

1. $\mathbf{k}_{\mathbb{R}} \star \mathbf{K}_\varepsilon \simeq \mathbf{k}_{\mathbb{R}}$
2. $\mathbf{k}_{[a,b]} \star \mathbf{K}_\varepsilon \simeq \mathbf{k}_{[a-\varepsilon, b+\varepsilon]}$,
3. $\mathbf{k}_{(a,b)} \star \mathbf{K}_\varepsilon \simeq \begin{cases} \mathbf{k}_{(a+\varepsilon, b-\varepsilon)} & \text{if } \varepsilon < \frac{|b-a|}{2} \\ \mathbf{k}_{[\frac{b-a}{2}-\varepsilon, \frac{b-a}{2}+\varepsilon]}[-1] & \text{if } \varepsilon \geq \frac{|b-a|}{2} \end{cases}$
4. $\mathbf{k}_{(a,b]} \star \mathbf{K}_\varepsilon \simeq \mathbf{k}_{(a+\varepsilon, b+\varepsilon]}$,
5. $\mathbf{k}_{[a,b)} \star \mathbf{K}_\varepsilon \simeq \mathbf{k}_{[a-\varepsilon, b-\varepsilon)}$.

PROOF We can obtain the computation for $\mathbf{k}_{(a,b)}$ by using the distinguished triangle $\mathbf{k}_{(a,b)} \longrightarrow \mathbf{k}_{\mathbb{R}} \longrightarrow \mathbf{k}_{\mathbb{R} \setminus (a,b)} \xrightarrow{+1}$, as $\mathbf{k}_{\mathbb{R} \setminus (a,b)}$ is the direct sum of one or two sheaves constant over closed intervals. Similarly for the case of $\mathbf{k}_{[a,b)}$, we can use the distinguished triangles $\mathbf{k}_{[a,b)} \longrightarrow \mathbf{k}_{[a,b]} \longrightarrow \mathbf{k}_{\{b\}} \xrightarrow{+1}$.

□

4 Structure of ε -interleavings

In this section we investigate the structure of ε -interleavings between constructible sheaves over \mathbb{R} . We start by giving explicit conditions on the support and degree of two indecomposables sheaves $\mathbf{k}_I[-i]$ and $\mathbf{k}_J[-j]$ to be ε -interleaved. Then, we introduce for any constructible sheaf F on \mathbb{R} its CLR decomposition (Proposition 4.4), which expresses F as a direct sum of three sheaves F_C , F_L and F_R whose interval decomposition have specific properties. We further show that the CLR decomposition decomposes the notion of interleaving in the following sense: F and G are ε -interleaved if and only if F_C and G_C , F_L and G_L , F_R and G_R are (Theorem 4.8).

4.1 Characterization of ε -interleavings between indecomposable sheaves

For any interval I and real number $\varepsilon \geq 0$, we will write $I^\varepsilon = \bigcup_{x \in I} B(x, \varepsilon)$ where $B(x, \varepsilon)$ is the euclidean closed ball centered at x with radius ε . Moreover if $I = (a, b)$ with $a, b \in \mathbb{R}$, and $\varepsilon < \frac{b-a}{2} = \frac{\text{diam}(I)}{2}$, define $I^{-\varepsilon} = (a + \varepsilon, b - \varepsilon)$. Furthermore, if I is bounded, we write $\text{cent}(I)$ for its center, that is $(b + a)/2$ where a, b are the boundary points of I .

The following proposition describes the condition for sheaves constant on open/closed intervals to be ε -interleaved.

PROPOSITION 4.1 (CLOSED/OPEN) Let S, T be two closed interval, and U, V be two open intervals and $\varepsilon > 0$. Assume that U, V, S, T are non-empty, then:

1. $\mathbf{k}_S \sim_\varepsilon \mathbf{k}_T$ if and only if $S \subset T^\varepsilon$ and $T \subset S^\varepsilon$,
2. $\mathbf{k}_U \sim_\varepsilon \mathbf{k}_V$ if and only if $U \subset V^\varepsilon$ and $V \subset U^\varepsilon$.
3. Assuming that S and U are bounded, $\mathbf{k}_S \sim_\varepsilon \mathbf{k}_U[-1]$ if and only if $\varepsilon \geq \frac{\text{diam}(U)}{2}$ and $S \subset \left[\text{cent}(U) - \left(\varepsilon - \frac{\text{diam}(U)}{2} \right), \text{cent}(U) + \left(\varepsilon - \frac{\text{diam}(U)}{2} \right) \right]$.

PROOF 1. Consider $f : \mathbf{k}_S \star K_\varepsilon \rightarrow \mathbf{k}_T$ and $g : \mathbf{k}_T \star K_\varepsilon \rightarrow \mathbf{k}_S$ the data of an ε -interleaving. Then f and g are in particular not zero since

$$\text{R}\Gamma(\mathbb{R}, \phi_{\mathbf{k}_S, 2\varepsilon}) = \text{R}\Gamma(\mathbb{R}, g) \circ \text{R}\Gamma(\mathbb{R}, f \star K_\varepsilon)$$

is an isomorphism between $\text{R}\Gamma(\mathbb{R}, \mathbf{k}_S \star K_{2\varepsilon})$ and $\text{R}\Gamma(\mathbb{R}, \mathbf{k}_T)$ which are non zero. Remark that $\mathbf{k}_S \star K_\varepsilon \simeq \mathbf{k}_{S^\varepsilon}$ and $\mathbf{k}_T \star K_\varepsilon \simeq \mathbf{k}_{T^\varepsilon}$ by Proposition 3.7. From our computations of morphisms (Proposition 3.5), we have necessarily $S \subset T^\varepsilon$ and $T \subset S^\varepsilon$. Conversely, if $S \subset T^\varepsilon$ and $T \subset S^\varepsilon$, it is easy to build an ε -interleaving.

2. Consider $f : \mathbf{k}_U \star K_\varepsilon \rightarrow \mathbf{k}_V$ and $g : \mathbf{k}_V \star K_\varepsilon \rightarrow \mathbf{k}_U$ the data of an ε -interleaving. For the same reason as above, f and g are not zero. Hence, $f \star K_{-\varepsilon} : \mathbf{k}_U \rightarrow \mathbf{k}_V \star K_{-\varepsilon}$ is not zero. As $\mathbf{k}_V \star K_{-\varepsilon} \simeq \mathbf{k}_{V^\varepsilon}$, by our computations of morphisms (Proposition 3.5) we get that $U \subset V^\varepsilon$. Similarly we have $V \subset U^\varepsilon$.

Conversely if we assume $U \subset V^\varepsilon$ and $V \subset U^\varepsilon$, it is easy to construct an ε -interleaving.

3. Let $f : \mathbf{k}_S \star K_\varepsilon \rightarrow \mathbf{k}_U[-1]$, $g : \mathbf{k}_U[-1] \star K_\varepsilon \rightarrow \mathbf{k}_S$ be the data of an ε -interleaving. For the same reason as above, f and g are not zero. Suppose $\varepsilon < \frac{\text{diam}(U)}{2}$, then Proposition 3.7 implies that $\mathbf{k}_U[-1] \star K_\varepsilon \simeq \mathbf{k}_{U^{-\varepsilon}[-1]}$, hence the fact that g is not zero is absurd.

Therefore we have $\varepsilon \geq \frac{\text{diam}(U)}{2}$, and

$$\mathbf{k}_U[-1] \star K_\varepsilon \simeq \mathbf{k}_{\left[\text{cent}(U) - \left(\varepsilon - \frac{\text{diam}(U)}{2} \right), \text{cent}(U) + \left(\varepsilon - \frac{\text{diam}(U)}{2} \right) \right]}.$$

Hence the existence of g implies that $S \subset \left[\text{cent}(U) - \left(\varepsilon - \frac{\text{diam}(U)}{2}\right), \text{cent}(U) + \left(\varepsilon - \frac{\text{diam}(U)}{2}\right) \right]$.

Also the existence of f implies that $S^\varepsilon \cap U \neq \emptyset$, but this condition is weaker than the previous one.

Conversely, if $\varepsilon \geq \frac{\text{diam}(U)}{2}$ and $S \subset \left[\text{cent}(U) - \left(\varepsilon - \frac{\text{diam}(U)}{2}\right), \text{cent}(U) + \left(\varepsilon - \frac{\text{diam}(U)}{2}\right) \right]$,

we can construct the desired morphisms (using Proposition 3.5) and have to check that their composition (after applying ε convolution to one of the two) is not zero, which can be obtained by taking stalks at any $x \in J$.

PROPOSITION 4.2 (HALF-OPEN) Let $I = [a, b)$ and $J = [c, d)$ with $a, c \in \mathbb{R}$ and $b, d \in \mathbb{R} \cup \{+\infty\}$, and $\varepsilon \geq 0$. Then $\mathbf{k}_I \sim_\varepsilon \mathbf{k}_J \iff |a - c| \leq \varepsilon$ and $|b - d| \leq \varepsilon$.

Similarly for $I = (a, b]$ and $J = (c, d]$, $\mathbf{k}_I \sim_\varepsilon \mathbf{k}_J \iff |a - c| \leq \varepsilon$ and $|b - d| \leq \varepsilon$.

PROOF The proof works exactly the same as the open/closed case, that is Proposition 4.1. □

4.2 CLR Decomposition

In order to define a matching between graded barcodes, we have to distinguish between the topological nature of their support interval as the existence of shifted morphisms between them depends on this nature according to Proposition 3.5.

DEFINITION 4.3 Let I be an interval.

1. I is said to be an interval of type C if there exists $(a, b) \in \mathbb{R}^2$ such that $I = [a, b]$ or $I = (a, b)$.
2. I is said to be an interval of type R if there exists $(a, b) \in \mathbb{R}^2$ such that either $I = [a, b)$, $I = (-\infty, b)$, $I = [a, +\infty)$ or $I = \mathbb{R}$.
3. I is said to be an interval of type L if there exists $(a, b) \in \mathbb{R}^2$ such that $I = (a, b]$, $I = (-\infty, b]$ or $I = (a, +\infty]$.

PROPOSITION 4.4 Let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ then there exists a unique decomposition up to isomorphism $F \simeq F_C \oplus F_R \oplus F_L$ such that :

1. the cohomology objects of F_C are direct sums of constant sheaves over intervals of type C,
2. the cohomology objects of F_R are direct sums of constant sheaves over intervals of type R,
3. the cohomology objects of F_L are direct sums of constant sheaves over intervals of type L.

We will call F_C (resp. F_R, F_L) the central (resp. right, left) part of F , and name this splitting the CLR decomposition of F .

PROOF Observe that the types C,L,R do form a partition of the set of intervals of \mathbb{R} , and apply the decomposition and structure theorems from section 2.14. □

DEFINITION 4.5 Let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$. F is said to be a central sheaf if $F \simeq F_C$. Similarly, F is a left (resp. right) sheaf if $F \simeq F_L$ (resp. $F \simeq F_R$).

We have the following easy properties.

PROPOSITION 4.6 Let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, $\varepsilon \in \mathbb{R}$ and $\alpha \in \{C, L, R\}$. Then F is of type α if and only if $F \star K_\varepsilon$ is of type α .

PROOF It is sufficient to prove the statement for sheaves of the form $\mathbf{k}_I[i]$ for $I \subset \mathbb{R}$ an interval and $i \in \mathbb{Z}$, which is a direct consequence of proposition 3.7. □

PROPOSITION 4.7 Let F and G be two sheaves of type $\alpha \in \{C, L, R\}$. Then any morphism $F \rightarrow G$ that factorizes through a sheaf H as $F \rightarrow H \rightarrow G$, with H of type $\beta \in \{C, L, R\} \setminus \{\alpha\}$ is necessarily zero.

PROOF It is sufficient to prove the statement for $F = \mathbf{k}_I[i]$, $G = \mathbf{k}_J[j]$ and $H = \mathbf{k}_L[l]$, with I and J two intervals of type α , L an interval of type β , and $i, j, l \in \mathbb{Z}$. This is then a direct consequence of our computations of morphisms in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ (proposition 3.5). □

The CLR decomposition is compatible with the relation of being ε -interleaved in the following sense.

THEOREM 4.8 Let $F, G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ and $\varepsilon \geq 0$, then the following holds :

$$F \sim_\varepsilon G \iff \begin{cases} F_C \sim_\varepsilon G_C \\ F_L \sim_\varepsilon G_L \\ F_R \sim_\varepsilon G_R \end{cases} .$$

PROOF The right to left implication is an immediate consequence of the additivity of the convolution functor.

We choose two isomorphisms, that will remain the same through all the proof:

$$F \simeq F_C \oplus F_L \oplus F_R \quad \text{and} \quad G \simeq G_C \oplus G_L \oplus G_R.$$

Then by proposition 4.6, with $\eta \in \mathbb{R}$, applying the functor $- \star K_\eta$ to the above isomorphisms gives us the CLR decompositions of $F \star K_\eta$ and $G \star K_\eta$ in terms of those of F and G . Now let us consider the data of an ε -interleaving between F and G , that is, two morphisms $F \star K_\varepsilon \xrightarrow{f} G$ and $G \star K_\varepsilon \xrightarrow{g} F$ such that $f \star K_\varepsilon \circ g : G \star K_{2\varepsilon} \rightarrow G$ is the smoothing morphism $\phi_{G, 2\varepsilon}$ (see definition 2.17) and similarly $g \star K_\varepsilon \circ f : F \star K_{2\varepsilon} \rightarrow F$ equals $\phi_{F, 2\varepsilon}$. For $\alpha, \beta \in \{C, L, R\}$, we denote by $f_{\alpha, \beta}$ the composition

$$f_{\alpha, \beta} = F_\alpha \star K_\varepsilon \rightarrow (F_C \oplus F_L \oplus F_R) \star K_\varepsilon \simeq F \star K_\varepsilon \xrightarrow{f} G \simeq G_C \oplus G_L \oplus G_R \rightarrow G_\beta.$$

We denote $f_{\alpha, \alpha}$ by f_α , and use similar notations for g and the smoothing morphisms of F and G . By proposition 4.7, we have:

$$\begin{aligned} g \star K_\varepsilon \circ f &= g_C \star K_\varepsilon \circ f_C + g_L \star K_\varepsilon \circ f_L + g_R \star K_\varepsilon \circ f_R, \\ f \star K_\varepsilon \circ g &= f_C \star K_\varepsilon \circ g_C + f_L \star K_\varepsilon \circ g_L + f_R \star K_\varepsilon \circ g_R. \end{aligned}$$

Therefore, for $\alpha \in \{C, L, R\}$, projecting the above equations onto the summands of type α gives:

$$g_\alpha \star K_\varepsilon \circ f_\alpha = \phi_{F_\alpha, 2\varepsilon} \quad \text{and} \quad f_\alpha \star K_\varepsilon \circ g_\alpha = \phi_{G_\alpha, 2\varepsilon},$$

since $(\phi_{F, 2\varepsilon})_\alpha = \phi_{F_\alpha, 2\varepsilon}$ and $(\phi_{G, 2\varepsilon})_\alpha = \phi_{G_\alpha, 2\varepsilon}$. Consequently, we deduce that $F_\alpha \sim_\varepsilon G_\alpha$. □

5 Isometry theorem and graded barcodes

This section presents the proof of the isometry theorem problem associated to the Krull-Schmidt metric category $(D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}}), d_C)$ (see section 2.1). The inequality $d_C \leq d_B$ is an easy consequence of the additivity of the convolution functor.

In order to prove the reverse inequality, we prove that an ε -interleaving between two sheaves induces a ε -matching between their graded-barcodes. To do so, we construct the matching according to the CLR decomposition. We reduce the construction of the matching between the left and right parts to the well-known case of persistence modules with one parameter. To this end, we first prove that interleavings between right (resp. left) parts of two sheaves happen degree-wise at the level of their cohomology objects. This enables us to define functors Ψ_R^j , that send the j -th cohomology of the right part of a sheaf to a one parameter persistence module. We prove that Ψ_R^j are barcode preserving, and send interleavings of sheaves to interleavings of persistence modules.

5.1 The easy inequality

We start by proving the easy direction of the inequality.

LEMMA 5.1 Let F and G two objects of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, then:

$$d_C(F, G) \leq d_B(\mathbb{B}(F), \mathbb{B}(G)).$$

PROOF If $d_B(\mathbb{B}(F), \mathbb{B}(G)) = +\infty$, then the inequality holds. Let us now assume that $d_B(\mathbb{B}(F), \mathbb{B}(G)) < +\infty$. Let $\delta \in \{\varepsilon \geq 0 \mid \text{there exists a } \varepsilon\text{-matching between } \mathbb{B}(F) \text{ and } \mathbb{B}(G)\}$. Then for any $\eta > \delta$ there exists two subsets $\mathcal{X} \subset \mathbb{B}(F)$ and $\mathcal{Y} \subset \mathbb{B}(G)$ and a bijection $\sigma : \mathcal{X} \rightarrow \mathcal{Y}$ such that for all $I^i \in \mathcal{X}$, with $\sigma(I^i) = J^j$ there exists an η -interleaving between $\mathbf{k}_I[-i]$ and $\mathbf{k}_J[-j]$ given by the two morphisms :

$$f_{I^i} : \mathbf{k}_I[-i] \star K_\eta \rightarrow \mathbf{k}_J[-j] \quad \text{and} \quad g_{I^i} : \mathbf{k}_J[-j] \star K_\eta \rightarrow \mathbf{k}_I[-i],$$

and for all $K^k \in \mathbb{B}(F) \setminus \mathcal{X} \sqcup \mathbb{B}(G) \setminus \mathcal{Y}$, $\mathbf{k}_K[-k] \sim_\eta 0$.

Therefore, the morphisms:

$$\begin{aligned} f : F \star K_\eta &\longrightarrow \bigoplus_{I^i \in \mathcal{X}} \mathbf{k}_I[-i] \star K_\eta \xrightarrow{\oplus f_{I^i}} \bigoplus_{J^j \in \mathcal{Y}} \mathbf{k}_J[-j] \longrightarrow G \\ g : G \star K_\eta &\longrightarrow \bigoplus_{J^j \in \mathcal{Y}} \mathbf{k}_J[-j] \star K_\eta \xrightarrow{\oplus g_{I^i}} \bigoplus_{I^i \in \mathcal{X}} \mathbf{k}_I[-i] \longrightarrow F \end{aligned}$$

form an η -interleaving between F and G . Consequently,

$$\begin{aligned} &\{\varepsilon \geq 0 \mid \text{there exists a } \varepsilon\text{-matching between } \mathbb{B}(F) \text{ and } \mathbb{B}(G)\} \\ &\subset \{\varepsilon \geq 0 \mid \text{there exists a } \varepsilon\text{-isomorphism between } F \text{ and } G\} \end{aligned}$$

which proves the lemma by taking the infimum of both sets. □

5.2 The cases $F_R \leftrightarrow G_R$ and $F_L \leftrightarrow G_L$

In this section, we give a description of the ε -interleavings between the right parts of two complexes of sheaves. The proofs and statements for the left parts are exactly the same.

5.2.1 Construction of Ψ_R^j

PROPOSITION 5.2 Let $F, G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ and $\varepsilon \geq 0$ with right parts F_R and G_R . The following holds :

$$F_R \sim_{\varepsilon} G_R \iff \forall j \in \mathbb{Z}, H^j(F_R) \sim_{\varepsilon} H^j(G_R).$$

PROOF The right to left implication is clear, so let us consider an ε -interleaving given by $F_R \star K_{2\varepsilon} \xrightarrow{f} G_R$ and $G_R \star K_{2\varepsilon} \xrightarrow{g} F_R$. Let $j \in \mathbb{Z}$ and pick \mathbf{k}_I a direct summand of $H^j(F_R)$ (I is a half-open interval of the type $[a, b)$). We consider again the composition :

$$\mathbf{k}_I[-j] \star K_{2\varepsilon} \xrightarrow{i_I^{F_R}} F_R \star K_{2\varepsilon} \xrightarrow{f \star K_{2\varepsilon}} G_R \star K_{2\varepsilon} \xrightarrow{g} F_R \xrightarrow{p_I^{F_R}} \mathbf{k}_I[-j]$$

From our computations of derived morphisms (Proposition 3.5), this is equal to :

$$\mathbf{k}_I[-j] \star K_{2\varepsilon} \xrightarrow{i_I^{F_R}} F_R \star K_{2\varepsilon} \xrightarrow{f \star K_{2\varepsilon}} H^j(G_R \star K_{2\varepsilon})[-j] \oplus H^{j+1}(G_R \star K_{2\varepsilon})[-j-1] \xrightarrow{g} F_R \xrightarrow{p_I^{F_R}} \mathbf{k}_I[-j].$$

We obtain using our computations of convolution (Proposition 3.7) that, since G_R has only half-open intervals in the decomposition of its cohomology objects, $H^j(G_R \star K_{2\varepsilon})[-j] \simeq H^j(G_R)[-j] \star K_{2\varepsilon}$ and $H^{j+1}(G_R \star K_{2\varepsilon})[-j-1] \simeq H^{j+1}(G_R)[-j-1] \star K_{2\varepsilon}$.

It follows again from Proposition 3.5 that any morphism of $\mathbf{k}_I[-j] \star K_{2\varepsilon} \rightarrow \mathbf{k}_I[-j]$ that factors through a complex concentrated in degree $j+1$ must be zero.

Finally, the first composition is thus equal to

$$\mathbf{k}_I[-j] \star K_{2\varepsilon} \xrightarrow{i_I^{F_R}} F_R \star K_{2\varepsilon} \xrightarrow{f \star K_{2\varepsilon}} H^j(G_R)[-j] \star K_{2\varepsilon} \xrightarrow{g} F_R \xrightarrow{p_I^{F_R}} \mathbf{k}_I[-j].$$

As this is true for any summand of $H^j(F_R)$ we get that the composition :

$$H^j(F_R)[-j] \star K_{2\varepsilon} \longrightarrow F_R \star K_{2\varepsilon} \xrightarrow{f \star K_{2\varepsilon}} G_R \star K_{2\varepsilon} \xrightarrow{g} F_R \longrightarrow H^j(F_R)[-j]$$

is equal to the composition

$$H^j(F_R)[-j] \star K_{2\varepsilon} \longrightarrow F_R \star K_{2\varepsilon} \longrightarrow H^j(G_R)[-j] \star K_{2\varepsilon} \longrightarrow F_R \longrightarrow H^j(F_R)[-j].$$

This gives the first part of the ε -interleaving. We get the second one by intertwining the roles of F_R and G_R . □

The result above shows that when one wants to understand a morphism between the right parts of two sheaves, it is sufficient to understand it at the level of each of their cohomology objects, degree wise. We will show that the behavior of ε -interleavings between sheaves with cohomologies concentrated in degree $j \in \mathbb{Z}$ decomposing into direct summands of type R , is essentially the same as looking at ε -interleavings in the opposite category of one-parameter persistence modules, which is well understood. We quickly introduce all the necessary definitions and results needed, but we refer to [?] for a detailed exposition about the isometry theorem for one-parameter persistence.

We denote $\text{Pers}_f(\mathbf{k}^{\mathbb{R}})$ the category of pointwise finite dimensional persistence modules over \mathbb{R} , that is, the category of functors $M : (\mathbb{R}, \leq) \rightarrow \text{Mod}_f(\mathbf{k})$ where $\text{Mod}_f(\mathbf{k})$ is the category of finite dimensional vector spaces over the field \mathbf{k} .

There is a notion of ε -interleaving (for $\varepsilon \geq 0$) in this context based on the shift functor $\cdot[\varepsilon]$ defined as $M[\varepsilon](s) = M(s + \varepsilon)$ and $M[\varepsilon](s \leq t) = M(s + \varepsilon \leq t + \varepsilon)$ for $s \leq t$ two real number. There is also a canonical natural transformation $s_\varepsilon^M : M \rightarrow M[\varepsilon]$. We will say that M and N in $\text{Pers}_f(\mathbf{k}^{\mathbb{R}})$ are ε -interleaved if there exists two morphisms $f : N \rightarrow N[\varepsilon]$ and $g : N \rightarrow N[\varepsilon]$ such that $g[\varepsilon] \circ f = s_{2\varepsilon}^M$ and $f[\varepsilon] \circ g = s_{2\varepsilon}^N$.

The pseudo-distance induced on $\text{Pers}_f(\mathbf{k}^{\mathbb{R}})$ by:

$$d_I(M, N) := \inf\{\varepsilon \geq 0 \mid M \text{ and } N \text{ are } \varepsilon\text{-interleaved}\}$$

is called the interleaving distance, and was first introduced in [?]. In this text, for $I \subset \mathbb{R}$ an interval, we will write \mathbf{k}^I the object of $\text{Pers}_f(\mathbf{k}^{\mathbb{R}})$ defined, for $s \leq t$, by:

$$\mathbf{k}^I(s) = \begin{cases} \mathbf{k} & \text{if } s \in I \\ 0 & \text{else} \end{cases} \quad \mathbf{k}^I(s \leq t) = \begin{cases} \text{id}_{\mathbf{k}} & \text{if } s, t \in I \\ 0 & \text{else} \end{cases}.$$

Let D_R^j be the full sub-category of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ of complexes of sheaves F such that $F \simeq F_R$ and $H^i(F) = 0$ for $i \neq j$.

PROPOSITION 5.3 There exists a functor $\Psi_R^j : D_R^j \rightarrow \text{Pers}(\mathbf{k}_{\mathbb{R}})^{op}$ such that :

1. for $F \in D_R^j$ such that $H^j(F) \simeq \bigoplus_{I \in B} \mathbf{k}_I$, we have $\Psi_R^j(F) = \bigoplus_{I \in B} \mathbf{k}^I$,
2. Ψ_R^j is fully faithful,
3. for $\varepsilon \geq 0$ and $F \in D_R^j$, $\Psi_R^j(F \star K_\varepsilon) = \Psi_R^j(F)[\varepsilon]$ and $\Psi_R^j(\phi_{F,\varepsilon}) = s_\varepsilon^{\Psi_R^j(F)}$,
4. Ψ_R^j is isometric with respect to $d_C(\cdot, \cdot)$ and $d_I(\cdot, \cdot)$.

PROOF This is a combination of the computations of morphisms and convolution (Propositions 3.5, 3.7, 4.4), together with the observation that for I, J two intervals of type R and $j \in \mathbb{Z}$, then we have the functorial isomorphisms :

$$\text{Hom}_{D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})}(\mathbf{k}_I[-j], \mathbf{k}_J[-j]) \simeq \text{Hom}_{\text{Mod}(\mathbf{k}_{\mathbb{R}})}(\mathbf{k}_I, \mathbf{k}_J) \simeq \text{Hom}_{\text{Pers}(\mathbf{k}_{\mathbb{R}})}(\mathbf{k}^J, \mathbf{k}^I).$$

5.2.2 Matching of the central parts

THEOREM 5.4 (MATCHING OF RIGHT PARTS) Let $F, G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ be ε -interleaved with respect to the morphisms $F \star K_\varepsilon \xrightarrow{f} G$ and $G \star K_\varepsilon \xrightarrow{g} F$. Let $j \in \mathbb{Z}$. Then there exists an ε -matching $\sigma_R^j : \mathbb{B}^j(F_R) \dashrightarrow \mathbb{B}^j(G_R)$.

PROOF Observe that $\Psi_R^j(F_R)$ (resp. $\Psi_R^j(G_R)$) is a persistence module with the same barcode than $H^j(F_R)$ (resp. $H^j(G_R)$). Also, from proposition 5.3, $\Psi_R^j(F_R)$ and $\Psi_R^j(G_R)$ are ε -interleaved as persistence modules. Hence, we can apply the isometry theorem for pointwise finite dimensional persistence modules [?, Theorem 4.11] to $\Psi_R^j(F_R)$ and $\Psi_R^j(G_R)$ and deduce the existence of a ε -matching of barcodes of persistence modules between $\mathbb{B}^j(F_R)$ and $\mathbb{B}^j(G_R)$. This matching is what we ask for σ_R^j by Proposition 4.2. □

5.3 The case $F_C \leftrightarrow G_C$

In this section, we construct the ε -matching between the central parts of two sheaves, assuming they are ε -interleaved. Using ideas of Bjerkevik [?, Section 4], we introduce a pre-order \leq_α on the set of graded-intervals of type C whose purpose is to prove the existence of the ε -matching using Hall's marriage theorem. To do so, we must prove that given a finite list of interval in the barcode of one of the two sheaves, there exists, at least, the same number of intervals in the barcode of the second sheaf which are at distance less than ε from an interval in the first list.

We will show that ordering the graded-barcodes of the central sheaves according to \leq_α will actually lead to a very nice expression of the interleaving morphisms, allowing us, by a rank argument, to deduce that this condition is satisfied.

5.3.1 Ordering graded-intervals of type C

Recall that we defined a graded interval to be an interval I together with an integer $j \in \mathbb{Z}$. It will be written I^j henceforth. For I of type C such that either $I = [a, b]$ or $I = (a, b)$ with $a, b \in \mathbb{R}$, define $\text{diam}(I) = b - a$ to be its diameter.

DEFINITION 5.5 The relation \leq_α on the set of graded intervals of type C is defined by :

1. For R^i, T^j two closed intervals in degree i and j : $R^i \leq_\alpha T^j \iff i = j$ and $\text{diam}(T) \leq \text{diam}(R)$,
2. for U^i, V^j two open intervals in degree i and j : $U^i \leq_\alpha V^j \iff i = j$ and $\text{diam}(U) \leq \text{diam}(V)$,
3. for R^i a closed interval in degree i , and V^j an open interval in degree j : $R^i \leq_\alpha V^j \iff i = j + 1$.

PROPOSITION 5.6 The relation \leq_α is a partial pre-order over the set of graded intervals of type C, that is, it is reflexive and transitive. Moreover, it is total if restricted to sets of graded intervals containing only, for a given $i \in \mathbb{Z}$, open intervals in degree i and closed intervals in degree $i + 1$.

The following is the analogous result in our setting to [?, Lemma 4.6].

PROPOSITION 5.7 Let I^i, J^j, S^l be three graded intervals of type C and $\varepsilon \geq 0$ such that $I^i \leq_\alpha J^j$ and there exists two non-zero morphisms $\chi : \mathbf{k}_S[-l] \star K_\varepsilon \rightarrow \mathbf{k}_I[-i]$ and $\xi : \mathbf{k}_J[-j] \star K_\varepsilon \rightarrow \mathbf{k}_S[-l]$. Then either $\mathbf{k}_S[-l] \sim_\varepsilon \mathbf{k}_I[-i]$ or $\mathbf{k}_S[-l] \sim_\varepsilon \mathbf{k}_J[-j]$.

PROOF By definition of the pre-order \leq_α , we only have to investigate the three cases of the above definition 5.5 :

1. Let $i \in \mathbb{Z}$ and R, T be two open intervals such that $R^i \leq_\alpha T^i$, that is, $\text{diam}(T) \leq \text{diam}(R)$. Let S^l be a graded interval such that there exists some non-zero χ and ξ . Then S must be a closed interval, and $l = i$. As a consequence, $R \subset S^\varepsilon$ and $S \subset T^\varepsilon$.

Assume that $\mathbf{k}_R[-i] \not\sim_\varepsilon \mathbf{k}_S[-i]$. Then, as $R \subset S^\varepsilon$, $S \not\subset R^\varepsilon$. So either $\min(S) < \min(R) - \varepsilon$, or $\max(S) > \max(R) + \varepsilon$. Assume the latter.

As $S \subset R^\varepsilon$, $\min(S) - \varepsilon < \min(R)$, we get subtracting the first inequality to this one : $\text{diam}(S) + \varepsilon > \text{diam}(R) + \varepsilon$. Hence $S <_\alpha R$. We get the same thing assuming $\min(S) < \min(R) - \varepsilon$.

Moreover, one can prove this way that $\mathbf{k}_T[-i] \not\sim_\varepsilon \mathbf{k}_S[-i]$ implies $S <_\alpha R$.

As we assumed $R^i \leq_\alpha T^i$, one has $\mathbf{k}_S[-l] \sim_\varepsilon \mathbf{k}_I[-i]$ or $\mathbf{k}_S[-l] \sim_\varepsilon \mathbf{k}_J[-j]$.

2. The proof for U^i, V^i where U and V are open intervals is similar.
3. Let R^i a closed interval, V^j an open interval, with $i = j + 1$. Let S^l be a graded interval and ε such that there exists χ and ξ such as in the proposition. Then S must be an open interval and $l = j$. By the existence of χ , we have that $\varepsilon \geq \frac{\text{diam}(U)}{2}$ and $R \subset [\text{cent}(U) - (\varepsilon - \frac{\text{diam}(U)}{2}), \text{cent}(U) + (\varepsilon - \frac{\text{diam}(U)}{2})]$, which, according to our characterization of ε -interleaving between indecomposable sheaves (proposition 4.1), is equivalent to $\mathbf{k}_R[-j-1] \sim_\varepsilon \mathbf{k}_S[-j]$.

5.3.2 Induced matching

We now have the ingredients to prove the theorem. We start by introducing a sign notation. Given two intervals I and J of type C, we define:

$$\delta(I, J) = \begin{cases} 0 & \text{if } I \text{ and } J \text{ are both closed or both open,} \\ 1 & \text{if } I \text{ is open and } J \text{ is closed,} \\ -1 & \text{if } I \text{ is closed and } J \text{ is open.} \end{cases}$$

THEOREM 5.8 (MATCHING OF CENTRAL PARTS) Let F_C and G_C be two central sheaves (definition 4.5), and $\varepsilon \geq 0$ be such that F_C and G_C are ε -interleaved with respect to maps $F_C \star K_\varepsilon \xrightarrow{f} G_C$ and $G_C \star K_\varepsilon \xrightarrow{g} F_C$. Then, there exists a bijection

$$\sigma_C : \mathbb{B}(F_C) \longrightarrow \mathbb{B}(G_C)$$

such that, for $I^j \in \mathbb{B}_C^j$, we have $\sigma_C(I)^{j+\delta(I,J)} \in \mathbb{B}^{j+\delta(I,J)}(G_C)$ and $\mathbf{k}_I \sim_\varepsilon \mathbf{k}_{\sigma_C(I)}[-\delta(I, J)]$.

Our proof will use a generalization of Hall's marriage theorem to the case of countable sets. It was first proved in 1976 by Podewski and Steffens [?]:

THEOREM 5.9 (HALL) Let X and Y be two countable sets, let $\mathcal{P}(Y)$ be the set of subsets of Y and $M : X \rightarrow \mathcal{P}(Y)$. Then the following are equivalent :

1. there exists an injective map $m : X \rightarrow Y$ satisfying $m(x) \in M(x)$ for every $x \in X$;
2. for every finite subset $A \subset X$, $|A| \leq |\cup_{x \in A} M(x)|$. Where $|A|$ is the cardinality of A .

We let F_C and G_C be two central sheaves. We set two isomorphisms :

$$F_C \simeq \bigoplus_{I^j \in \mathbb{B}(F_C)} \mathbf{k}_I[-j] \quad \text{and} \quad G_C \simeq \bigoplus_{I^j \in \mathbb{B}(G_C)} \mathbf{k}_I[-j].$$

For any morphism $f : F_C \rightarrow G_C$, given $I^i \in \mathbb{B}(F_C)$ and $J^j \in \mathbb{B}(G_C)$, we will write :

$$f_{I^i, J^j} = \mathbf{k}_I[-i] \longrightarrow F_C \xrightarrow{f} G_C \longrightarrow \mathbf{k}_J[-j].$$

Similarly for $A \subset \mathbb{B}(F)$, let $f|_A$ be the composition :

$$\bigoplus_{I^i \in A} \mathbf{k}_I[-i] \longrightarrow F_C \xrightarrow{f} G_C.$$

We now assume that F_C and G_C are ε -interleaved with respect to $F_C \star K_\varepsilon \xrightarrow{f} G_C$ and $G_C \star K_\varepsilon \xrightarrow{g} F_C$. For $I^i \in \mathbb{B}(F_C)$ and $J^j \in \mathbb{B}^j(G_C)$, we deduce from our computations of propositions 3.5 and 4.1 that:

$$(f \star K_\varepsilon)_{I^i, J^j} \circ g_{J^j, I^i} \neq 0 \text{ implies either : } \begin{cases} I, J \text{ are closed and } i = j, \\ I \text{ is open, } J \text{ is closed and } j = i + 1, \\ J \text{ is open, } I \text{ is closed and } i = j + 1. \end{cases}$$

PROOF (MATCHING OF CENTRAL PARTS) Our strategy is to adapt Bjerkevik's proof of [?, Theorem 4.2] to our setting. The pre-order \leq_α we have defined has exactly the same properties as the one defined in his proof.

To define σ_C , we will apply Hall's theorem. From the local finiteness properties follows the fact that the graded-barcodes of F_C and G_C are countable. We here consider multi-sets as sets, to make the proof easier to understand. Nevertheless, it would not be difficult to write the proof properly using multi-sets. Let $M : \mathbb{B}_C(F_C) \rightarrow \mathcal{P}(\mathbb{B}_C(G_C))$ defined by :

$$M(I^i) = \{J^j \in \mathbb{B}(G_C) \mid \mathbf{k}_I[-i] \sim_\varepsilon \mathbf{k}_J[-j - \delta(I, J)]\}$$

for $I^i \in \mathbb{B}(F_C)$.

We define the following partitions : $\mathbb{B}(F_C) = \bigsqcup_{i \in \mathbb{Z}} \Sigma_{F_C}^i$ and $\mathbb{B}(G_C) = \bigsqcup_{i \in \mathbb{Z}} \Sigma_{G_C}^i$ where,

$$\Sigma_{F_C}^i = \{J^j \in \mathbb{B}(F_C) \mid J \text{ is open and } j = i \text{ or } J \text{ is closed and } j = i + 1\},$$

$$\Sigma_{G_C}^i = \{J^j \in \mathbb{B}(G_C) \mid J \text{ is open and } j = i \text{ or } J \text{ is closed and } j = i + 1\}.$$

We will define σ_C according to these partitions, that is, we will construct some bijections $\sigma_C^i : \Sigma_{F_C}^i \rightarrow \Sigma_{G_C}^i$ for all $i \in \mathbb{Z}$ and set $\sigma_C = \bigsqcup_{i \in \mathbb{Z}} \sigma_C^i$.

Let $i \in \mathbb{Z}$, A be a finite subset of $\Sigma_{F_C}^i$ and $M(A) = \cup_{I^i \in A} M(I^i)$. To apply Hall's theorem and deduce the existence of σ_C , we need to prove that $|A| \leq |M(A)|$.

By proposition 5.6, \leq_α is a total pre-order on A . Hence, with $r = |A|$, there exists an enumeration $A = \{I_1^{i_1}, \dots, I_r^{i_r}\}$, where $i_l = n$ if I_l is an open interval and $i_l = n + 1$ if I_l is a closed interval, such that for $1 \leq i \leq j \leq r$ we have $I_i \leq_\alpha I_j$.

We have by assumption $g \circ (f \star K_\varepsilon) = \phi_{F, 2\varepsilon}$ (see definition 2.18), also, the additivity of the convolution functor implies the following equality for $I_l^{i_l} \in A$:

$$\phi_{\mathbf{k}_I[-i_l], 2\varepsilon} = \mathbf{k}_I[-i_l] \star K_\varepsilon \longrightarrow F \star K_\varepsilon \xrightarrow{\phi_{F, 2\varepsilon}} F \rightarrow \mathbf{k}_I[-i_l].$$

Therefore :

$$\begin{aligned} \phi_{\mathbf{k}_I[-i_l], 2\varepsilon} &= \sum_{J^j \in \mathbb{B}(G)} g_{J^j, I_l^{i_l}} \circ (f \star K_\varepsilon)_{I_l^{i_l}, J^j} \\ &= \sum_{J^j \in \mathbb{B}^l(G)} g_{J^j, I_l^{i_l}} \circ \left[f_{I_l^{i_l}, J^j} \star K_\varepsilon \right]. \end{aligned}$$

Now observe that if $g_{J^j, I_l^{i_l}} \circ \left[f_{I_l^{i_l}, J^j} \star K_\varepsilon \right] \neq 0$ then $\mathbf{k}_{I_l}[-i_l] \sim_\varepsilon \mathbf{k}_J[-j]$, hence :

$$\phi_{\mathbf{k}_I[-i_l], 2\varepsilon} = \sum_{J^j \in M(A)} g_{J^j, I_l^{i_l}} \circ (f \star K_\varepsilon)_{I_l^{i_l}, J^j}$$

Similarly for $I_m \neq I_{m'}$ in A ,

$$0 = \sum_{J^j \in \mathbb{B}^l(G)} g_{J^j, I_m^{i_m}} \circ \left[f_{I_{m'}, J^j}^{i_{m'}} \star K_\varepsilon \right].$$

Hence if $m < m'$ and $g_{J^j, I_m^{i_m}} \circ \left[f_{I_{m'}, J^j}^{i_{m'}} \star K_\varepsilon \right] \neq 0$, then $\mathbf{k}_J[-j]$ is ε -interleaved with either $\mathbf{k}_{I_m}[-m]$ or $\mathbf{k}_{I_{m'}}[-m']$. Therefore :

$$0 = \sum_{J^j \in M(A)} g_{J^j, I_m^{i_m}} \circ \left[f_{I_{m'}, J^j}^{i_{m'}} \star K_\varepsilon \right].$$

For $m > m'$, we can't say anything about the value of $\sum_{J^j \in M(A)} g_{J^j, I_m^{i_m}} \circ \left[f_{I_{m'}, J^j}^{i_{m'}} \star K_\varepsilon \right]$.

Writing those equalities in matrix form, we get :

$$\begin{pmatrix} g_{J^1, I_1^{i_1}} & \cdots & g_{J^1, I_r^{i_r}} \\ \vdots & \ddots & \vdots \\ g_{J^s, I_1^{i_1}} & \cdots & g_{J^s, I_r^{i_r}} \end{pmatrix} \begin{pmatrix} f_{I_1^{i_1}, J^1} \star K_\varepsilon & \cdots & f_{I_r^{i_r}, J^1} \star K_\varepsilon \\ \vdots & \ddots & \vdots \\ f_{I_1^{i_1}, J^s} \star K_\varepsilon & \cdots & f_{I_r^{i_r}, J^s} \star K_\varepsilon \end{pmatrix} = \begin{pmatrix} \phi_{I_1^{i_1}, 2\varepsilon} & ? & ? & ? \\ 0 & \phi_{I_2^{i_2}, 2\varepsilon} & ? & ? \\ \vdots & \vdots & \ddots & ? \\ 0 & 0 & \cdots & \phi_{I_r^{i_r}, 2\varepsilon} \end{pmatrix}$$

Now recall that $\mathrm{R}\Gamma(\mathbb{R}, -)$ is an additive functor. Hence, applying $\mathrm{R}\Gamma(\mathbb{R}, -)$ to the above equality, we get :

$$\begin{aligned} & \begin{pmatrix} \mathrm{R}\Gamma(\mathbb{R}, g_{J^1, I_1^{i_1}}) & \cdots & \mathrm{R}\Gamma(\mathbb{R}, g_{J^1, I_r^{i_r}}) \\ \vdots & \ddots & \vdots \\ \mathrm{R}\Gamma(\mathbb{R}, g_{J^s, I_1^{i_1}}) & \cdots & \mathrm{R}\Gamma(\mathbb{R}, g_{J^s, I_r^{i_r}}) \end{pmatrix} \begin{pmatrix} \mathrm{R}\Gamma(\mathbb{R}, f_{I_1^{i_1}, J^1} \star K_\varepsilon) & \cdots & \mathrm{R}\Gamma(\mathbb{R}, f_{I_r^{i_r}, J^1} \star K_\varepsilon) \\ \vdots & \ddots & \vdots \\ \mathrm{R}\Gamma(\mathbb{R}, f_{I_1^{i_1}, J^s} \star K_\varepsilon) & \cdots & \mathrm{R}\Gamma(\mathbb{R}, f_{I_r^{i_r}, J^s} \star K_\varepsilon) \end{pmatrix} \\ &= \begin{pmatrix} \mathrm{R}\Gamma(\mathbb{R}, \phi_{I_1^{i_1}, 2\varepsilon}) & ? & ? & ? \\ 0 & \mathrm{R}\Gamma(\mathbb{R}, \phi_{I_2^{i_2}, 2\varepsilon}) & ? & ? \\ \vdots & \vdots & \ddots & ? \\ 0 & 0 & \cdots & \mathrm{R}\Gamma(\mathbb{R}, \phi_{I_r^{i_r}, 2\varepsilon}) \end{pmatrix} \\ &= \begin{pmatrix} 1 & ? & ? & ? \\ 0 & 1 & ? & ? \\ \vdots & \vdots & \ddots & ? \\ 0 & 0 & \cdots & 1 \end{pmatrix}. \end{aligned}$$

Each entry in those matrices is uniquely characterized by one scalar. Hence, we can consider their rank. The left hand side has rank at most equal to the minimum of r and s , in particular it is less or equal to $|M(A)|$. The right-hand side has rank $r = |A|$. Therefore we obtain the inequality we wanted. \square

5.4 Isometry theorem

In this section, we put together the results proved before to prove that the convolution distance between two sheaves is exactly the same as the bottleneck distance between their graded-barcodes.

THEOREM 5.10 (ISOMETRY) Let F, G be two objects of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, then :

$$d_C(F, G) = d_B(\mathbb{B}(F), \mathbb{B}(G)).$$

PROOF By Lemma 5.1, there only remains to prove that $d_C(F, G) \geq d_B(\mathbb{B}(F), \mathbb{B}(G))$, or equivalently, that any ε -interleaving between F and G induces an ε -matching between $\mathbb{B}(F)$ and $\mathbb{B}(G)$.

According to sections 5.3 and 5.4, this interleaving induces a ε -matching between the central, left and right parts of F and G , which proves the theorem.

6 Applications

In this section, we expose some corollaries of the isometry theorem. We start with some explicit computations on an example, showing the fundamentally derived nature of our graded-bottleneck distance. Then, we prove that d_C is closed, that is, two sheaves are ε -close if and only if they are ε -interleaved, which in particular implies that d_C induces a metric on the isomorphism classes of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$. We then provide a counter-example of two non constructible sheaves being at convolution distance zero, but which are not isomorphic. These results answer an open question asked by Kashiwara-Schapira in [?] in the one dimensional case. The fact that d_C is closed allows us to consider the set of isomorphism classes of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ as a topological metric space. We prove that it is locally path-connected and give a characterization of its connected components.

6.1 Example : projection from the circle

We aim here to explain and compute an explicit example that was pointed to us by Justin Curry. It consists of two simple maps from the euclidean circle to the real line. Understanding this example has been at the origin of our work. It is simple yet general enough to exhibit the phenomenons and issues that can happen with the matchings of graded barcodes.


Let $\mathbb{S}^1 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$ be the one dimensional circle seen as a sub-manifold in \mathbb{R}^2 . Let $f : \mathbb{S}^1 \rightarrow \mathbb{R}$ be the first coordinate projection and $g : \mathbb{S}^1 \rightarrow \mathbb{R}$ be the constant map with value zero. Let $F = Rf_*\mathbf{k}_{\mathbb{S}^1}$ and $G = Rg_*\mathbf{k}_{\mathbb{S}^1}$. Since $\|f - g\| = 1$, the stability theorem by Kashiwara and Schapira [?, theorem 2.7] implies :

$$d_C(F, G) \leq 1.$$

The CLR decomposition (Definition 4.3) of this two complexes of sheaves is easy to compute (and depicted in the figure below).

PROPOSITION 6.1 The complexes F and G have non-zero cohomology spaces at most in degree 0 and 1. Moreover :

1. $H^0(F) \simeq \mathbf{k}_{(-1,1)} \oplus \mathbf{k}_{[-1,1]}$ and $H^1(F) \simeq 0$
2. $H^0(G) \simeq \mathbf{k}_{\{0\}}$ and $H^1(G) \simeq \mathbf{k}_{\{0\}}$



Hence, F and G are central sheaves and $\mathbb{B}(F) = \{[-1, 1]^0, (-1, 1)^0\}$, $\mathbb{B}(G) = \{\{0\}^0, \{0\}^1\}$. Even in this simple example, there could be *no* ε -matching between the graded-barcodes if one was working in the ordinary graded category. Indeed, $d_C(\mathbf{k}_{\{0\}}[-1], 0) = +\infty$. Hence this will not agree both with Kashiwara-Schapira (naturally derived) convolution distance and the intuition from persistence.

However, using our derived notion of interleavings and matching distance we get the expected answer and in fact prove that in this case the bound given by the L_∞ -norm between the function is optimal.

Indeed, let σ be the matching defined by :

$$\sigma([-1, 1]^0) = \{0\}^0 \quad \text{and} \quad \sigma((-1, 1)^0) = \{0\}^1.$$

Then we claim that σ is a 1-matching between $\mathbb{B}(F)$ and $\mathbb{B}(G)$. Moreover, since the convolution distances between any pair of graded intervals is at least 1, there can not exist an ε -matching between $\mathbb{B}(F)$ and $\mathbb{B}(G)$ for $0 \leq \varepsilon < 1$. Hence we have $F \sim_1 G$ and further

PROPOSITION 6.2 The convolution distance of $F = Rf_*\mathbf{k}_{S^1}$ and $G = Rg_*\mathbf{k}_{S^1}$ is

$$d_C(Rf_*\mathbf{k}_{S^1}, Rg_*\mathbf{k}_{S^1}) = 1.$$

6.2 About the closedness of d_C

In this section we apply our isometry Theorem 5.10 to answer an open question of Kashiwara-Schapira on the closedness of the convolution distance (see Remark 2.3 of [?]) in the one dimensional case. More precisely, we show that the convolution distance is closed between constructible sheaves over \mathbb{R} . We also provide a counter-example to this statement without constructibility assumption.

THEOREM 6.3 The convolution distance is closed on $D_{\mathbb{R}^c}^b(\mathbf{k}_{\mathbb{R}})$. That is, for $F, G \in D_{\mathbb{R}^c}^b(\mathbf{k}_{\mathbb{R}})$ and $\varepsilon \geq 0$: $d_C(F, G) \leq \varepsilon \iff F \sim_\varepsilon G$.

We start with the following easy lemma, whose proof is left to the reader.

LEMMA 6.4 Let I^i, J^j two graded intervals (possibly empty, we set $\mathbf{k}_\emptyset = 0$) and $\varepsilon \geq 0$. Then :

$$d_C(\mathbf{k}_I[-i], \mathbf{k}_J[-j]) \leq \varepsilon \iff \mathbf{k}_I[-i] \sim_\varepsilon \mathbf{k}_J[-j]$$

PROOF (OF THE THEOREM) Suppose $d_C(F, G) \leq \varepsilon$. Then by definition there exists a decreasing sequence (ε_n) such that $\varepsilon_n \rightarrow \varepsilon$ when n goes to infinity and for every $n \in \mathbb{N}$, $F \sim_{\varepsilon_n} G$. For simplicity of the proof, we will assume the graded-barcodes of F and G to be finite, but the proof generalizes to the locally finite case. Then by applying the isometry theorem, for $n \geq 0$, there exists a ε_n matching $\sigma_n : \mathbb{B}(F) \rightarrow \mathbb{B}(G)$.

Now by finiteness of the graded-barcodes, the set of matchings between $\mathbb{B}(F)$ and $\mathbb{B}(G)$ is finite. Hence, we can extract from (σ_n) a constant sequence, say $(\sigma_{\varphi(n)})$. Applying lemma 4.1 and making n going to infinity, we see that $\sigma := \sigma_{\varphi(0)}$ is an ε -matching between $\mathbb{B}(F)$ and $\mathbb{B}(G)$.

REMARK 6.5 One must observe that in the case of persistence modules, the interleaving distance is *not* closed. There exists some ephemeral modules at distance 0 from 0 : consider the one parameter persistence module $\mathbf{k}^{\{0\}}$ (keeping notations of section 5.2). To avoid this issue, Chazal, Crawley-Boevey and de Silva introduced the observable category of persistence modules $\text{Obs}(\text{Pers}(\mathbf{k}_{\mathbb{R}}))$ in [?]. It is defined as the quotient category of $\text{Pers}(\mathbf{k}_{\mathbb{R}})$ by the full sub-category of ephemeral persistent modules, which has objects $M \in \text{Pers}(\mathbb{R})$ such that $M(s < t) = 0$ for every $s < t \in \mathbb{R}$. By construction, the interleaving distance on $\text{Pers}(\mathbf{k}_{\mathbb{R}})$ induces a closed metric on $\text{Obs}(\text{Pers}(\mathbf{k}_{\mathbb{R}}))$. Note that this construction has since been generalized by the first author in [?].

COROLLARY 6.6 The functors $\Psi_R^j : D_R^j \rightarrow \text{Pers}(\mathbb{R})$ (see proposition 5.3) induces an isometric equivalence of categories between D_R^j and $\text{Obs}(\text{Pers}(\mathbb{R}))^{op}$.

We now explicit a counter-example to the closedness of d_C without constructibility assumptions. More precisely, we will construct two sheaves $F, G \in D^b(\mathbf{k}_{\mathbb{R}})$ such that $d_C(F, G) = 0$ but $F \not\cong G$. We consider the sets $X = \mathbb{Q} \cap [0, 1]$ and $Y = \sqrt{2}\mathbb{Q} \cap [0, 1] = \{\sqrt{2}q \mid q \in \mathbb{Q}\} \cap [0, 1]$.

PROPOSITION 6.7 There exists a sequence of functions $(r_n)_{n \in \mathbb{Z}_{>0}}$ from X to Y satisfying:

1. for any $n \in \mathbb{Z}_{>0}$, $r_n : X \rightarrow Y$ is bijective,
2. $\sup_{x \in X} |r_n(x) - x| \xrightarrow{n \rightarrow +\infty} 0$.

PROOF Let $n \in \mathbb{Z}_{>0}$. We define r_n piecewise on $[0, 1 - \frac{1}{n}] \cap \mathbb{Q}$ and $]1 - \frac{1}{n}, 1] \cap \mathbb{Q}$. For $q \in [0, 1 - \frac{1}{n}] \cap \mathbb{Q}$, we set $r_n(q) = \frac{\sqrt{2}}{\lceil \sqrt{2} \rceil_{10}(n)} q$, with $\lceil \sqrt{2} \rceil_{10}(n) = \frac{\lceil 10^n \sqrt{2} \rceil}{10^n}$ the n -th ceil decimal approximation of $\sqrt{2}$. Then $r_n|_{[0, 1 - \frac{1}{n}] \cap \mathbb{Q}}$ is injective, and

$$\begin{aligned} r_n \left([0, 1 - \frac{1}{n}] \cap \mathbb{Q} \right) &= \left\{ \sqrt{2}q \mid q \in \left[0, \frac{1 - 1/n}{\lceil \sqrt{2} \rceil_{10}(n)} \right] \cap \mathbb{Q} \right\} \\ &= Y \cap \left[0, \sqrt{2} \frac{1 - 1/n}{\lceil \sqrt{2} \rceil_{10}(n)} \right] \\ &\subsetneq Y. \end{aligned}$$


Now, since $]1 - \frac{1}{n}, 1] \cap \mathbb{Q}$ and $Y \setminus r_n([0, 1 - \frac{1}{n}] \cap \mathbb{Q})$ are both infinite subsets of \mathbb{Q} , there exists a bijection

$$\varphi_n : \left] 1 - \frac{1}{n}, 1 \right] \cap \mathbb{Q} \xrightarrow{\sim} Y \setminus r_n \left([0, 1 - \frac{1}{n}] \cap \mathbb{Q} \right).$$

We define $r_n|_{[1-\frac{1}{n}, 1] \cap \mathbb{Q}} = \varphi_n$. Then $r_n|_{[1-\frac{1}{n}, 1] \cap \mathbb{Q}}$ is injective and

$$r_n([1 - \frac{1}{n}, 1] \cap \mathbb{Q}) = Y \setminus r_n \left([0, 1 - \frac{1}{n}] \cap \mathbb{Q} \right).$$

Finally, r_n is indeed a bijective function from X to $Y = \sqrt{2}\mathbb{Q} \cap [0, 1]$.



Graphical representation of r_n

Observe that :

$$\sup_{x \in X} |r_n(x) - x| = \max \left(\sup_{x \in X \cap [0, 1 - \frac{1}{n}]} |r_n(x) - x|, \sup_{x \in X \cap [1 - \frac{1}{n}, 1]} |r_n(x) - x| \right).$$

The first term of the maximum is worth $\left(\frac{\sqrt{2}}{\lceil \sqrt{2} \rceil_{10}(n)} - 1 \right) (1 - 1/n)$, and the second term is bounded from above by the diameter of the interval $[\sqrt{2} \frac{1-1/n}{\lceil \sqrt{2} \rceil_{10}(n)}, 1]$ which is worth $1 - \sqrt{2} \frac{1-1/n}{\lceil \sqrt{2} \rceil_{10}(n)}$. Since both of these terms go to 0 as n goes to infinity, we deduce the desired property :

$$\sup_{x \in X} |r_n(x) - x| \xrightarrow{n \rightarrow +\infty} 0.$$

PROPOSITION 6.8 Let $(F_i)_{i \in I}$ and $(G_j)_{j \in J}$ two families of objects of $D^b(\mathbf{k}_{\mathbb{V}})$. Assume that there exists a bijective function $\sigma : I \rightarrow J$, and $\varepsilon \geq 0$ such that for all $i \in I$, $d_C(F_i, G_{\sigma(i)}) \leq \varepsilon$. Then :

$$d_C \left(\bigoplus_{i \in I} F_i, \bigoplus_{j \in J} G_j \right) \leq \varepsilon.$$

PROOF Let $\varepsilon' > \varepsilon$ and $i \in I$. Then by assumptions, there exists ε' -interleaving morphisms between F_i and $G_{\sigma(i)}$, $\varphi_i : F_i \star K_{\varepsilon'} \rightarrow G_{\sigma(i)}$ and $\psi_i : G_{\sigma(i)} \star K_{\varepsilon'} \rightarrow F_i$. Since $-\star K_{\varepsilon'}$ is a left-adjoint functor, it commutes with arbitrary colimits. Therefore, by taking direct sums of the previous ε' -interleaving morphisms, we get ε' -interleaving morphisms between $\bigoplus_{i \in I} F_i$ and $\bigoplus_{i \in I} G_{\sigma(i)} \simeq \bigoplus_{j \in J} G_j$, which proves the result. □

Let $F = \bigoplus_{x \in X} \mathbf{k}_{\{x\}}$ and $G = \bigoplus_{y \in Y} \mathbf{k}_{\{y\}}$.

PROPOSITION 6.9 F is not isomorphic to G and $d_C(F, G) = 0$.

PROOF F and G cannot be isomorphic since $F_1 \simeq \mathbf{k}$ and $G_1 \simeq 0$.

Let $r_n : X \rightarrow Y$ be as in proposition 6.7. Using proposition 6.8, and the fact that, for $x, y \in \mathbb{R}$, $d_C(\mathbf{k}_{\{x\}}, \mathbf{k}_{\{y\}}) = |x - y|$, we obtain that, for any $n \in \mathbb{Z}_{>0}$, :

$$d_C(F, G) \leq \sup_{x \in X} |r_n(x) - x|.$$

Taking the limit as n goes to infinity, we deduce that $d_C(F, G) = 0$. □

6.3 Description of the connected components of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$

In this section, we study the connected components of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ with respect to the metric. In order to make sense of this, we introduce the *small* category Barcode which is a combinatorial description of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ and is equivalent to $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$. The category Barcode is shown to be skeletal (any two isomorphic objects are equal), and is equipped with the graded bottleneck distance. From Barcode we thus obtain an extended metric space, which will be proven to be locally path-connected, see Theorem 6.12. To do so, we first prove an interpolation lemma in the same fashion as Chazal et al. [?, Theorem 3.5], which stands that if two sheaves are ε -interleaved, there exists a 1-lipschitz path in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ between them.

LEMMA 6.10 (INTERPOLATION) Let $F, G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ be such that $F \sim_{\varepsilon} G$ for some $\varepsilon \geq 0$. Then there exists a family of sheaves $(U_t)_{t \in [0, \varepsilon]}$ in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ such that :

1. $U_0 = F$ and $U_{\varepsilon} = G$.
2. For $t \in [0, \varepsilon]$, $d_c(F, U_t) \leq t$ and $d_c(G, U_t) \leq t - \varepsilon$.
3. For $(t, t') \in [0, \varepsilon]^2$, $d_c(U_t, U_{t'}) \leq |t - t'|$.

PROOF Let $F \star K_{\varepsilon} \xrightarrow{\varphi} G$ and $G \star K_{\varepsilon} \xrightarrow{\psi} F$ be the interleaving morphisms between F and G .

We start by constructing U_t for $t \in [0, \frac{\varepsilon}{2}]$. The interleaving morphism and the canonical maps in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ give (by Proposition 2.16 and (2)) the following diagram \mathbb{D}_t :

$$\begin{array}{ccc} G \star K_{t-\varepsilon} & & F \star K_{-t} \\ \downarrow \phi_{G, 2t \star K_{t-\varepsilon}} & \begin{array}{c} \swarrow \psi \star K_{t-\varepsilon} \\ \searrow \phi \star K_{-t} \end{array} & \downarrow \phi_{F, 2\varepsilon - 2t \star K_{-t}} \\ G \star K_{-t-\varepsilon} & & F \star K_{t-2\varepsilon} \end{array} .$$

Taking resolutions in $\text{Mod}(\mathbf{k}_{\mathbb{R}})$, one can assume this diagram is actually given by a diagram still denoted \mathbb{D}_t in $C(\text{Mod}(\mathbf{k}_{\mathbb{R}}))$ which we assume from now on. One can note that this diagram defines the two maps $\theta_t, \tilde{\phi}_t : (G \star K_{t-\varepsilon}) \oplus (F \star K_{-t}) \rightarrow (G \star K_{-t-\varepsilon}) \oplus (F \star K_{t-2\varepsilon})$ given by $(x, y) \xrightarrow{\theta_t} (\varphi \star K_{-t}(y), \psi \star K_{t-\varepsilon}(x))$ and $(x, y) \xrightarrow{\tilde{\phi}_t} (\phi_{G, 2t \star K_{t-\varepsilon}}(x), \phi_{F, 2\varepsilon - 2t \star K_{-t}}(y))$. The limit $\varprojlim \mathbb{D}_t$ of the diagram is precisely (isomorphic to) the equalizer of the two maps and thus to the kernel $\ker(\theta_t - \tilde{\phi}_t)$ of their difference.

Since we are dealing with a diagram in $C(\text{Mod}(\mathbf{k}_{\mathbb{R}}))$ that we wish to see in the derived category we essentially only need to replace the limit by its homotopy limit. Namely,

we define $\tilde{U}_t := \text{ho} \varprojlim \mathbb{D}_t$ to be the homotopy limit in (the model category of sheaves [?]) $C(\text{Mod}(\mathbf{k}_{\mathbb{R}}))$ of the diagram \mathbb{D}_t . For the reader who wish to avoid the use of the technique of homotopy limits in model category (see [?]), in view of the above identification, an explicit model for this homotopy limit is given by the cocone of the map $(G \star K_{t-\varepsilon}) \oplus (F \star K_{-t}) \xrightarrow{\theta_t - \tilde{\phi}_t} (G \star K_{-t-\varepsilon}) \oplus (F \star K_{t-2\varepsilon})$ and hence passes to $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$.

We need to prove that \tilde{U}_t is t -interleaved with F . Note that by definition of an homotopy limit we have a canonical map $\tilde{U}_t \rightarrow (G \star K_{t-\varepsilon}) \oplus (F \star K_{-t})$, simply given by the obvious map on the cocone above. Hence we have a morphism to either factor, in particular we have $\tilde{U}_t \xrightarrow{f} F \star K_{-t}$ and hence (by proposition 2.16) a map

$$\tilde{U}_t \star K_t \xrightarrow{f} F. \quad (3)$$

We now need to define a map $g : F \star K_t \rightarrow \tilde{U}_t$. By definition of an homotopy limit, we have a canonical factorization $\varprojlim \mathbb{D}_t \rightarrow \tilde{U}_t \rightarrow (G \star K_{t-\varepsilon}) \oplus (F \star K_{-t})$ of the canonical map defined by the limit in $C(\text{Mod}(\mathbf{k}_{\mathbb{R}}))$. At the explicit cocone level this is just given by the canonical map of the limit on its first summand $(G \star K_{t-\varepsilon}) \oplus (F \star K_{-t})$

The interleaving map $\varphi : F \star K_{\varepsilon} \rightarrow G$ induces the map

$$F \star K_t \xrightarrow{(\varphi \star K_{t-\varepsilon}, \phi_{F,2t \star K_t})} (G \star K_{t-\varepsilon}) \oplus (F \star K_{-t}) \quad (4)$$

which makes the following diagram

$$\begin{array}{ccc} & F \star K_t & \\ \varphi \star K_{t-\varepsilon} \swarrow & & \searrow \phi_{F,2t \star K_t} \\ G \star K_{t-\varepsilon} & & F \star K_{-t} \\ \downarrow \phi_{G,2t \star K_{t-\varepsilon}} & \psi \star K_{t-\varepsilon} \swarrow & \searrow \varphi \star K_{-t} \\ G \star K_{-t-\varepsilon} & & F \star K_{t-2\varepsilon} \\ & \downarrow \phi_{F,2\varepsilon-2t \star K_{-t}} & \end{array}$$

commutative since φ, ψ defines a ε -interleaving. This implies that the map (4) factors through $\varprojlim \mathbb{D}_t$ and hence we get a map $g : F \star K_t \rightarrow \varprojlim \mathbb{D}_t \rightarrow \tilde{U}_t$ in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$.

For $t \in]\frac{\varepsilon}{2}, \varepsilon]$, we construct U_t in a similar fashion by intertwining the roles of F and G in the diagram $C(\text{Mod}(\mathbf{k}_{\mathbb{R}}))$.

Let $\Delta_{\varepsilon} = \{(x, y) \in \mathbb{R}^2 \mid 0 \leq y - x \leq \varepsilon\}$ be equipped with the standard product order of $\mathbb{R}^2 : (x, y) \leq (x', y') \iff x \leq x' \text{ and } y \leq y'$. Observe that the mapping :

$$\Delta_{\varepsilon} \ni (x, y) \rightsquigarrow U_{y-x} \star K_{-x-y}$$

induces a well defined functor $(\Delta_{\varepsilon}, \leq) \longrightarrow D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ whose restriction to the poset $\{(x, y) \in \mathbb{R}^2 \mid y - x = t\}$ is the functor : $(x, y) \longrightarrow U_t \star K_{-x-y}$ with internal maps given by the natural morphisms $(\phi_{U_t, \varepsilon})$. Hence, for $\varepsilon \geq t, t' \geq 0$, U_t and $U_{t'}$ are $|t - t'|$ interleaved. □

We will now define the category Barcode we mentioned earlier.

Let $\text{Int}(\mathbb{R})$ be the set of intervals of \mathbb{R} and p_1, p_2 be the two first coordinate projections of $\text{Int}(\mathbb{R}) \times \mathbb{Z} \times \mathbb{Z}_{\geq 0}$. Let \mathbb{B} be a subset of $\text{Int}(\mathbb{R}) \times \mathbb{Z} \times \mathbb{Z}_{\geq 0}$. Then \mathbb{B} is said to be

— *locally finite* if $p_1(\mathbb{B}) \cap K$ is finite for all compact subsets of \mathbb{R} ;

- *bounded* if $p_2(\mathbb{B}) \subset \mathbb{Z}$ is bounded;
- *well-defined* if the fibers of the projection (p_1, p_2) have cardinality at most 1.

In a triple $(I, j, n) \in \mathbb{B}$, the first integer will stand for the degree on which the interval I is seen and the second non-negative integer n stands for its multiplicity.

DEFINITION 6.11 The category *Barcode* has objects the set

$$\text{Obj}(\text{Barcode}) = \{\mathbb{B} \subset \text{Int}(\mathbb{R}) \times \mathbb{Z} \times \mathbb{Z}_{\geq 0} \mid \mathbb{B} \text{ is bounded, locally finite and well-defined}\}.$$

For any \mathbb{B} and $\mathbb{B}' \in \text{Barcode}$, the set of their morphisms is

$$\text{Hom}_{\text{Barcode}}(\mathbb{B}, \mathbb{B}') = \prod_{\substack{(I, j, n) \in \mathbb{B} \\ (I', j', n') \in \mathbb{B}'}} \text{Hom}_{D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})}(\mathbf{k}_I^n[-j], \mathbf{k}_{I'}^{n'}[-j']).$$

We define the composition in *Barcode* so that the mapping :

$$\iota : \text{Obj}(\text{Barcode}) \ni \mathbb{B} \mapsto \bigoplus_{(I, j, n) \in \mathbb{B}} \mathbf{k}_I^n[-j] \in \text{Obj}(D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}}))$$

becomes a fully faithful functor :

$$\iota : \text{Barcode} \longrightarrow D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}}).$$

Note that this is possible only because the objects of *Barcode* are locally finite. Theorems 2.12 and 2.9 assert that ι is essentially surjective, therefore is an equivalence. We also deduce from these theorems that *Barcode* is a skeletal category: it satisfies for any $\mathbb{B}, \mathbb{B}' \in \text{Barcode}$,

$$\mathbb{B} \simeq \mathbb{B}' \text{ if and only if } \mathbb{B} = \mathbb{B}'.$$

The notion of equality is well-defined here since $\text{Obj}(\text{Barcode})$ is a set. Therefore ι identifies its image as a skeleton of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$, a full-subcategory which is dense and skeletal.

Moreover, Theorems 2.12 and 2.9 allows us to equip the set $\text{Obj}(\text{Barcode})$ with the graded-bottleneck distance (definition 2.5). The derived isometry theorem 5.10 implies that, for any $\mathbb{B}, \mathbb{B}' \in \text{Barcode}$, one has :

$$d_C(\iota(\mathbb{B}), \iota(\mathbb{B}')) = d_B(\mathbb{B}, \mathbb{B}').$$

THEOREM 6.12 The following assertions hold :

1. $(\text{Obj}(\text{Barcode}), d_B)$ is an extended metric space,
2. $(\text{Obj}(\text{Barcode}), d_B)$ is locally path-connected.

PROOF 1. The fact that d_B is a pseudo-extended metric is inherited from the properties of d_C (proposition 2.21) by the derived isometry theorem. Moreover, if $d_B(\mathbb{B}, \mathbb{B}') = 0$ then $\mathbb{B} = \mathbb{B}'$ by theorem 6.3.

2. We will prove that open balls are path-connected, that is, any two barcodes at finite distance can be connected by a continuous path. Let \mathbb{B}_0 and \mathbb{B}_ε in *Barcode* such that $d_B(\mathbb{B}_0, \mathbb{B}_\varepsilon) = \varepsilon$. According to the interpolation lemma 6.10, there exists a family of objects $(F_t)_{t \in [0, \varepsilon]}$ of $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ such that $F_0 = \iota(\mathbb{B}_0)$, $F_\varepsilon = \iota(\mathbb{B}_\varepsilon)$, and for any $t, t' \in [0, \varepsilon]$, $d_C(F_t, F_{t'}) \leq |t - t'|$. Given $t \in [0, \varepsilon]$, define \mathbb{B}_t to be the graded-barcode of F_t . Then, it is clear thanks to the derived isometry theorem that $(t \mapsto \mathbb{B}_t)$ defines a 1-lipschitz path between \mathbb{B}_0 and \mathbb{B}_ε .

REMARK 6.13 One shall be aware that it is not sufficient for being in the same connected component to have graded-barcodes of the same “type”, as one can see in the following counter-example. Define :

$$F = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} \mathbf{k}_{\left(\frac{n(n+1)}{2}, \frac{(n+1)(n+2)}{2}\right)} \quad \text{and} \quad G = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} \mathbf{k}_{\left(\frac{(n+1)^2}{2}-1, \frac{(n+1)^2}{2}+1\right)}$$

Then $d_C(F, G) = +\infty$, hence F and G do not belong to the same connected component, whereas one might think that the obvious matching between their barcodes leads to a continuous path between the two sheaves.

6.4 Algorithmic remarks on computing one best matching

The formulation of the convolution distance as a matching distance we obtained in Section 5 turns the computation of an algebraic problem into minimizing the cost of a matching, which is of combinatorial nature. This is in fact a variant of a very classical problem of linear programming, for which there exists an abundant literature that can be solved in polynomial time using the *Hungarian algorithm* [?]. Hence, distances in $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{R}})$ can be implemented in a computer and computed.