

# MAHARAM EXTENSION AND STATIONARY STABLE PROCESSES

BY EMMANUEL ROY

*Université Paris*

We give a second look at stationary stable processes by interpreting the self-similar property at the level of the Lévy measure as characteristic of a Maharam system. This allows us to derive structural results and their ergodic consequences.

**1. Introduction.** In a fundamental paper [9], Rosiński revealed the hidden structure of stationary symmetric  $\alpha$ -stable ( $S\alpha S$ ) processes. Namely, he proved that, following Hardin [5], through what is called a *minimal spectral representation*, such a process is driven by a nonsingular dynamical system.

Such a result was proved to classify those processes according to their ergodic properties such as various kinds of mixing. In [13], we used a different approach as we considered the whole family of stationary infinitely divisible processes without Gaussian part (called *IDp processes*). The key tool there was the Lévy measure system of the process, which was measure-preserving and not just merely nonsingular. So far, in the stable case, the connection between the Lévy measure and the nonsingular system was not clear. This is the purpose of this paper, to fill the gap and go beyond both approaches.

Indeed, we will prove that Lévy measure systems of  $\alpha$ -stable processes have the form of a so-called Maharam system. This observation has some interesting consequences as it allows us to derive very quickly minimal spectral representations in the  $S\alpha S$  case, to reinforce factorization results, and to refine ergodic classification.

Let us explain very loosely the mathematical features of stable distributions we will be using. Observe that stable distributions are characterized by a self-similar property which is obvious when observing the corresponding Lévy process:

If  $X_t$  is an  $\alpha$ -stable Lévy process, then  $b^{-1/\alpha} X_{bt}$  has the same distribution.

However, if not obvious or useful, this property is also present for any  $\alpha$ -stable object but takes another form. The common feature is to be found in the Lévy measure:

Loosely speaking, if  $\{X_t\}_{t \in S}$  is an  $\alpha$ -stable process indexed by a set  $S$ , then for any positive number  $c$ , the image of the Lévy measure  $Q$  by the map  $R_c := \{x_t\}_{t \in S} \mapsto \{cx_t\}_{t \in S}$  is  $c^{-\alpha} Q$ .

This property of the Lévy measure is characteristic of  $\alpha$ -stable processes and can be translated into an ergodic theoretic statement:

---

Received January 2010; revised March 2011.

*MSC2010 subject classifications.* Primary 60G52, 60G10, 37A40; secondary 37A50.

*Key words and phrases.* Stable stationary processes, Maharam system, ergodic properties.

The measurable nonsingular flow  $\{R_c\}_{c \in \mathbb{R}_+}$  is dissipative and the multiplicative coefficient  $c^{-\alpha}$  has an outstanding importance in this matter, since it reveals the structure of a Maharam transformation. The importance is even greater when there is more invariance involved (stationary  $\alpha$ -stable processes, etc. . . .), as in the present paper.

The paper is organized as follows. In Section 2 we recall what a spectral representation is, and in Section 3, we give the necessary background in nonsingular ergodic theory. Maharam systems are introduced in Section 4, and the link with Lévy measures of stable processes, together with spectral representations is exposed in Section 5. Section 6 is a refinement of the structure of stable processes. We deduce from the preceding results some ergodic properties in Section 7.

**2. Spectral representation.** We warn the reader that we will, most of the time, omit the implicit “ $\mu$ -a.e.” or “modulo null sets” throughout the document.

A family of functions  $\{f_t\}_{t \in T} \subset L^\alpha(\Omega, \mathcal{F}, \mu)$  where  $(\Omega, \mathcal{F}, \mu)$  is a  $\sigma$ -finite Lebesgue space is said to be a spectral representation of  $S\alpha S$  process  $\{X_t\}_{t \in T}$  if

$$\{X_t\}_{t \in T} = \left\{ \int_{\Omega} f_t(\omega) M(d\omega) \right\}_{t \in T}$$

holds in distribution,  $M$  being an independently scattered  $S\alpha S$ -random measure on  $(\Omega, \mathcal{F})$  with intensity measure  $\mu$ .

We will say that a spectral representation is *proper* if  $\text{Supp}\{f_t, t \in T\} = \Omega$ . Of course we obtain a proper representation from a general one by removing the complement of  $\text{Supp}\{f_t, t \in T\}$ .

To express that a representation contains the strict minimum to define the process, the notion of minimality has been introduced (Hardin [5]):

A spectral representation is said to be  $\{f_t\}_{t \in T} \subset L^\alpha(\Omega, \mathcal{F}, \mu)$  *minimal* if it is proper and  $\sigma(\frac{f_t}{f_s} 1_{\{f_s \neq 0\}}, s, t \in T) = \mathcal{F}$ .

Hardin proved in [5] the existence of minimal representations for  $S\alpha S$  processes.

In the stationary case ( $T = \mathbb{R}$  or  $\mathbb{Z}$ ), Rosiński has explained the form of the spectral representation:

**THEOREM 1 (Rosiński).** *Let  $\{f_t\}_{t \in T} \subset L^\alpha(\Omega, \mathcal{F}, \mu)$  be a minimal representation of a stationary  $S\alpha S$ -process; then there exists nonsingular flow  $\{\phi_t\}_{t \in T}$  on  $(\Omega, \mathcal{F}, \mu)$  and a cocycle  $\{a_t\}_{t \in T}$  for this flow with values in  $\{-1, 1\}$  (or in  $|z| = 1$  in the complex case) such that, for each  $t \in T$ ,*

$$f_t = a_t \left\{ \frac{d\mu \circ \phi_t}{d\mu} \right\}^{1/\alpha} (f_0 \circ \phi_t).$$

**3. Some terminology.** A quadruplet  $(\Omega, \mathcal{F}, \mu, T)$  is called a *dynamical system* or shortly a *system* if  $T$  is a *nonsingular automorphism*, that is, a bijective bi-measurable map such that  $T^* \mu \sim \mu$ . If  $T_* (\mu) = \mu$ , then  $(\Omega, \mathcal{F}, \mu, T)$  is a *measure-preserving* (abr. *m.p.*) dynamical system.

A system  $(\Omega_2, \mathcal{F}_2, \mu_2, T_2)$  is said to be a *nonsingular (resp. measure preserving) factor* of the system  $(\Omega_1, \mathcal{F}_1, \mu_1, T_1)$  if there exists a measurable *nonsingular (resp. measure preserving) homomorphism* between them, that is, a measurable map  $\Phi$  from  $\Omega_1$  to  $\Omega_2$  such that  $\Phi T_1 = T_2 \Phi$  and  $\Phi^* \mu_1 \sim \mu_2$  (resp.  $\Phi^* \mu_1 = \mu_2$ ). If  $\Phi$  is invertible and bi-measurable it is called a *nonsingular (resp. measure preserving) isomorphism*, and the system is said to be *nonsingular (resp. measure preserving) isomorphic*.

3.1. *Krieger types.* Consider a nonsingular dynamical system  $(\Omega, \mathcal{F}, \mu, T)$ . A set  $A \in \mathcal{F}$  such that  $\mu(A) > 0$  is said to be *periodic* of period  $n$  if  $T^i A, 0 \leq i \leq n - 1$ , are disjoint and  $T^n A = A$  and *wandering* if  $T^i A, i \in \mathbb{Z}$  are disjoint. A set is *exhaustive* if  $\bigcup_{k \in \mathbb{Z}} T^k A = \Omega$ . A system is *conservative* if there is no wandering set and *dissipative* if there is an exhaustive wandering set.

$(\Omega, \mathcal{F}, \mu, T)$  is said to be of *Krieger type*:

- $I_n$  if there exists an exhaustive set of period  $n$ ;
- $I_\infty$  if it is dissipative;
- $\Pi_1$  if there is no periodic set and exists an equivalent finite  $T$ -invariant measure;
- $\Pi_\infty$  if it is conservative with an equivalent infinite  $T$ -invariant continuous measure but no absolutely continuous finite  $T$ -invariant measure;
- $\text{III}$  if there is no absolutely continuous  $T$ -invariant measure.

**4. Maharam transformation.**

DEFINITION 2. An m.p. dynamical system is said to be *Maharam* if it is isomorphic to  $(\Omega \times \mathbb{R}, \mathcal{F} \otimes \mathcal{B}, \mu \otimes e^s ds, \tilde{T})$ , where  $T$  is a nonsingular automorphism of  $(\Omega, \mathcal{F}, \mu)$ , and  $\tilde{T}$  is defined by

$$\tilde{T}(\omega, s) := \left( T(\omega), s - \ln \frac{dT_*^{-1} \mu}{d\mu}(\omega) \right).$$

Observe that the dissipative flow  $\{\tau_t\}_{t \in \mathbb{R}}$  defined by  $\tau_t := (\omega, s) \mapsto (\omega, s - t)$  commutes with  $\tilde{T}$ .

Note that we have chosen the usual additive representation but we could (and eventually will!) use the following multiplicative representation of a Maharam system. Take  $0 < \alpha < 2$ . We can represent  $(\Omega \times \mathbb{R}, \mathcal{F} \otimes \mathcal{B}, \mu \otimes e^s ds, \tilde{T})$  by the system  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$ , where  $\tilde{T}_\alpha$  is defined by

$$\tilde{T}_\alpha(\omega, s) := \left( T(\omega), s \left( \frac{dT_*^{-1} \mu}{d\mu}(\omega) \right)^{1/\alpha} \right).$$

The isomorphism is provided by the map  $(\omega, s) \mapsto (\omega, (2 - \alpha)^{-1/(2-\alpha)} e^{(2-\alpha)s})$ . Observe that, under this isomorphism,  $\{\tau_t\}_{t \in \mathbb{R}}$  is changed into  $\{S_{e^{t/\alpha}}\}_{t \in \mathbb{R}_+^*}$ , where  $S_t$  is the multiplication by  $t$  on the second coordinate.

In [2], the authors proved the following characterization, as a straightforward application of Krengel’s representation of dissipative transformations:

**THEOREM 3.** *A system  $(X, \mathcal{A}, \nu, \gamma)$  is Maharam if and only if there exists a measurable flow  $\{Z_t\}_{t \in \mathbb{R}}$  commuting with  $\gamma$  such that  $(Z_t)_* \nu = e^t \nu$ .  $\{Z_t\}_{t \in \mathbb{R}}$  corresponds to  $\{\tau_t\}_{t \in \mathbb{R}}$  under the isomorphism with the Maharam system under the additive representation.*

In the original theorem they assumed ergodicity of  $\gamma$  to prove that the resulting nonsingular transformation  $T$  in the above representation was actually living on a nonatomic measure space  $(\Omega, \mathcal{F}, \mu)$ . The ergodicity assumption is therefore not necessary in the way we present this theorem.

We end this section with a very natural lemma which is part of folklore. We omit the proof.

**LEMMA 4.** *Consider two Maharam systems  $(\Omega_1 \times \mathbb{R}_+^*, \mathcal{F}_1 \otimes \mathcal{B}, \mu_1 \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_1)$  and  $(\Omega_2 \times \mathbb{R}_+^*, \mathcal{F}_2 \otimes \mathcal{B}, \mu_2 \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_2)$ , and denote by  $\{S_t\}_{t \in \mathbb{R}_+^*}$  and  $\{Z_t\}_{t \in \mathbb{R}_+^*}$  their respective multiplicative flows. Assume there exists a (measure-preserving) factor map (resp. isomorphism)  $\Phi$  between the two systems such that, for all  $t \in \mathbb{R}_+^*$ ,  $S_t \Phi = Z_t \Phi$ . Then  $\Phi$  induces a nonsingular factor map (resp. isomorphism)  $\phi$  between  $(\Omega_1, \mathcal{F}_1, \mu_1, T_1)$  and  $(\Omega_2, \mathcal{F}_2, \mu_2, T_1)$ .*

**REMARK 5.** Observe also that the Maharam systems associated to  $(\Omega, \mathcal{F}, \mu_1, T)$  and  $(\Omega, \mathcal{F}, \mu_2, T)$  where  $\mu_1 \sim \mu_2$  are isomorphic.

4.1. *Refinements of type III (see [3]).* Since the flow  $\{S_t\}_{t \in \mathbb{R}}$  commutes with  $\tilde{T}$ , it acts nonsingularly on the space  $(Z, \nu)$  of ergodic components of  $\tilde{T}$  and is called the *associated flow* of  $T$ . This flow is ergodic whenever  $T$  is ergodic, and its form allows us to classify ergodic type III systems:

- $T$  is of type  $\text{III}_\lambda$ ,  $0 < \lambda < 1$ , if the associated flow is the periodic flow  $x \mapsto x + t \pmod{-\log \lambda}$ ;
- $T$  is of type  $\text{III}_0$ , if the associated flow is free;
- $T$  is of type  $\text{III}_1$ , if the associated flow is the trivial flow on a singleton.

In particular  $\tilde{T}$  is ergodic if and only if  $T$  is of type  $\text{III}_1$ .

**5. Lévy measure as Maharam system and spectral representations.**

5.1. *Lévy measure of  $\alpha$ -stable processes.* For simplicity we will only consider discrete time stationary processes.

Let us recall, following [8] (see also [13]), that the Lévy measure of stationary IDp process  $X$  of distribution  $\mathbb{P}$  is the shift-invariant  $\sigma$ -finite measure on  $\mathbb{R}^{\mathbb{Z}}$ ,  $Q$ , such that  $Q(0_{\mathbb{R}^{\mathbb{Z}}}) = 0$ ,  $\int_{\mathbb{R}^{\mathbb{Z}}} (x_0^2 \wedge 1) Q(d\{x_n\}_{n \in \mathbb{Z}}) < \infty$  and

$$\begin{aligned} & \mathbb{E} \left[ \exp \left( i \sum_{k=n_1}^{n_2} a_k X_k \right) \right] \\ &= \exp \left[ \int_{\mathbb{R}^{\mathbb{Z}}} \left( \exp \left( i \sum_{k=n_1}^{n_2} a_k x_k \right) - 1 - i \sum_{k=n_1}^{n_2} a_k c(x_k) \right) Q(d\{x_n\}_{n \in \mathbb{Z}}) \right] \end{aligned}$$

for any choice of  $-\infty < n_1 \leq n_2 < +\infty$ ,  $\{a_k\}_{n_1 \leq n_2} \in \mathbb{R}^{n_2 - n_1}$ .

$c$  is defined by:

$$\begin{aligned} c(x) &= -1 && \text{if } x < -1; \\ c(x) &= x && \text{if } -1 \leq x \leq 1; \\ c(x) &= 1 && \text{if } x > 1. \end{aligned}$$

The system  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$  where  $S$  is the shift on  $\mathbb{R}^{\mathbb{Z}}$  is called the *Lévy measure system* associated to the process  $X$ .

The  $\alpha$ -stable stationary processes,  $0 < \alpha < 2$ , are (see Chapter 3 in [17]) completely characterized as those IDp processes such that their Lévy measure satisfies

$$(5.1) \quad (R_t)_* Q = t^{-\alpha} Q$$

for any positive  $t$ ,  $R_t$  being the multiplication by  $t$ , that is,

$$\{x_n\}_{n \in \mathbb{Z}} \mapsto \{tx_n\}_{n \in \mathbb{Z}}.$$

We also recall the fundamental result of Maruyama that allows to represent any IDp process with Lévy measure  $Q$  as a stochastic integral with respect to a Poisson measure with intensity  $Q$ .

**THEOREM 6 (Maruyama representation [8]).** *Let  $\mathbb{P}$  be the distribution of a stationary IDp process with Lévy measure  $Q$  and  $((\mathbb{R}^{\mathbb{Z}})^*, (\mathcal{B}^{\otimes \mathbb{Z}})^*, Q^*, S_*)$  the Poisson measure over the Lévy measure system  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$ . Set  $X_0$  as  $\{x_n\}_{n \in \mathbb{Z}} \mapsto x_0$  and define, on  $(\mathbb{R}^{\mathbb{Z}})^*$ , the stochastic integral  $I(X_0)$  as the limit in probability, as  $n$  tends to infinity, of the random variables*

$$v \mapsto \int_{|X_0| > 1/n} X_0 dv - \int_{|X_0| > 1/n} c(X_0) dQ.$$

*Then the process  $\{I(X_0) \circ S_*^n\}_{n \in \mathbb{Z}}$  has distribution  $\mathbb{P}$ .*

5.2. Lévy measure as Maharam system.

**THEOREM 7.** *Let  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$  be the Lévy measure system of an  $\alpha$ -stable stationary process. Then there exists a probability space  $(\Omega, \mathcal{F}, \mu)$ , a nonsingular transformation  $T$ , a function  $f \in L^\alpha(\mu)$  such that, if  $M$  denotes the map  $(\omega, t) \mapsto tf(\omega)$ , then the map  $\Theta := (\omega, t) \mapsto \{M \circ \tilde{T}_\alpha^n(\omega, t)\}_{n \in \mathbb{Z}}$  yields an isomorphism of the Maharam system  $(\Omega \times \mathbb{R}_+, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$  with  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$ .*

**PROOF.** First observe that Theorem 3 can be applied to  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$  since the measurable and (obviously) dissipative flow  $\{R_{e^{t/\alpha}}\}_{t \in \mathbb{R}}$  satisfies the hypothesis, thanks to equation (5.1). Therefore, there exists an isomorphism  $\Psi$  between the Maharam system  $(\Omega \times \mathbb{R}_+, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$  and  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$  for an appropriate nonsingular system  $(\Omega, \mathcal{F}, \mu, T)$ . Set  $f := \Psi(\omega, 1)_0$  [i.e.,  $\Psi(\omega, 1)_0$  is the 0th coordinate of the sequence  $\Psi(\omega, 1)$ ], and let us check that  $f \in L^\alpha(\mu)$ . Indeed, as  $Q$  is a Lévy measure, we have

$$\int_{\mathbb{R}^{\mathbb{Z}}} x_0^2 \wedge 1 Q(d\{x_n\}_{n \in \mathbb{Z}}) < \infty,$$

but since  $\Psi$  is an isomorphism and  $\Psi(\omega, t) = \Psi \circ S_t(\omega, 1) = R_t \circ \Psi(\omega, 1) = t\Psi(\omega, 1)$ , we have

$$\begin{aligned} & \int_{\mathbb{R}^{\mathbb{Z}}} x_0^2 \wedge 1 Q(d\{x_n\}_{n \in \mathbb{Z}}) \\ &= \int_{\Omega} \int_{\mathbb{R}_+} \Psi(\omega, t)_0^2 \wedge 1 \frac{1}{t^{\alpha+1}} dt \mu(d\omega), \\ & \int_{\Omega} \int_{\mathbb{R}_+} (t^2 \Psi(\omega, 1)_0^2) \wedge 1 \frac{1}{t^{\alpha+1}} dt \mu(d\omega) \\ &= \left( \int_{\mathbb{R}_+} z^2 \wedge 1 \frac{1}{z^{\alpha+1}} dz \right) \int_{\Omega} |\Psi(\omega, 1)_0|^\alpha \mu(d\omega) \end{aligned}$$

after the change of variable  $z := t|\Psi(\omega, 1)_0|$ . Therefore  $\int_{\Omega} |\Psi(\omega, 1)_0|^\alpha \mu(d\omega) < \infty$ .  $\square$

In the symmetric case we can make the theorem more precise:

**THEOREM 8.** *Let  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$  be the Lévy measure system of a symmetric  $\alpha$ -stable stationary process. Then there exists a probability space  $(X, \mathcal{A}, \nu)$ , a nonsingular transformation  $R$ , a function  $f \in L^\alpha(\nu)$  and a measurable map  $\xi: X \rightarrow \{-1, 1\}$  such that, if  $M$  denotes the map  $(x, t) \mapsto tf(x)$ , then the map  $(x, t) \mapsto \{M \circ \overline{R}_\alpha^n(x, t)\}_{n \in \mathbb{Z}}$  yields an isomorphism between  $(X \times \mathbb{R}^*, \mathcal{A} \otimes \mathcal{B}, \nu \otimes \frac{1}{|s|^{1+\alpha}} ds, \overline{T}_\alpha)$  with  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$ ,  $\overline{R}_\alpha$  being defined by  $(x, t) \mapsto (Rx, \xi(x)t(\frac{dR_x^{-1}\mu}{d\mu}(x))^{1/\alpha})$ .*

PROOF. Start by applying Theorem 7 to the Lévy measure.

Observe that the symmetry involves the presence of a measure preserving involution  $I$ , namely  $I\{x_n\}_{n \in \mathbb{Z}} = \{-x_n\}_{n \in \mathbb{Z}}$ .  $I$  also preserves the Lévy measure of the process. Observe also that  $I$  commutes with the shift and with the flow  $R_t$ . Therefore  $\tilde{T} := \Theta^{-1}I\Theta$  is a measure preserving automorphism of  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T})$ , and we can apply Lemma 4 to deduce that  $\tilde{T}$  induces a nonsingular involution  $\phi$  on  $(\Omega, \mathcal{F}, \mu, T)$ . It is standard that such transformation admits an equivalent finite invariant measure, so, up to another measure preserving isomorphism, we can assume that  $\phi$  preserves the probability measure  $\mu$ .

Using Rohklin’s structure theorem, the compact factor associated to the compact group  $\{\text{Id}, \phi\}$  tells us that we can represent  $(\Omega, \mathcal{F}, \mu, T)$  as  $(X \times \{-1, 1\}, \mathcal{A} \otimes \mathcal{P}\{-1, 1\}, \nu \otimes m, R_\xi)$ , where  $R$  is a nonsingular automorphism of  $(X, \mathcal{A}, \nu)$ ,  $m$  is the uniform measure on  $(\{-1, 1\}, \mathcal{P}\{-1, 1\})$ ,  $\xi$  a cocycle from  $X$  to  $\{-1, 1\}$  and  $R_\xi := (x, \varepsilon) \mapsto (Rx, \xi(x)\varepsilon)$ .

It is now clear that  $(X \times \{-1, 1\} \times \mathbb{R}_+^*, (\mathcal{A} \otimes \mathcal{P}\{-1, 1\}) \otimes \mathcal{B}, \nu \otimes m \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{S}_\xi)$  is isomorphic to  $(X \times \mathbb{R}^*, \mathcal{A} \otimes \mathcal{B}, \nu \otimes \frac{1}{|s|^{1+\alpha}} ds, \overline{R}_\alpha)$  thanks to the mapping  $(x, \varepsilon, t) \mapsto (x, 2^{1/\alpha}\varepsilon t)$  and  $\overline{R}_\alpha := (x, t) \mapsto (Rx, \xi(x)(\frac{dR_x^{-1}\mu}{d\mu}(x))^{1/\alpha}t)$ .  $\square$

5.3. *Spectral representation.* It is now very easy to derive spectral representations from the above results. In particular, if  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$  is the Lévy measure system of an  $S\alpha S$  stationary process, under the notation of Theorem 8,  $(X, \mathcal{A}, \nu)$  together with the function  $f \in L^\alpha(\nu)$ , the cocycle  $\phi$  and the nonsingular automorphism  $T$  yields a spectral representation of the process. Indeed, by building the Poisson measure over  $(X \times \mathbb{R}^*, \mathcal{A} \otimes \mathcal{B}, \nu \otimes \frac{1}{|s|^{1+\alpha}} ds, \overline{R}_\alpha)$  and by applying to it,  $f$  and  $\xi$  (Theorem 3.12.2, page 156 in [16]), we recover the  $S\alpha S$  process with Lévy measure  $Q$ , which proves the validity of the spectral representation. The minimality can be obtained without difficulty thanks to Proposition 2.2 in [10].

5.4. *Maharam systems as Lévy measure.* We can ask if whether a Maharam system  $(\Omega \times \mathbb{R}_+, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$  can be coded into a Lévy measure system of a stable process. We can answer this question affirmatively in the only interesting case, that is, when the Maharam system has no finite absolutely continuous  $\tilde{T}_\alpha$ -invariant measure, that is, when the resulting Lévy measure system leads to an ergodic stable process.

Recall that a Maharam system  $(\Omega \times \mathbb{R}_+, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$  has no finite absolutely continuous  $\tilde{T}_\alpha$ -invariant measure if and only if the nonsingular system  $(\Omega, \mathcal{F}, \mu, T)$  has the same property. But then a famous theorem of Krengel [7] shows that such a system possesses a 2-generator, that is, there exists a measurable function  $f : \Omega \rightarrow \{0, 1\}$ ,  $\mu\{f = 1\} < \infty$ , such that  $\sigma\{f \circ T^n, n \in \mathbb{Z}\} = \mathcal{F}$ .

To be more precise, this means that, up to isomorphism,  $(\Omega, \mathcal{F}, \mu, T)$  can be represented as  $(\{0, 1\}^{\mathbb{Z}}, \mathcal{B}(\{0, 1\}^{\mathbb{Z}}), \nu, S)$  for an appropriate measure  $\nu$ , where

$S$  is the shift transformation. Then the Maharam system can be represented as  $(\{0, 1\}^{\mathbb{Z}} \times \mathbb{R}_+^*, \mathcal{B}(\{0, 1\}^{\mathbb{Z}}) \otimes \mathcal{B}, \nu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{S}_\alpha)$ . But if  $\varphi$  is the map  $(\{x_n\}_{n \in \mathbb{Z}}, t) \mapsto \{tx_n\}_{n \in \mathbb{Z}}$  and  $Q = \varphi_*(\nu \otimes \frac{1}{s^{1+\alpha}} ds)$ , we obtain a Lévy measure system  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$  of an  $\alpha$ -stable system, as  $\{x_n\}_{n \in \mathbb{Z}} \mapsto \{x_0\}$  is in  $L^\alpha(\mu)$  (see the proof of Theorem 7). Moreover, the sequence  $\{y_n\}_{n \in \mathbb{Z}}$  takes only two values, 0 or  $\sup\{y_n\}_{n \in \mathbb{Z}}$   $Q$ -almost everywhere ( $\{y_n\}_{n \in \mathbb{Z}}$  can not be identically zero with positive measure as such a constant sequence forms a shift-invariant set of finite measure); therefore,  $\varphi^{-1}$  exists and is defined by  $\{y_n\}_{n \in \mathbb{Z}} \mapsto (\sup\{y_n\}_{n \in \mathbb{Z}}, \{\frac{y_n}{\sup\{y_n\}_{n \in \mathbb{Z}}}\}_{n \in \mathbb{Z}})$ .

$(\{0, 1\}^{\mathbb{Z}} \times \mathbb{R}_+^*, \mathcal{B}(\{0, 1\}^{\mathbb{Z}}) \otimes \mathcal{B}, \nu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{S}_\alpha)$  is isomorphic to  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$ .

**6. Refinements of the representation.** Ergodic stationary processes are building blocks of stationary processes; prime numbers are the building blocks of integers; factors are building blocks of Von Neumann algebras etc. What are the building blocks of stationary infinitely divisible processes? Let us get more precise.

Given a stationary ID process  $X$ , what are the solutions to the equation (in distribution)

$$X = X_1 + X_2,$$

where  $X_1$  and  $X_2$  are independent stationary ID processes. Of course, if  $Q$  is the Lévy measure of  $X$ , then taking  $X_1$  with Lévy measure  $c_1 Q$  and  $X_2$  with Lévy measure  $c_2 Q$  with  $c_1 + c_2 = 1$  gives a solution. If these are the only solutions, we said in [12] that  $X$  is *pure*, meaning that is impossible to reduce  $X$  to “simpler” pieces. It was then very easy to show that  $X$  is pure if and only if its Lévy measure is ergodic.

**PROPOSITION 9.** *A stationary IDp process  $X$  is pure if and only if its Lévy measure  $Q$  is ergodic.*

**PROOF.** Assume  $Q$  is not ergodic. There exists a partition of  $\mathbb{R}^{\mathbb{Z}}$  into two shift invariant sets  $A$  and  $B$  both of positive measure. Therefore,  $Q|_A$  and  $Q|_B$  can be taken as Lévy measures of two stationary IDp processes  $X_A$  and  $X_B$  and taking them independently leads to

$$X = X_A + X_B$$

in distribution, as  $Q = Q|_A + Q|_B$ .

In the converse, assume  $Q$  is ergodic, and suppose there exist independent stationary IDp processes  $X_1$  and  $X_2$  with Lévy measure  $Q_1$  and  $Q_2$  such that

$$X = X_1 + X_2$$

holds in distribution. As  $Q = Q_1 + Q_2$  we get  $Q_1 \ll Q$ . But as  $Q$  is ergodic, this in turns implies that there exists  $c > 0$  such that  $Q_1 = cQ$  and thus  $Q_2 = (1 - c)Q$ . □

In this section, we will try to comment the above equation according to the Krieger type of the associated nonsingular transformation. A description of the interesting class of those stable processes driven by nonsingular transformations of type  $\text{III}_0$  is unknown.

6.1. *The type  $\text{III}_1$  case, pure stable processes.* It was an open question whether there exist pure stable processes. It can now be solved thanks to the Maharam structure of the Lévy measure: an  $\alpha$ -stable process is pure if and only if the underlying nonsingular system is of type  $\text{III}_1$ .

The existence of pure stable processes (guaranteed by the comments made in Section 5.4) is reassuring as it validates the specific study of stable processes.

6.2. *The type  $\text{III}_\lambda$  case,  $0 < \lambda < 1$ .* In this section we derive the form of those  $\alpha$ -stable processes associated with an ergodic, type  $\text{III}_\lambda$  nonsingular automorphism,  $0 < \lambda < 1$ .

6.2.1. *Semi-stable stationary processes.* An infinitely divisible probability measure  $\mu$  on  $\mathbb{R}^d$  is called  $\alpha$ -semi-stable with span  $b$  if its Fourier transform satisfies

$$\hat{\mu}(z)^{b^\alpha} = \hat{\mu}(bz)e^{i\langle c, z \rangle}$$

for some  $c \in \mathbb{R}^d$ .

By extension, an  $\alpha$ -semi-stable process process is a process whose finite-dimensional distributions are  $\alpha$ -semi-stable. Using once again results of Chapter 3 in [17], one gets the following characterization of  $\alpha$ -semi-stable stationary processes:

A shift-invariant Lévy measure  $Q$  on  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}})$  is the Lévy measure of an  $\alpha$ -semi-stable stationary process of span  $b > 0$  if and only if it satisfies

$$(R_b)_* Q = b^{-\alpha} Q,$$

where  $R_b$  is the multiplication by  $b$

$$\{x_n\}_{n \in \mathbb{Z}} \mapsto \{bx_n\}_{n \in \mathbb{Z}}.$$

Of course by iterating  $R_b$ , we easily observe that  $(R_{b^n})_* Q = b^{-n\alpha} Q$  for all  $n \in \mathbb{Z}$ .

6.2.2. *Discrete Maharam extension.* Assume  $(\Omega, \mathcal{F}, \mu, T)$  is a nonsingular system such that there exists  $\lambda > 0$  so that  $\frac{dT_*^{-1}\mu}{d\mu} \in \{\lambda^n, n \in \mathbb{Z}\}$   $\mu$ -almost everywhere. We can form its *discrete Maharam extension*, that is, the m.p. system  $(\Omega \times \mathbb{Z}, \mathcal{F} \otimes \mathcal{B}, \mu \otimes \lambda^n dn, \tilde{T})$  where  $\lambda^n dn$  stands for the measure  $\sum_{n \in \mathbb{Z}} \lambda^n \delta_n$  on  $(\mathbb{Z}, \mathcal{B})$  and  $\tilde{T}$  is defined by

$$\tilde{T}(\omega, n) = \left( T\omega, n - \log_\lambda \frac{dT_*^{-1}\mu}{d\mu}(\omega) \right).$$

6.2.3. *Ergodic decomposition of Maharam extension of type III<sub>λ</sub> transformations.* Let  $(\Omega, \mathcal{F}, \mu, T)$  be an ergodic type III<sub>λ</sub> system. Up to a change of measure we can assume that the Radon–Nykodim derivative take its values in the group  $\{\lambda^n, n \in \mathbb{Z}\}$  where  $r(T) = \{0, \lambda^n, n \in \mathbb{Z}, +\infty\}$  is the ratio set of  $T$  (see [6]). Therefore, the discrete Maharam extension  $(\Omega \times \mathbb{Z}, \mathcal{F} \otimes \mathcal{B}, \beta\mu \otimes \lambda^n dn, \tilde{T})$ , where  $\beta := \int_0^{-\ln \lambda} e^{-s} ds$ , exists. Now form the product system

$$\left( \Omega \times \mathbb{Z} \times [0, -\ln \lambda[, \mathcal{F} \otimes \mathcal{B} \otimes \mathcal{B}([0, -\ln \lambda]), \beta\mu \otimes \lambda^n dn \otimes \frac{e^{-s}}{\beta} ds, \tilde{T} \times \text{Id} \right).$$

The dissipative nonsingular flow  $S_t : (\omega, n, s) \mapsto (\omega, n + \lfloor \frac{s-t}{-\ln \lambda} \rfloor, s - t + \ln \lambda \lfloor \frac{s-t}{-\ln \lambda} \rfloor)$  satisfies  $S_t \circ \tilde{T} \times \text{Id} = \tilde{T} \times \text{Id} \circ S_t$  and  $(S_t)_* \mu \otimes \lambda^n dn \otimes e^{-s} ds = e^{-t} \mu \otimes \lambda^n dn \otimes e^{-s} ds$ , and it is very easy to see that  $(\mathbb{R} \times [0, -\ln \lambda[, \mathcal{B} \otimes \mathcal{B}([0, -\ln \lambda]), \lambda^n dn \otimes e^{-s} ds)$  is just a reparametrization of  $(\mathbb{R}, \mathcal{B}, e^s ds)$ , thanks to the mapping  $(n, s) \mapsto -n \ln \lambda - s$ .

Therefore  $(\Omega \times \mathbb{Z} \times [0, -\ln \lambda[, \mathcal{F} \otimes \mathcal{B} \otimes \mathcal{B}([0, -\ln \lambda]), \mu \otimes \lambda^n dn \otimes e^{-s} ds, \tilde{T} \times \text{Id})$  can be seen as the Maharam extension of  $(\Omega, \mathcal{F}, \mu, T)$ .

It remains to prove the ergodicity of  $(\Omega \times \mathbb{Z}, \mathcal{F} \otimes \mathcal{B}, \beta\mu \otimes \lambda^n dn, \tilde{T})$ . This follows, for example, from Corollary 5.4 in [18], as the ratio set is precisely the set of essential values of the Radon–Nykodim cocycle.

We then obtain the ergodic decomposition of the Maharam extension: it is the discrete Maharam extension  $(\Omega \times \mathbb{Z}, \mathcal{F} \otimes \mathcal{B}, \beta\mu \otimes \lambda^n dn, \tilde{T})$  randomized by the measure  $\frac{e^{-s}}{\beta} ds$  on  $[0, -\ln \lambda[$ .

6.2.4. *Application to stable processes.* Let  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$  be the Lévy measure system of an  $\alpha$ -stable process driven by an ergodic type III<sub>λ</sub> system  $(\Omega, \mathcal{F}, \mu, T)$ , and let  $f \in L^\alpha(\mu)$  be given as in Theorem 7. Let  $b > 1$  so that  $b^{-\alpha} = \lambda$ , we need to obtain a multiplicative version of the above structure adapted to our parameters. Up to a change of measure we can assume that  $(\frac{dT_*^{-1}\mu}{d\mu})^{1/\alpha} \in \{b^n, n \in \mathbb{Z}\}$   $\mu$ -almost everywhere. Consider the discrete Maharam extension  $(\Omega \times G_b, \mathcal{F} \otimes \mathcal{B}, \beta\mu \otimes m_b, \tilde{T})$  (in a multiplicative representation) where  $\beta = \int_1^b \frac{1}{s^{1+\alpha}} ds$ ,  $m_b$  is the measure  $\sum_{g \in G_b} g^{-\alpha} \delta_g$  on  $G_b := \{b^n, n \in \mathbb{Z}\}$  and  $\tilde{T} := (\omega, g) \mapsto (T\omega, g(\frac{dT_*^{-1}\mu}{d\mu}(\omega))^{1/\alpha})$ . Form the system  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q^r, S)$  as a factor of  $(\Omega \times G_b, \mathcal{F} \otimes \mathcal{B}, \beta\mu \otimes m_b, \tilde{T})$  given by the mapping  $\varphi := (\omega, g) \mapsto \{M \circ \tilde{T}^n(\omega, g)\}_{n \in \mathbb{Z}}$  where  $M(\omega, g) = gf(\omega)$  and  $Q^r = \varphi_*(\beta\mu \otimes m_b)$ .

Now, as above, we recover the Maharam extension of  $(\Omega, \mathcal{F}, \mu, T)$  by considering  $(\Omega \times G_b \times [1, b[, \mathcal{F} \otimes \mathcal{B} \otimes \mathcal{B}([1, b]), \beta\mu \otimes m_b \otimes \frac{1}{\beta s^{1+\alpha}} ds, \tilde{T} \times \text{Id})$ . As the system  $(G_b \times [1, b[, \mathcal{B} \otimes \mathcal{B}([1, b]), m_b \otimes \frac{1}{s^{1+\alpha}} ds)$  is isomorphic to  $(\mathbb{R}_+^*, \mathcal{B}, \frac{1}{s^{1+\alpha}} ds)$  thanks to  $(g, t) \mapsto gt$ , we obtain  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$  by applying the map  $(\{x_n\}_{n \in \mathbb{Z}}, t) \mapsto \{tx_n\}_{n \in \mathbb{Z}}$  to  $(\mathbb{R}^{\mathbb{Z}} \times [1, b[, \mathcal{B}^{\otimes \mathbb{Z}} \otimes \mathcal{B}([1, b]), Q^r \otimes \frac{1}{\beta s^{1+\alpha}} ds, S \times$

Id). At last, we can check that  $Q^r$  is a Lévy measure, and indeed we know that

$$\int_{\mathbb{R}^{\mathbb{Z}}} (x_0^2 \wedge 1) Q(d\{x_n\}_{n \in \mathbb{Z}}) < +\infty,$$

but

$$\int_{\mathbb{R}^{\mathbb{Z}}} (x_0^2 \wedge 1) Q(d\{x_n\}_{n \in \mathbb{Z}}) = \int_1^b \left( \int_{\mathbb{R}^{\mathbb{Z}}} ((sx_0)^2 \wedge 1) Q^r(d\{x_n\}_{n \in \mathbb{Z}}) \right) \frac{1}{\beta s^{1+\alpha}} ds;$$

therefore, for some  $1 \leq s < b$ ,  $\int_{\mathbb{R}^{\mathbb{Z}}} ((sx_0)^2 \wedge 1) Q^r(d\{x_n\}_{n \in \mathbb{Z}}) < +\infty$ , and this is enough to prove that  $Q^r$  is a Lévy measure.

$(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q^r, S)$  is the Lévy measure system of an  $\alpha$ -semi-stable stationary process with span  $b$ . Heuristically, if  $X$  has Lévy measure  $Q$ ,  $X$  can be thought of as the continuous sum of independent processes  $Y^t$ ,  $1 \leq t < b$  weighted by the probability measure  $\frac{1}{\beta s^{1+\alpha}} ds$  where  $\frac{1}{t} Y^t$  has Lévy measure  $Q^r$ . More formally, if  $((\Omega \times G_b \times [1, b]^*, (\mathcal{F} \otimes \mathcal{B} \otimes \mathcal{B}([1, b]))^*, (\beta\mu \otimes m_b \otimes \frac{1}{\beta s^{1+\alpha}} ds)^*, (\tilde{T} \times \text{Id})_*)$  denotes the Poisson suspension over  $(\Omega \times G_b \times [1, b], \mathcal{F} \otimes \mathcal{B} \otimes \mathcal{B}([1, b]), \beta\mu \otimes m_b \otimes \frac{1}{\beta s^{1+\alpha}} ds, \tilde{T} \times \text{Id})$ , then, if  $I$  denotes the stochastic integral as in Theorem 6,  $X := \{I\{M_1\} \circ (\tilde{T} \times \text{Id})_*^n\}_{n \in \mathbb{Z}}$  has Lévy measure  $Q$  and  $Y := \{I\{M_2\} \circ (\tilde{T} \times \text{Id})_*^n\}_{n \in \mathbb{Z}}$  where  $M_1(\omega, g, s) = s g f(\omega)$  and  $M_2(\omega, g, s) = g f(\omega)$ .

We therefore observe that  $X$  is entirely determined by a pure  $\alpha$ -semi-stable stationary process with span  $b$ ,  $Y$ . It is very easy to see that  $X$  and  $Y$  share the same type of mixing.

6.2.5. *Examples.* It is not difficult to exhibit examples of stable processes of the kind described above, as the structure detailed allows us to build such processes. We can, for example, consider the systems  $T_p$ ,  $\frac{1}{2} < p < 1$  introduced in [4]. We will follow the presentation given in ([1], page 104).

Let  $\Omega$  be the group of dyadic integers, let  $\tau$  acts by translation by  $\underline{1}$  on  $\Omega$  and for  $\frac{1}{2} < p < 1$ , let  $\mu_p$  be a probability measure on  $\Omega$  defined on cylinders by

$$\mu_p([\varepsilon_1, \dots, \varepsilon_n]) = \prod_{k=1}^n p(\varepsilon_k),$$

where  $p(0) = 1 - p$  and  $p(1) = p$ .

If we set  $\frac{1-p}{p} = \lambda$ , we get

$$\frac{d\tau_*^{-1} \mu_p}{d\mu_p} = \lambda^\phi,$$

where  $\phi(x) = \min\{n \in \mathbb{N}, x_n = 0\} - 2$ . It is proved in [4] that the discrete Maharam extension  $(\Omega \otimes \mathbb{Z}, \mathcal{F} \otimes \mathcal{B}, \mu \otimes \lambda^n dn, \tilde{\tau})$  is ergodic.

We can form a new system, which will be the Lévy measure system of a stationary semi-stable process with span  $\lambda^\alpha$ , thanks to the following map:

$$f : (\omega, n) \mapsto \lambda^{\alpha n} \sum_{i \geq 1} \omega_i 2^{-i}.$$

The Lévy measure  $Q^r$  is the image of  $\mu \otimes \lambda^n dn$  by the map  $(\omega, n) \mapsto \{f \circ \tilde{\tau}^k(\omega, n)\}_{k \in \mathbb{Z}}$ .

By randomizing this Lévy measure as explained above, we obtain the Lévy measure  $Q$  of a stationary  $\alpha$ -stable process; that is,  $Q$  is the image measure of  $Q^r \otimes \frac{1}{\beta s^{1+\alpha}} ds$  by the map

$$(\{x_n\}_{n \in \mathbb{Z}}, t) \mapsto \{tx_n\}_{n \in \mathbb{Z}}.$$

To obtain a realization of these two processes as stochastic integrals over Poisson suspensions, we can proceed as explained at the end of the preceding section.

Anticipating the next sections, we derive the ergodic properties of these processes:

$\tau$  being of type  $\text{III}_\lambda$ , the Maharam extension is of type  $\text{II}_\infty$  which means that the Lévy measure system of the corresponding stationary  $\alpha$ -stable process (with Lévy measure  $Q$ ) is of type  $\text{II}_\infty$ . Therefore the associated Poisson suspension is weakly mixing. As stochastic integrals with respect to this Poisson suspension, both processes (with Lévy measures  $Q$  and  $Q^r$ ) are weakly mixing.

Thanks to (Lemma 1.2.10, page 30 in [1]),  $\tau$  is rigid for the sequence  $\{2^n\}_{n \in \mathbb{N}}$ . Therefore, by Theorem 18 (or with a slight adaptation for the semi-stable case), both processes are also rigid for the same sequence.

**6.3. The type I and II cases.** This case is easy to deal with as we can assume the associated ergodic nonsingular system is actually measure preserving, that is, the Lévy measure system  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$  is isomorphic to  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T})$  where  $T$  preserves  $\mu$  and  $\tilde{T}$  acts as  $T \times \text{Id}$ , that is,  $\tilde{T}(\omega, t) = (T\omega, t)$ . Considering  $f \in L^\alpha(\mu)$  furnished by Theorem 7,  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T})$  is isomorphic to  $(\mathbb{R}^{\mathbb{Z}} \times \mathbb{R}_+^*, \mathcal{B}^{\otimes \mathbb{Z}} \otimes \mathcal{B}, Q^s \otimes \frac{1}{s^{1+\alpha}} ds, S \times \text{Id})$  through the map  $(\omega, t) \mapsto (\{f \circ T^n(\omega)\}_{n \in \mathbb{Z}}, t)$  and

$$\begin{aligned} \int_{\mathbb{R}^{\mathbb{Z}}} (x_0^2 \wedge 1) Q(d\{x_n\}_{n \in \mathbb{Z}}) &= \int_{\Omega} \int_{\mathbb{R}_+} ((tf(\omega))^2 \wedge 1) \frac{1}{t^{\alpha+1}} dt \mu(d\omega) \\ &= \int_{\mathbb{R}_+} \left( \int_{\mathbb{R}^{\mathbb{Z}}} ((tx_0)^2 \wedge 1) Q^s(d\{x_n\}_{n \in \mathbb{Z}}) \right) \frac{1}{t^{\alpha+1}} dt < +\infty. \end{aligned}$$

For the same reason as above,  $Q^s$  is a Lévy measure. We draw the same conclusions as in the preceding section, taking into account that the weight is now the infinite measure  $\frac{1}{t^{\alpha+1}} dt$  on  $\mathbb{R}_+^*$ , and  $Q^s$  can be any Lévy measure (of a stationary IDp process).

**7. Ergodic properties.** Some ergodic properties of general IDp stationary processes have been given in terms of ergodic properties of the Lévy measure system in [13]. For an  $\alpha$ -stable stationary processes, it is more interesting to give them in terms of the associated nonsingular system  $(\Omega, \mathcal{F}, \mu, T)$ . This work has been undertaken in the symmetric ( $S\alpha S$ ) case in a series of papers (see, in particular, [11] and [15]).

We have a new tool to deal with this problem:

As the Lévy measure of an  $\alpha$ -stable stationary processes can now be seen as the Maharam extension  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T})$  of the system  $(\Omega, \mathcal{F}, \mu, T)$ , it suffices to connect ergodic properties of  $T$  and  $\tilde{T}$ , and then apply the general results relating ergodic properties of a stationary IDp process with respect to those of its Lévy measure system.

Observe that linking ergodic properties of  $T$  and  $\tilde{T}$  is a general problem in nonsingular ergodic theory which is of great interest.

We will illustrate this in the following sections dealing with mixing,  $K$ -property and rigidity, the last two having been neglected in the  $\alpha$ -stable literature.

*7.1. Mixing.* First recall that if  $S$  is a nonsingular transformation of a measure space  $(X, \mathcal{A}, m)$ , it induces a unitary operator  $U_S$  on  $L^2(m)$  by

$$U_S f(x) = \sqrt{\frac{dS_*^{-1}\mu}{d\mu}(x)} f \circ S(x).$$

We first give a general result:

**PROPOSITION 10.** *The Maharam system  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$  is of zero type if and only if for all  $f \in L^2(\mu)$ ,  $\langle U_T^n f, f \rangle_{L^2(\mu)} \rightarrow 0$  as  $n$  tends to infinity.*

The proof can be extracted from [11] but follows also from the observation that the conditions below are equivalent (we assume that  $\mu$  is a probability):

- (1) for all  $f \in L^2(\mu)$ ,  $\langle U_T^n f, f \rangle_{L^2(\mu)} \rightarrow 0$  as  $n$  tends to infinity;
- (2)  $|\log \frac{dT_*^{-n}\mu}{d\mu}| \rightarrow \infty$  in probability;
- (3)  $m(A_\varepsilon \cap \tilde{T}_\alpha^n A_\varepsilon) \rightarrow 0$  for all  $0 < \varepsilon < 1$  where  $m = \mu \otimes \frac{1}{s^{1+\alpha}} ds$  and  $A_\varepsilon := \Omega \times [\varepsilon, \frac{1}{\varepsilon}]$ ;
- (4)  $\tilde{T}_\alpha$  is of zero type.

Combining this result with the characterization of the Lévy measure system as a Maharam system and the mixing criteria found in [13], we obtain the following theorem, already known in the  $S\alpha S$ -case (see [11]):

**THEOREM 11.** *A stationary stable process  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, \mathbb{P}, S)$  with associated system  $(\Omega, \mathcal{F}, \mu, T)$  is mixing if and only if for all  $f \in L^2(\mu)$ ,  $\langle U_T^n f, f \rangle_{L^2(\mu)} \rightarrow 0$  as  $n$  tends to infinity.*

In the forthcoming sections, we are interested in less-known ergodic properties ( $K$  property and rigidity) that have been neglected in the  $\alpha$ -stable literature.

7.2.  $K$  property.

**DEFINITION 12** (see [19]). A conservative nonsingular system  $(\Omega, \mathcal{F}, \mu, T)$  is a  $K$ -system if there exists a sub- $\sigma$ -algebra  $\mathcal{G} \subset \mathcal{F}$  such that  $T^{-1}\mathcal{G} \subset \mathcal{G}$ ,  $T^{-n}\mathcal{G} \downarrow \{\Omega, \emptyset\}$ ,  $T^n\mathcal{G} \uparrow \mathcal{F}$  and  $\frac{d\mu}{dT_*\mu}$  is  $\mathcal{G}$ -measurable.

A  $K$ -system is always ergodic (see [19]).

**DEFINITION 13.** A measure-preserving system  $(X, \mathcal{A}, m, S)$  is remotely infinite if there exists a sub- $\sigma$ -algebra  $\mathcal{C} \subset \mathcal{A}$  such that  $T^{-1}\mathcal{C} \subset \mathcal{C}$ ,  $S^n\mathcal{C} \uparrow \mathcal{A}$  and  $\bigcap_{n \geq 1} S^{-n}\mathcal{C}$  contains zero or infinite measure sets only.

**PROPOSITION 14.** *If  $(\Omega, \mathcal{F}, \mu, T)$  is a  $K$ -system which is not of type  $\text{II}_1$ , then its Maharam extension  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$  is remotely infinite.*

**PROOF.** Let  $\mathcal{G}$  be as in Definition 12. Observe that, as  $\frac{d\mu}{dT_*\mu}$  is  $\mathcal{G}$ -measurable,  $\mathcal{G} \otimes \mathcal{B}_+$  is  $\tilde{T}_\alpha$ -invariant, that is,  $\tilde{T}_\alpha^{-1}\mathcal{G} \otimes \mathcal{B}_+ \subset \mathcal{G} \otimes \mathcal{B}_+$ . Indeed, take  $g \in \mathcal{G}$ -measurable and  $f \in \mathcal{B}_+$ -measurable, and we get

$$\begin{aligned} g \otimes f(\tilde{T}_\alpha(\omega, s)) &= \left( g(T\omega), s \left( \frac{dT_*^{-1}\mu}{d\mu}(\omega) \right)^{1/\alpha} \right) \\ &= \left( g(T\omega), s \left( \frac{d\mu}{dT_*\mu}(T\omega) \right)^{1/\alpha} \right). \end{aligned}$$

We are going to show that  $\mathcal{P} := \bigcap_{n \in \mathbb{N}} \tilde{T}_\alpha^{-n}\mathcal{G} \otimes \mathcal{B}_+$  only contains sets of zero or infinite measure. Observe that, as  $S_t$  commutes with  $\tilde{T}_\alpha$  and preserves  $\mathcal{G} \otimes \mathcal{B}_+$  for all  $t > 0$ , then  $S_t^{-1}(\tilde{T}_\alpha^{-n}\mathcal{G} \otimes \mathcal{B}_+) \subset \tilde{T}_\alpha^{-n}\mathcal{G} \otimes \mathcal{B}_+$  and therefore  $S_t^{-1}\mathcal{P} \subset \mathcal{P}$  for all  $t > 0$ . Now consider the measurable union, say  $K$ , of  $\mathcal{P}$ -measurable sets of finite and positive measure. It is a  $\tilde{T}_\alpha$ -invariant set and a  $S_t$ -invariant set as well. Recall that the nonsingular action of the flow  $S_t$  on the ergodic components of  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$  is ergodic; therefore, if  $K \neq \emptyset$ , then  $K = \Omega \times \mathbb{R}_+^* \text{ mod. } \nu \otimes \frac{1}{s^{1+\alpha}} ds$ .

Assume  $K = \Omega \times \mathbb{R}_+^*$ . This implies that the measure  $\mu \otimes \frac{1}{s^{1+\alpha}} ds$  is  $\sigma$ -finite on  $\mathcal{P}$ , and therefore  $\mathcal{P}$  is a factor of  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$ . Now

consider the quotient space  $(\Omega \times \mathbb{R}_+^*)/\mathcal{P}$  that we can endow, with a slight abuse of notation with the  $\sigma$ -algebra  $\mathcal{P}$ . Let  $\rho$  be the image measure of  $\mu \otimes \frac{1}{s^{1+\alpha}} ds$  by the projection map  $\pi$ . On  $((\Omega \times \mathbb{R}_+^*)/\mathcal{P}, \mathcal{P}, \rho)$   $\tilde{T}_\alpha$  and the dissipative flow  $S_t$  induce a transformation  $U$  and a dissipative flow  $Z_t$  that satisfy

$$\pi \circ \tilde{T}_\alpha = U \circ \pi, \pi \circ S_t = U \circ \pi \quad \text{and} \quad U \circ Z_t = Z_t \circ U.$$

Of course, thanks to Theorem 3,  $((\Omega \times \mathbb{R}_+^*)/\mathcal{P}, \mathcal{P}, \rho, U)$  is a Maharam system; therefore, we can represent it as  $(Y \times \mathbb{R}_+^*, \mathcal{K} \otimes \mathcal{B}_+, \sigma \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{L}_\alpha)$  for a non-singular system  $(Y, \mathcal{K}, \sigma, L)$ . Applying Lemma 4,  $\pi$  induces a nonsingular factor map  $\Gamma$  from  $(\Omega, \mathcal{G}, \mu, T)$  to  $(Y, \mathcal{K}, \sigma, L)$ , which means that there exists an  $R$ -invariant  $\sigma$ -algebra  $\mathcal{Z} \subset \mathcal{G}$  such that  $\Gamma^{-1}\mathcal{K} = \mathcal{Z}$ . But we can observe, that for all  $n > 0$ , the factor  $\tilde{T}_\alpha^{-n}\mathcal{G} \otimes \mathcal{B}_+$  corresponds to a Maharam system that corresponds to the factor  $T^{-n}\mathcal{G}$  of  $(\Omega, \mathcal{G}, \mu, T)$ . Therefore, for all  $n > 0$ ,  $\mathcal{Z} \subset T^{-n}\mathcal{G}$ , that is,  $\mathcal{Z} \subset \bigcap_{n \in \mathbb{N}} T^{-n}\mathcal{G} = \{\Omega, \emptyset\}$ . This means that  $\mathcal{K} = \{Y, \emptyset\}$ , or, in other words, that  $(Y, \mathcal{K}, \sigma, L)$  is the trivial (one-point) system.  $(Y \times \mathbb{R}_+^*, \mathcal{K} \otimes \mathcal{B}_+, \sigma \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{L}_\alpha)$  then possesses lots of invariant sets of positive finite measure, for example  $A := Y \times [1, 2]$ . But  $\pi^{-1}(A)$  is in turn a positive and finite measure invariant set for the system  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$ , and the existence of such set is impossible in a Maharam extension of an ergodic system which does not posses a finite  $T$ -invariant probability measure  $\nu \ll \mu$ . We can conclude that  $K = \emptyset$ .

To prove that  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$  is remotely infinite, it remains to show that  $\bigvee_{n \in \mathbb{Z}} \tilde{T}_\alpha^{-n}\mathcal{G} \otimes \mathcal{B}_+ = \mathcal{F} \otimes \mathcal{B}_+$ . We only sketch the proof which consists of verifying that the operation of taking natural extension and Maharam extension commute.

Of course, we have  $\bigvee_{n \in \mathbb{Z}} \tilde{T}_\alpha^{-n}\mathcal{G} \otimes \mathcal{B}_+ \subset \mathcal{F} \otimes \mathcal{B}_+$ . It is not difficult to check that  $\bigvee_{n \in \mathbb{Z}} \tilde{T}_\alpha^{-n}\mathcal{G} \otimes \mathcal{B}_+$  corresponds to a Maharam system that comes from a  $\sigma$ -algebra  $\mathcal{H} \subset \mathcal{F}$ . But we also have  $\mathcal{G} \subset \mathcal{H}$  and as  $T^{-1}\mathcal{H} = \mathcal{H}$ , we get  $\bigvee_{n \in \mathbb{Z}} T^{-n}\mathcal{G} \subset \mathcal{H}$ . By assumption,  $\bigvee_{n \in \mathbb{Z}} T^{-n}\mathcal{G} = \mathcal{F}$ , and we deduce  $\mathcal{H} = \mathcal{F}$  which implies  $\bigvee_{n \in \mathbb{Z}} \tilde{T}_\alpha^{-n}\mathcal{G} \otimes \mathcal{B}_+ = \mathcal{F} \otimes \mathcal{B}_+$ .  $\square$

As before we deduce the following result for  $\alpha$ -stable stationary processes:

**THEOREM 15.** *Let  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, \mathbb{P}, S)$  be a stationary stable process with associated system  $(\Omega, \mathcal{F}, \mu, T)$ . If  $(\Omega, \mathcal{F}, \mu, T)$  is  $K$  and not of type  $\text{II}_1$ , then  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, \mathbb{P}, S)$  is  $K$ .*

**PROOF.** From Proposition 14, we know that the Lévy measure system of the stable process is remotely infinite. The corresponding Poisson suspension is  $K$  by a result from [14]. By applying Maruyama’s representation Theorem (Theorem 6), we recover the stable process as a factor of the suspension, which therefore inherits the  $K$  property.  $\square$

Recall that in the probability preserving context,  $K$  is strictly stronger than mixing. In [11], to produce examples of mixing  $\alpha$ -stable stationary processes that were not based on dissipative nonsingular systems, the authors considered indeed null recurrent Markov chains as base systems. These systems are well-known examples of  $K$ -systems; therefore Theorem 15 shows that the associated  $\alpha$ -stable stationary processes are not just merely mixing but are indeed  $K$ .

7.3. *Rigidity.* We recall that a system  $(\Omega, \mathcal{F}, \mu, T)$  is rigid if there exists an increasing sequence  $n_k$  such that  $T^{n_k} \rightarrow \text{Id}$  in the group of nonsingular automorphism on  $(\Omega, \mathcal{F}, \mu)$  [the convergence being equivalent to the weak convergence in  $L^2(\mu)$  of the associated unitary operators  $U_{T^{n_k}} : f \mapsto \sqrt{\frac{dT_*^{-n_k}\mu}{d\mu}} f \circ T^{n_k}$  to the identity]. Observe that in the finite measure case, rigidity does not imply ergodicity but prevents mixing.

PROPOSITION 16. *The Maharam system  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds, \tilde{T}_\alpha)$  is rigid for the sequence  $n_k$  if and only  $(\Omega, \mathcal{F}, \mu, T)$  is rigid for the sequence  $n_k$ .*

PROOF. First observe that the map  $T \mapsto \tilde{T}_\alpha$  is a continuous group homomorphism from the group of nonsingular automorphism of  $(\Omega, \mathcal{F}, \mu)$  to the group of measure preserving automorphism of  $(\Omega \times \mathbb{R}_+^*, \mathcal{F} \otimes \mathcal{B}_+, \mu \otimes \frac{1}{s^{1+\alpha}} ds)$ . As  $T^{n_k} \rightarrow \text{Id}$ , then  $\tilde{T}_\alpha^{n_k} \rightarrow \text{Id}$ ; therefore  $\tilde{T}_\alpha^{n_k}$  is rigid for the sequence  $n_k$ .

Conversely, if  $\tilde{T}_\alpha$  is rigid for the same sequence, then, as

$$\langle U_{\tilde{T}_\alpha}^{n_k} f \otimes g, f \otimes g \rangle_{L^2(\mu \otimes 1/(s^{1+\alpha}) ds)} \leq \|g\|_2^2 \langle U_T^{n_k} f, f \rangle_{L^2(\mu)} \leq \|g\|_2^2 \|f\|_2^2$$

and  $\langle U_{\tilde{T}_\alpha}^{n_k} f \otimes g, f \otimes g \rangle_{L^2(\mu \otimes 1/(s^{1+\alpha}) ds)} \rightarrow \|g\|_2^2 \|f\|_2^2$ , we get  $\langle U_T^{n_k} f, f \rangle_{L^2(\mu)} \rightarrow \|f\|_2^2$ ; thus  $T$  is rigid.  $\square$

We need the following general result:

PROPOSITION 17. *A stationary IDp stationary process  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, \mathbb{P}, S)$  is rigid for the sequence  $n_k$  if and only if its Lévy measure system  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, Q, S)$  is rigid for the sequence  $n_k$ .*

PROOF. Consider  $X := \{X_n\}_{n \in \mathbb{Z}}$  where  $X_n := \{x_k\}_{k \in \mathbb{Z}} \mapsto x_n$  on  $\mathbb{R}^{\mathbb{Z}}$  and let  $\langle a, X \rangle$  be a finite linear combination of the coordinates.  $\exp i \langle a, X \rangle - \mathbb{E}[\exp i \langle a, X \rangle]$  is a centered square integrable vector under  $\mathbb{P}$  whose spectral measure (under  $\mathbb{P}$ ) is  $\lambda_a := |\mathbb{E}[\exp i \langle a, X \rangle]|^2 \sum_{k=1}^\infty \frac{1}{k!} \sigma_a^{*k}$  where  $\sigma_a$  is the spectral measure of  $\exp i \langle a, X \rangle - 1$  under  $Q$  (see [13]). Therefore  $\widehat{\sigma}_a(n_k) \rightarrow \widehat{\sigma}_a(0)$  if and only if  $\widehat{\lambda}_a(n_k) \rightarrow \widehat{\lambda}_a(0)$ . This implies that  $\exp i \langle a, X \rangle - \mathbb{E}[\exp i \langle a, X \rangle]$  is a rigid vector for  $n_k$  under  $\mathbb{P}$  if and only if  $\exp i \langle a, X \rangle - 1$  is a rigid vector for  $n_k$  under  $Q$ . Observe now that the smallest  $\sigma$ -algebra generated by vectors of the kind

$\exp i\langle a, X \rangle - \mathbb{E}[\exp i\langle a, X \rangle]$  under  $\mathbb{P}$  is  $\mathcal{B}^{\otimes \mathbb{Z}}$ , and the same is true with vectors of the kind  $\exp i\langle a, X \rangle - 1$  under  $Q$ . As in any dynamical system if there exists a rigid vector for the sequence  $n_k$ , there exists a nontrivial factor which is rigid for the sequence  $n_k$ , we get the announced result.  $\square$

**THEOREM 18.** *A stationary stable process  $(\mathbb{R}^{\mathbb{Z}}, \mathcal{B}^{\otimes \mathbb{Z}}, \mathbb{P}, S)$  with associated system  $(\Omega, \mathcal{F}, \mu, T)$  is rigid for the sequence  $n_k$  if and only  $(\Omega, \mathcal{F}, \mu, T)$  is rigid for the sequence  $n_k$ .*

**PROOF.** This is the combination of the last two results.  $\square$

## REFERENCES

- [1] AARONSON, J. (1997). *An Introduction to Infinite Ergodic Theory. Mathematical Surveys and Monographs* **50**. Amer. Math. Soc., Providence, RI. [MR1450400](#)
- [2] AARONSON, J., LEMAŃCZYK, M. and VOLNÝ, D. (1998). A cut salad of cocycles. *Fund. Math.* **157** 99–119. [MR1636882](#)
- [3] DANILENKO, A. I. and SILVA, C. E. Ergodic theory: Nonsingular transformations. Preprint. Available at <http://dblp.uni-trier.de/rec/bibtex/reference/complexity/DanilenkoS09>.
- [4] HAJIAN, A., ITO, Y. and KAKUTANI, S. (1972). Invariant measures and orbits of dissipative transformations. *Adv. Math.* **9** 52–65. [MR0302860](#)
- [5] HARDIN, C. D. JR. (1982). On the spectral representation of symmetric stable processes. *J. Multivariate Anal.* **12** 385–401. [MR0666013](#)
- [6] KATZNELSON, Y. and WEISS, B. (1991). The classification of nonsingular actions, revisited. *Ergodic Theory Dynam. Systems* **11** 333–348. [MR1116644](#)
- [7] KRENGEL, U. (1970). Transformations without finite invariant measure have finite strong generators. In *Contributions to Ergodic Theory and Probability (Proc. Conf., Ohio State Univ., Columbus, Ohio, 1970)* 133–157. Springer, Berlin. [MR0269808](#)
- [8] MARUYAMA, G. (1970). Infinitely divisible processes. *Theory Probab. Appl.* **15** 1–22.
- [9] ROSIŃSKI, J. (1995). On the structure of stationary stable processes. *Ann. Probab.* **23** 1163–1187. [MR1349166](#)
- [10] ROSIŃSKI, J. (2006). Minimal integral representations of stable processes. *Probab. Math. Statist.* **26** 121–142. [MR2301892](#)
- [11] ROSIŃSKI, J. and SAMORODNITSKY, G. (1996). Classes of mixing stable processes. *Bernoulli* **2** 365–377. [MR1440274](#)
- [12] ROY, E. (2005). Mesures de Poisson, infinie divisibilité et propriétés ergodiques. Ph.D. thesis, Univ. Paris 6.
- [13] ROY, E. (2007). Ergodic properties of Poissonian ID processes. *Ann. Probab.* **35** 551–576. [MR2308588](#)
- [14] ROY, E. (2009). Poisson suspensions and infinite ergodic theory. *Ergodic Theory Dynam. Systems* **29** 667–683. [MR2486789](#)
- [15] SAMORODNITSKY, G. (2005). Null flows, positive flows and the structure of stationary symmetric stable processes. *Ann. Probab.* **33** 1782–1803. [MR2165579](#)
- [16] SAMORODNITSKY, G. and TAQQU, M. S. (1994). *Stable Non-Gaussian Random Processes: Stochastic Models with Infinite Variance*. Chapman & Hall, New York. [MR1280932](#)
- [17] SATO, K.-I. (1999). *Lévy Processes and Infinitely Divisible Distributions. Cambridge Studies in Advanced Mathematics* **68**. Cambridge Univ. Press, Cambridge. [MR1739520](#)
- [18] SCHMIDT, K. (1977). *Cocycles on Ergodic Transformation Groups. Macmillan Lectures in Mathematics* **1**. Macmillan Company of India, Ltd., Delhi. [MR0578731](#)

- [19] SILVA, C. E. and THIEULLEN, P. (1995). A skew product entropy for nonsingular transformations. *J. Lond. Math. Soc.* (2) **52** 497–516. [MR1363816](#)

LABORATOIRE ANALYSE GÉOMÉTRIE ET APPLICATIONS  
UMR 7539  
UNIVERSITÉ PARIS 13  
99 AVENUE J.-B. CLÉMENT  
F-93430 VILLETANEUSE  
FRANCE  
E-MAIL: [roy@math.univ-paris13.fr](mailto:roy@math.univ-paris13.fr)