

# On the ranked excursion heights of a Kiefer process

by

Endre CSÁKI<sup>1</sup> and Yueyun HU<sup>2</sup>

**Abstract:** Let  $(K(s, t), 0 \leq s \leq 1, t \geq 1)$  be a Kiefer process, i.e. a continuous two-parameter centered Gaussian process indexed by  $[0, 1] \times \mathbb{R}_+$  whose covariance function is given by  $\mathbb{E}(K(s_1, t_1)K(s_2, t_2)) = (s_1 \wedge s_2 - s_1 s_2) t_1 \wedge t_2$ ,  $0 \leq s_1, s_2 \leq 1, t_1, t_2 \geq 0$ . For each  $t > 0$ , the process  $K(\cdot, t)$  is a Brownian bridge on the scale of  $\sqrt{t}$ . Let  $M_1^*(t) \geq M_2^*(t) \geq \dots M_j^*(t) \geq \dots 0$  be the ranked excursion heights of  $K(\cdot, t)$ . In this paper, we study the path properties of the process  $t \rightarrow M_j^*(t)$ . Two laws of the iterated logarithm are established to describe the asymptotic behaviors of  $M_j^*(t)$  as  $t$  goes to infinity.

**Keywords.** Kiefer process, excursions, ranked heights.

**AMS Classification 2000.** 60F15, 60G55.

---

<sup>1</sup>A. Rényi Institute of Mathematics, Hungarian Academy of Sciences, Reáltanoda u. 13–15, P.O.B. 127, Budapest, H–1364, Hungary. E-mail: csaki@renyi.hu. Research supported by the Hungarian National Foundation for Scientific Research, Grants T 029621 and T 037886

<sup>2</sup>Laboratoire de Probabilités et Modèles Aléatoires (CNRS UMR–7599), Université Paris VI, 4 Place Jussieu, F–75252 Paris cedex 05, France. E-mail: hu@ccr.jussieu.fr

# 1 Introduction

Let  $\{B(t), t \geq 0\}$  be a standard one-dimensional Brownian motion, i.e. a continuous centered Gaussian process with covariance

$$\mathbb{E}\left(B(t_1)B(t_2)\right) = t_1 \wedge t_2, \quad t_1, t_2 \geq 0.$$

We consider also standard Brownian bridge  $\{p(s), 0 \leq s \leq 1\}$ , i.e. a centered Gaussian process with covariance

$$\mathbb{E}(p(s_1)p(s_2)) = s_1 \wedge s_2 - s_1s_2, \quad 0 \leq s_1, s_2 \leq 1.$$

It is well-known that almost all sample paths of  $B$  consists of countable many zero-free intervals called excursions. Let  $(a, b)$  an excursion interval, i.e.  $B(a) = B(b) = 0$  and either  $B(s) > 0, a < s < b$  called positive excursion, or  $B(s) < 0, a < s < b$  called negative excursion. The height of excursion is defined by either

$$H \stackrel{\text{def}}{=} \max_{a \leq s \leq b} B(s)$$

or

$$H^* \stackrel{\text{def}}{=} \max_{a \leq s \leq b} |B(s)|.$$

Clearly,  $H > 0$  holds only for positive excursions. Pitman and Yor [11] introduced the ranked heights of excursions up to time  $t$ : let

$$H_1(t) \geq H_2(t) \geq \dots H_j(t) \geq \dots$$

and

$$H_1^*(t) \geq H_2^*(t) \geq \dots H_j^*(t) \geq \dots$$

be the heights of positive and all excursions respectively, of  $\{B(s), 0 \leq s \leq t\}$ , including the meander heights  $\sup_{g_t \leq s \leq t} B(s)$  and  $\sup_{g_t \leq s \leq t} |B(s)|$ , where  $g_t$  denotes the last zero of  $B$  before  $t$ . The ranked heights of excursions of  $p$  can be defined similarly.

Let furthermore  $\{K(s, t), 0 \leq s \leq 1, t \geq 0\}$  be a Kiefer process, i.e. a continuous two-parameter centered Gaussian process indexed by  $[0, 1] \times \mathbb{R}_+$  whose covariance function is given by

$$\mathbb{E}\left(K(s_1, t_1)K(s_2, t_2)\right) = (s_1 \wedge s_2 - s_1s_2)t_1 \wedge t_2, \quad 0 \leq s_1, s_2 \leq 1, t_1, t_2 \geq 0.$$

Kiefer [7] introduced this process  $K$  to approximate the empirical process. See Csörgő and Révész [4] for detailed studies and related references on Kiefer process and on the invariance

principle between empirical process and Kiefer process. Note that for fixed  $t > 0$ , the process  $s \in [0, 1] \rightarrow \frac{K(s,t)}{\sqrt{t}}$  is a standard Brownian bridge. Denote by

$$M_1(t) \geq M_2(t) \geq \dots \geq M_j(t) \geq \dots$$

the ranked heights of the positive excursions of the Brownian bridge  $K(\cdot, t)$  over the whole time interval  $[0, 1]$ . Denote by

$$M_1^*(t) \geq M_2^*(t) \geq \dots \geq M_j^*(t) \geq \dots$$

the ranked heights of the excursions of  $|K(\cdot, t)|$ . By scaling properties, the distributions of  $(\frac{M_j(t)}{\sqrt{t}}, j \geq 1)$  and  $(\frac{M_j^*(t)}{\sqrt{t}}, j \geq 1)$  are the same as that of the ranked excursion heights of a standard Brownian bridge. See Pitman and Yor [12] for studies on these distribution.

We are interested in the path properties of the processes  $t \rightarrow M_j(t)$  and  $t \rightarrow M_j^*(t)$ . In particular, we aim at the asymptotic behaviors of  $M_j(t)$  and  $M_j^*(t)$  as  $t \rightarrow \infty$ .

Observe that  $M_1(t) = \sup_{0 \leq s \leq 1} K(s, t)$  and  $M_1^*(t) = \sup_{0 \leq s \leq 1} |K(s, t)|$ . The following laws of the iterated logarithm are known, see respectively Csörgő and Révész ([4], pp. 81), Mogul'skii [8] and Csáki and Shi [3]:

**Theorem A** ([4], [8], [3]). *We have*

$$\limsup_{t \rightarrow \infty} \frac{M_1^*(t)}{\sqrt{t \log \log t}} = \frac{1}{\sqrt{2}}, \quad \text{a.s.} \quad (1.1)$$

$$\liminf_{t \rightarrow \infty} \sqrt{\frac{\log \log t}{t}} M_1^*(t) = \frac{\pi}{\sqrt{8}}, \quad \text{a.s.} \quad (1.2)$$

$$\liminf_{t \rightarrow \infty} \frac{(\log t)^\chi}{\sqrt{t}} M_1(t) = \begin{cases} 0 & \text{if } \chi \leq \frac{1}{2} \\ \infty & \text{if } \chi > \frac{1}{2} \end{cases} \quad \text{a.s.} \quad (1.3)$$

In (1.1) we may replace  $M_1^*(t)$  by  $M_1(t)$ .

The almost sure behavior of  $H_j^*(t)$  was studied in Csáki and Hu [2]:

**Theorem B** ([2]). *We have*

$$\limsup_{t \rightarrow \infty} \frac{H_j^*(t)}{\sqrt{t \log \log t}} = \frac{\sqrt{2}}{2j-1}, \quad \text{a.s.} \quad j \geq 1 \quad (1.4)$$

$$\liminf_{t \rightarrow \infty} \frac{(\log t)^\chi}{\sqrt{t}} H_j^*(t) = \begin{cases} 0 & \text{if } \chi \leq 1 \\ \infty & \text{if } \chi > 1 \end{cases} \quad \text{a.s.} \quad j \geq 2. \quad (1.5)$$

A natural question is to ask what happens with  $(M_j^*(t), t \geq 0)$  for  $j \geq 2$ . As a process indexed by  $t$ , the  $j$ -highest heights  $M_j^*(t)$  may share some unusual properties different from  $M_1^*(t)$ . For instance,  $t \rightarrow M_j^*(t)$  is not continuous for  $j \geq 2$  in contrast with the continuity of  $M_1^*(\cdot)$ .

**Theorem 1.1** Fix  $j \geq 1$ . We have

$$\limsup_{t \rightarrow \infty} \frac{M_j^*(t)}{\sqrt{t \log \log t}} = \frac{1}{j\sqrt{2}}, \quad \text{a.s.}$$

The same result remains true when  $M_j^*(t)$  is replaced by  $M_j(t)$ .

It is also of interest to find the liminf behavior of  $M_j(\cdot)$ :

**Theorem 1.2** Fix  $j \geq 2$ . We have

$$\liminf_{t \rightarrow \infty} \frac{(\log t)^\chi}{\sqrt{t}} M_j^*(t) = \begin{cases} 0 & \text{if } \chi \leq \frac{1}{2} \\ \infty & \text{if } \chi > \frac{1}{2} \end{cases} \quad \text{a.s.}$$

The same result remains true when we replace  $M_j^*(t)$  by  $M_j(t)$ .

Comparing (1.2) with Theorem 1.2, we can see that the liminf behaviors of  $M_1^*$  and  $M_j^*$  ( $j > 1$ ) are completely different.

The proof of Theorem 1.1 is based on an estimate on the downcrossings of a Brownian bridge, this estimate will be given in Section 2. To show Theorem 1.2, a usual way would be to estimate  $\mathbb{P}(\inf_{1 \leq t \leq 2} M_j(t) < \epsilon)$  as  $\epsilon$  goes to 0. This problem remains open to our best knowledge. To overcome this difficulty, we shall adopt the method of Csáki and Shi [3], which consists of reducing the problem for the Kiefer process to that for an Ornstein-Uhlenbeck process. Section 2 also contains several preliminary results to complete the proofs of Theorems 1.1 and 1.2, which will be presented respectively in Sections 3 and 4.

Throughout this paper,  $(C_k, 1 \leq k \leq 6)$  denote some positive constants whose exact values are unimportant.

## 2 Downcrossings

Consider a continuous function  $f : I = [a, b] \rightarrow \mathbb{R}$  with  $a, b \in \mathbb{R}$ . For two real numbers  $x < y$ , we define inductively

$$\alpha_1 = \alpha_1(y) \stackrel{\text{def}}{=} \inf\{v \geq a : f(v) \geq y\}, \quad (2.1)$$

$$\beta_k = \beta_k(x) \stackrel{\text{def}}{=} \inf\{v \geq \alpha_k : f(v) \leq x\}, \quad k \geq 1, \quad (2.2)$$

$$\alpha_k = \alpha_k(y) \stackrel{\text{def}}{=} \inf\{v \geq \beta_{k-1} : f(v) \geq y\}, \quad k \geq 2, \quad (2.3)$$

with the convention  $\inf \emptyset = \infty$ . Define the number of downcrossings of  $(x, y)$  by  $f$  during the time interval  $I$  as

$$D_f(x, y; I) = \sup\{k : \alpha_k(y) \leq b\}. \quad (2.4)$$

We adopt the above definition of downcrossings, which is slightly different from the usual one, to keep the following equivalence:

$$\sup_{v \in I} f(v) \geq y \quad \iff \quad D_f(x, y; I) \geq 1.$$

Remark that the condition  $\{D_f(x, y; I) \geq 1\}$  does not depend on  $x$ . In the following two subsections, we shall discuss respectively the numbers of downcrossings by a standard Brownian motion, a Brownian bridge and by an Ornstein-Uhlenbeck process.

## 2.1 Brownian motion and Brownian bridge

Let  $\{B(s), s \geq 0\}$  be a standard Brownian motion and let  $\{p(s), 0 \leq s \leq 1\}$  be a standard Brownian bridge from 0 to 0. First, we present a preliminary result based on the reflection principle.

**Lemma 2.1** *Fix  $j \geq 1$  and  $\max(x, 0) < y$ . We have*

$$\mathbb{P}\left(D_B(x, y; [0, 1]) \geq j, B(1) \in dz\right) = \begin{cases} \varphi(2jy - 2(j-1)x - z)dz & \text{if } z \leq y, \\ \varphi(2(j-1)y - 2(j-1)x + z)dz & \text{if } z > y, \end{cases} \quad (2.5)$$

where  $\varphi$  is the standard normal density function.

**Proof:** We use the reflection principle formulated by (cf., e.g. [5])

**Fact 2.2** *Let  $\{B(s), s \geq 0\}$  be a standard Brownian motion and let  $\tau$  be a stopping time for  $B$ . Then*

$$B^{(\tau)}(s) \stackrel{\text{def}}{=} \begin{cases} B(s) & \text{if } 0 \leq s \leq \tau \\ 2B(\tau) - B(s) & \text{if } \tau \leq s \end{cases}$$

is also a standard Brownian motion.

Let us make use of the stopping times  $\alpha_k = \alpha_k(y)$  and  $\beta_k = \beta_k(x)$  introduced in (2.1)–(2.3), corresponding to  $f(t) = B(t)$ ,  $I = [0, 1]$ .

Our Lemma 2.1 is well-known for  $j = 1$ .

We illustrate the proof in the simple case  $j = 2$ , using the reflection principle subsequently for our stopping times. Let  $\{B(s), 0 \leq s \leq 1\}$  be a Brownian motion such that  $\alpha_2 < 1$  and  $B(1) = z \leq y$ . Then by Fact 2.2,  $B_1(s) \stackrel{\text{def}}{=} B^{(\alpha_1)}(s)$ ,  $0 \leq s \leq 1$  is a Brownian motion with  $B_1(1) = 2y - z$ ,  $\beta_1$  is its first hitting time of  $2y - x$  and  $\alpha_2$  is its first hitting time of  $y$  after

$\beta_1$ . In the next step consider  $B_2(s) \stackrel{\text{def}}{=} B_1^{(\beta_1)}(s)$ ,  $0 \leq s \leq 1$ . Then  $B_2(1) = 2y - 2x + z$ , and  $\alpha_2$  is its first hitting time of  $3y - 2x$ . Finally, consider  $B_3(s) = B_2^{(\alpha_2)}(s)$ ,  $0 \leq s \leq 1$  for which we have  $B_3(1) = 4y - 2x - z$ . By reversing this procedure, starting from a Brownian motion with endpoints  $4y - 2x - z$  at  $s = 1$ , we get a Brownian motion with  $\alpha_1 < 1$  and  $B(1) = z$ . This proves the first equality of (2.5) in the case  $j = 2$ . The procedure is similar for  $z > y$ , except that we stop with  $B_2$ , so the last reflection (at  $\alpha_2$ ) is not performed. Using this idea in obvious manner for the general case  $j > 2$ , yields our lemma.  $\blacksquare$

Since a Brownian bridge  $\{p(s), 0 \leq s \leq 1\}$  is a Brownian motion conditioned to  $B(1) = 0$ , we have the following

**Corollary 2.3** *For  $j \geq 1$  and  $\max(x, 0) < y$ , we have*

$$\mathbb{P}\left(D_p(x, y; [0, 1]) \geq j\right) = \exp\left(-2(jy - (j-1)x)^2\right).$$

**Proof:** Putting  $z = 0$  in (2.5) we get

$$\mathbb{P}\left(D_p(x, y; [0, 1]) \geq j\right) = \frac{\varphi(2jy - 2(j-1)x)}{\varphi(0)} = \exp\left(-2(jy - (j-1)x)^2\right).$$

$\blacksquare$

Taking  $x = 0$ , we recover Pitman and Yor [12]'s formula for the distribution of  $M_j(1)$ :

$$\mathbb{P}\left(M_j(1) > y\right) = \mathbb{P}\left(D_p(0, y; [0, 1]) \geq j\right) = \exp\left(-2j^2y^2\right). \quad (2.6)$$

Another corollary can be obtained by taking  $x = 0$  and integrating with respect to  $z$ :

**Corollary 2.4** *For  $j \geq 1$ ,  $y > 0$  we have*

$$\mathbb{P}\left(H_j(1) > y\right) = 2(1 - \Phi((2j-1)y)),$$

where  $\Phi$  is the standard normal distribution function, and  $H_j(1)$  denotes the height of the  $j$ -th highest positive Brownian excursion up to time 1.

Now we present an estimate on  $\sup_{0 \leq t \leq T} M_j^*(t)$ .

**Proposition 2.5** *Fix  $j \geq 2$ . There exists some constant  $C_1 > 1$  such that for all  $u > 0$  and  $\lambda \geq \sqrt{u}$ , we have*

$$\mathbb{P}\left(\sup_{0 \leq t \leq u} M_j^*(t) > \lambda\right) \leq C_1 \exp\left(-2\left(\frac{j\lambda}{\sqrt{u}} - \frac{2j-1}{2}\right)^2\right)$$

In the proof of Proposition 2.5, we need the following lemma:

**Lemma 2.6** For  $0 \leq x < y$ ,  $j \geq 1$ , we have

$$\mathbb{P}\left(D_{|p|}(x, y; [0, 1]) \geq j\right) \leq 2^j \mathbb{P}\left(D_p(x, y; [0, 1]) \geq j\right).$$

**Proof of Lemma 2.6:** Again, we present the proof for  $j = 2$ . Upcrossings from  $x$  to  $y$  by  $|p|$  are either upcrossings by  $p$  from  $x$  to  $y$  or downcrossings by  $p$  from  $-x$  to  $-y$ . Define the following events:

$$\begin{aligned} A^{++} &\stackrel{\text{def}}{=} \{\text{There are at least two upcrossings by } p \text{ from } x \text{ to } y\} \\ A^{+-} &\stackrel{\text{def}}{=} \{\text{There is at least one downcrossing by } p \text{ from } -x \text{ to } -y \\ &\quad \text{after an upcrossing by } p \text{ from } x \text{ to } y\} \\ A^{-+} &\stackrel{\text{def}}{=} \{\text{There is at least one upcrossing by } p \text{ from } x \text{ to } y \\ &\quad \text{after a downcrossing by } p \text{ from } -x \text{ to } -y\} \\ A^{--} &\stackrel{\text{def}}{=} \{\text{There are at least two downcrossings by } p \text{ from } -x \text{ to } -y\}. \end{aligned}$$

Obviously

$$\mathbb{P}\left(D_{|p|}(x, y; [0, 1]) \geq 2\right) \leq \mathbb{P}(A^{++}) + \mathbb{P}(A^{+-}) + \mathbb{P}(A^{-+}) + \mathbb{P}(A^{--})$$

and by symmetry,  $\mathbb{P}(A^{++}) = \mathbb{P}(A^{--})$ ,  $\mathbb{P}(A^{+-}) = \mathbb{P}(A^{-+})$ . Moreover,  $\mathbb{P}(A^{+-}) \leq \mathbb{P}(A^{++})$ , since by Corollary 2.3 we have

$$\mathbb{P}(A^{++}) = \exp(-2(2y - x)^2)$$

and an argument, similar to the proof of Lemma 2.1 shows that

$$\mathbb{P}(A^{+-}) = \exp(-8y^2).$$

Hence,

$$\mathbb{P}\left(D_{|p|}(x, y; [0, 1]) \geq 2\right) \leq 4\mathbb{P}(A^{++}) = 2^2 \exp(-2(2y - x)^2),$$

proving Lemma 2.6 for  $j = 2$ . Extension of the above argument in an obvious manner for  $j > 2$ , proves our Lemma 2.6.  $\blacksquare$

Now we proceed with the proof of Proposition 2.5.

**Proof of Proposition 2.5:** For  $t > 0$ , we define  $\sigma_0^{(t)}(0) = 0$  and for  $i \geq 1$ ,

$$\begin{aligned} \tau_i^{(t)}(x) &\stackrel{\text{def}}{=} \inf\{s \geq \sigma_{i-1}^{(t)}(0) : |K(s, t)| = x\}, \\ \sigma_i^{(t)}(0) &\stackrel{\text{def}}{=} \inf\{s \geq \tau_i^{(t)}(x) : K(s, t) = 0\}, \end{aligned}$$

(write  $\tau_i^{(t)}(x) = 1$  if such  $s$  does not exist). Therefore,

$$\mathbb{P}\left(\sup_{0 \leq t \leq u} M_j^*(t) > \lambda\right) = \mathbb{P}\left(\exists t \in [0, u] : \tau_j^{(t)}(\lambda) < 1\right) = \mathbb{P}\left(\Theta \leq u\right),$$

where we define  $\Theta \stackrel{\text{def}}{=} \inf\{t \geq 0 : M_j^*(t) > \lambda\}$ . Let  $\mathcal{F}_t \stackrel{\text{def}}{=} \sigma\{K(s, u), 0 \leq s \leq 1, 0 \leq u \leq t\}$ , the sigma-algebra generated by  $\{K(s, u), 0 \leq s \leq 1, 0 \leq u \leq t\}$ . Then  $\Theta$  is a stopping time with respect to  $(\mathcal{F}_t)$ . Notice that the process  $t \rightarrow (K(\cdot, \Theta + t) - K(\cdot, \Theta))$  is independent of  $\mathcal{F}_\Theta$  and has the same law as  $(K(\cdot, t), t \geq 0)$ . Using the self similarity:  $K(\cdot, v + \Theta) - K(\cdot, \Theta) \stackrel{\text{law}}{=} \sqrt{v}K(\cdot, 1)$  for any fixed  $v > 0$ , we get

$$\mathbb{P}\left(\sup_{0 \leq s \leq 1} |K(s, u) - K(s, \Theta)| < \frac{\sqrt{u}}{2} \mid \Theta \leq u\right) \geq \mathbb{P}\left(\sup_{0 \leq s \leq 1} |K(s, 1)| < \frac{1}{2}\right) \stackrel{\text{def}}{=} \frac{2^j}{C_1} > 0.$$

Denote by

$$E_1 \stackrel{\text{def}}{=} \left\{ \sup_{0 \leq s \leq 1} |K(s, u) - K(s, \Theta)| < \frac{\sqrt{u}}{2} \right\} \cap \left\{ \Theta \leq u \right\},$$

we have shown that

$$\mathbb{P}\left(\Theta \leq u\right) \leq 2^{-j} C_1 \mathbb{P}\left(E_1\right).$$

On  $E_1$ , we can decompose  $K(s, u) = K(s, \Theta) + \widehat{K}(s)$  with  $\sup_{0 \leq s \leq 1} |\widehat{K}(s)| \leq \frac{\sqrt{u}}{2}$ . Since  $|K(\tau_i^{(\Theta)}(\lambda), \Theta)| = \lambda$  and  $K(\sigma_i^{(\Theta)}(0), \Theta) = 0$  for  $1 \leq i \leq j$ , it follows that for such random times  $0 < s_1 \stackrel{\text{def}}{=} \tau_1^{(\Theta)}(\lambda) < v_1 \stackrel{\text{def}}{=} \sigma_1^{(\Theta)}(0) < \dots < s_j \stackrel{\text{def}}{=} \tau_j^{(\Theta)}(\lambda) < 1$ , we have respectively,

$$|K(s_1, u)| \geq \lambda - \frac{\sqrt{u}}{2}, |K(v_1, u)| \leq \frac{\sqrt{u}}{2}, \dots, |K(s_j, u)| \geq \lambda - \frac{\sqrt{u}}{2}.$$

Namely, we have

$$E_1 \subset \left\{ D_{|K(\cdot, u)|} \left( \frac{\sqrt{u}}{2}, \lambda - \frac{\sqrt{u}}{2}; [0, 1] \right) \geq j \right\}.$$

It follows from scaling, Corollary 2.3 and Lemma 2.6 that

$$\begin{aligned} \mathbb{P}\left(E_1\right) &\leq \mathbb{P}\left(D_{|K(\cdot, u)|} \left( \frac{\sqrt{u}}{2}, \lambda - \frac{\sqrt{u}}{2}; [0, 1] \right) \geq j\right) \\ &= \mathbb{P}\left(D_{|p|} \left( \frac{1}{2}, \frac{\lambda}{\sqrt{u}} - \frac{1}{2}; [0, 1] \right) \geq j\right) \leq 2^j \exp\left(-2 \frac{(j\lambda - (2j-1)\frac{\sqrt{u}}{2})^2}{u}\right), \end{aligned}$$

proving the result. ■



## 2.2 Ornstein-Uhlenbeck process

Let us consider a stationary Ornstein-Uhlenbeck process  $(U(t), t \geq 0)$  with parameter  $\frac{1}{2}$ , which is a stationary centered Gaussian process with covariance  $\mathbb{E}(U(t)U(s)) = e^{-\frac{|t-s|}{2}}$ . We mention a paper by Pitman and Yor [10] for the study of distributions of excursion lengths of  $U$ .

Recall some known facts on the hitting times of  $U$ . Fix  $-\infty \leq z_1 < z_2 \leq \infty$  and define

$$\sigma(z_1, z_2) = \inf\{s \geq 0 : U(s) \notin [z_1, z_2]\}$$

to be the first exit time from the interval  $[z_1, z_2]$ . Consider the Sturm-Liouville equation:

$$\frac{1}{2}\phi''(x) - \frac{x}{2}\phi'(x) = -\lambda\phi(x), \quad x \in (z_1, z_2); \quad \phi(z_i) = 0 \text{ if } |z_i| < \infty, \quad i = 1, 2.$$

**Fact 2.7** ([14], [6], [9]) *Assume that  $\min(|z_1|, |z_2|) < \infty$ . There is a sequence of simple eigenvalues  $0 < \lambda_1(z_1, z_2) < \dots < \lambda_n(z_1, z_2) < \dots$  whose corresponding eigenfunctions  $\psi_1(z_1, z_2; x), \dots, \psi_n(z_1, z_2; x), \dots$  form a complete orthonormal system with respect to  $m(dx) = e^{-x^2/2}dx$ . The function  $(z_1, z_2) \rightarrow \lambda_1(z_1, z_2)$  is strictly positive and jointly continuous on  $\Xi = \{(z_1, z_2) \in [-\infty, \infty]^2 : z_1 < z_2, \min(|z_1|, |z_2|) < \infty\}$ , strictly increasing in  $z_1 \in (-\infty, z_2]$  for  $z_2 \leq \infty$  and strictly decreasing in  $z_2 \in [z_1, \infty)$  for  $z_1 \geq -\infty$ :*

$$\lambda_1(-\infty, 0) = \lambda_1(0, \infty) = \frac{1}{2}, \quad \lim_{(z_1, z_2) \rightarrow 0} \lambda_1(z_1, z_2) = \infty, \quad \lim_{(z_1, z_2) \rightarrow (-\infty, \infty)} \lambda_1(z_1, z_2) = 0.$$

**Fact 2.8** ([14], [6], [9], [1], [3]) *Assume that  $\min(|z_1|, |z_2|) < \infty$ . There exists some constant  $C_2 > 0$  such that uniformly on  $x \in \mathbb{R}$ ,*

$$\mathbb{P}\left(\sigma(z_1, z_2) > t \mid U(0) = x\right) = e^{-\lambda_1(z_1, z_2)t} \left(\theta(z_1, z_2)\psi_1(z_1, z_2; x) + r(t, x)\right),$$

where  $\theta(z_1, z_2) = \int_{z_1}^{z_2} \psi_1(z_1, z_2; x)m(dx)$  and

$$|r(t, x)| \leq C_2 \exp\left(\frac{x^2}{9} - \frac{t}{2}\right).$$

When  $z_1 = -z_2 = -z$  with  $z > 0$ , we get

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}\left(\sup_{0 \leq s \leq t} |U(s)| < z\right) = -\lambda_1(-z, z). \quad (2.7)$$

Moreover,  $\lim_{z \rightarrow \infty} \lambda_1(-z, z) = 0$ .

We shall need the probability that the process  $U$  downcrosses a given interval  $(z_1, z_2)$  only a few times during  $[-t, t]$ . This is stated in the following lemma:

**Lemma 2.9** *Fix  $-\infty < z_1 < z_2 < \infty$  and  $k \geq 1$ ; We have*

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}\left(D_U(z_1, z_2; [-t, t]) \leq k\right) = \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}\left(D_U(z_1, z_2; [0, 2t]) \leq k\right) = -2\mu(z_1, z_2),$$

where  $\mu(z_1, z_2) \stackrel{\text{def}}{=} \min(\lambda_1(-\infty, z_2), \lambda_1(z_1, \infty)) > 0$ . Moreover, we have

$$\lim_{z_1, z_2 \rightarrow 0} \mu(z_1, z_2) = \frac{1}{2}.$$

Note that the constant  $k$ , arbitrary but fixed, does not influence the rate of exponential decay of the two probability terms in the above lemma.

**Proof:** The above first equality is due to the stationarity of the Ornstein-Uhlenbeck process. Using again the stopping times  $\alpha_j$  and  $\beta_j$  defined in (2.1)–(2.3) associated with  $a = 0$ ,  $b = 2t$ ,  $x = z_1$ ,  $y = z_2$ ,  $I = [0, 2t]$  and  $f(v) = U(v)$ , we have  $\mathbb{P}\left(D_U(z_1, z_2; [0, 2t]) \leq k\right) = \mathbb{P}\left(\alpha_{k+1} > 2t\right)$ .

Remark that  $\alpha_1 = \inf\{s \geq 0 : U(s) \geq z_2\} = \sigma(-\infty, z_2)$ . The strong Markov property implies that the random variables of the family  $\{\alpha_1, \beta_j - \alpha_j, \alpha_{j+1} - \beta_j, j \geq 1\}$  are mutually independent. Furthermore,  $\beta_1 - \alpha_1 = \sigma(z_1, \infty) \circ \theta_{\alpha_1}$  where  $\theta$  is the usual shift operator. And for  $j \geq 2$ ,  $\beta_j - \alpha_j$  (resp:  $\alpha_j - \beta_{j-1}$ ) has the same law as  $T_{z_2 \rightarrow z_1}$  (resp:  $T_{z_1 \rightarrow z_2}$ ), where  $T_{x \rightarrow y}$  denotes the hitting time of  $y$  by an Ornstein-Uhlenbeck process starting from  $x$ . Based on Fact 2.8, simple convolution computation yields that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}\left(\alpha_{k+1} > t\right) = -\mu(z_1, z_2),$$

and the desired conclusion follows. ■

## 2.3 A technical lemma

Recall that  $\{p(s), 0 \leq s \leq 1\}$  denotes a standard Brownian bridge. Let  $0 \leq y < z/4$  and consider the event

$$G_{y,z} = \left\{ \exists 0 < a_1 < c_1 < b_1 < a_2 < c_2 < b_2 < 1 : \right. \\ \left. |p(a_i)| \leq y, |p(b_i)| \leq y, |p(c_i)| \geq z, i = 1, 2 \right\} \quad (2.8)$$

Remark that  $G_{y,z} \supset G_{0,z}$  and that  $G_{0,z}$  is in fact the event that the height of the second highest excursion of  $|p(\cdot)|$  is larger than  $z$ . We shall need to bound  $\mathbb{P}(G_{y,z})$  in the proof of the upper bound of Theorem 1.2.

**Lemma 2.10** *There exists an absolute constant  $C_3 > 0$  such that for all  $0 \leq y < \frac{z}{4}$  and  $0 < z < \frac{1}{2}$ ,*

$$\mathbb{P}(G_{y,z}) \leq 1 - C_3 z^2.$$

We note that this estimate is nearly sharp, since we can also obtain a lower bound from (2.6) as follows:

$$\mathbb{P}(G_{y,z}) \geq \mathbb{P}(G_{0,z}) = \mathbb{P}(M_2^*(1) \geq z) \geq \mathbb{P}(M_2(1) \geq z) \geq 1 - C_4 z^2, \quad 0 < z < \frac{1}{2}.$$

**Proof of Lemma 2.10:** Fix  $(y, z)$  such that  $0 \leq y < \frac{z}{4}$  and  $0 < z < \frac{1}{2}$ . Define two stopping times for any continuous process  $X(\cdot)$ :

$$\begin{aligned} T_z^*(X) &\stackrel{\text{def}}{=} \inf\{t \geq 0 : |X(t)| \geq z\} \\ \Upsilon(X) &\stackrel{\text{def}}{=} \inf\{t > T_z^*(X) : |X(t)| \leq y\}, \end{aligned}$$

with  $\inf \emptyset = \infty$ . Observe that

$$G_{y,z}^c \supset \left\{ T_z^*(p) < z^2; 1 - 2z^2 \leq \Upsilon(p) \leq 1 - z^2; \sup_{\Upsilon(p) \leq t \leq 1} |p(t)| \leq \frac{z}{2} \right\}.$$

Applying the strong Markov property at  $\Upsilon(p)$ , we deduce from the symmetry that

$$\mathbb{P}(G_{y,z}^c) \geq \mathbb{E}\left(\mathbf{1}_{(T_z^*(p) < z^2; 1 - 2z^2 \leq \Upsilon(p) \leq 1 - z^2)} f(y, 1 - \Upsilon(p); z)\right),$$

because  $p(\Upsilon(p)) = \pm y$  on the event  $\{\Upsilon(p) < 1\}$ , and where the function  $f$  is given by

$$f(y, s; z) \stackrel{\text{def}}{=} \mathbb{P}\left(\text{The Brownian bridge from } y \text{ to } 0 \text{ of length } s \text{ always lives in } \left[-\frac{z}{2}, \frac{z}{2}\right]\right).$$

It follows from the scaling property that for all  $z^2 \leq s \leq 2z^2$ ,

$$f(y, s; z) = f\left(\frac{y}{\sqrt{s}}, 1; \frac{z}{\sqrt{s}}\right) \geq \inf_{0 \leq a \leq \frac{1}{4}} f\left(a, 1, \frac{1}{\sqrt{2}}\right) \stackrel{\text{def}}{=} C_5 > 0,$$

because  $a = \frac{y}{\sqrt{s}} \leq \frac{1}{4}$  and  $\frac{z}{\sqrt{s}} \geq \frac{1}{\sqrt{2}}$ . Hence we have shown that

$$\mathbb{P}(G_{y,z}^c) \geq C_5 \mathbb{P}(T_z^*(p) < z^2; 1 - 2z^2 \leq \Upsilon(p) \leq 1 - z^2).$$

Recall the following absolute continuity between the law of a standard Brownian bridge and that of a standard Brownian motion: Denote by  $\mathbb{P}_{0,0}$  the law of  $p(\cdot)$  and by  $\mathbb{P}_0$  that of  $B(\cdot)$ , on the canonical space  $(\mathcal{C}([0, 1] \rightarrow \mathbb{R}), (X(t), 0 \leq t \leq 1), (\mathcal{X}_t)_{0 \leq t \leq 1})$ , we have

$$d\mathbb{P}_{0,0} |_{\mathcal{X}_t} = \frac{1}{\sqrt{1-t}} \exp\left(-\frac{X^2(t)}{2(1-t)}\right) d\mathbb{P}_0 |_{\mathcal{X}_t}, \quad t < 1.$$

Applying the above formula to the stopping time  $\Upsilon(X)$ , we have

$$\begin{aligned}
& \mathbb{P}\left(T_z^*(p) < z^2; 1 - 2z^2 \leq \Upsilon(p) \leq 1 - z^2\right) \\
&= \mathbb{E}_0\left(\mathbf{1}_{(T_z^*(X) < z^2; 1 - 2z^2 \leq \Upsilon(X) \leq 1 - z^2)} \frac{1}{\sqrt{1 - \Upsilon(X)}} \exp\left(-\frac{y^2}{2(1 - \Upsilon(X))}\right)\right) \\
&\geq \frac{e^{-1/32}}{\sqrt{2}} \frac{1}{z} \mathbb{P}_0\left(T_z^*(X) < z^2; 1 - 2z^2 \leq \Upsilon(X) \leq 1 - z^2\right) \\
&\geq \frac{e^{-1/32}}{\sqrt{2}} \frac{1}{z} \mathbb{P}_0\left(T_z^*(X) < \frac{z^2}{2}\right) \mathbb{P}_z\left(1 - 2z^2 \leq T_y(X) \leq 1 - \frac{3z^2}{2}\right),
\end{aligned}$$

where  $\mathbb{P}_z$  means that the Brownian motion  $X(\cdot)$  starts from  $z$  and  $T_y(X)$  denotes the first hitting time at  $y$  of  $X$ . Thanks to the scaling property, the first probability in the above inequality  $\mathbb{P}_0\left(T_z^*(X) < \frac{z^2}{2}\right)$  is bounded below by some numerical constant. Using the well-known distribution of the Brownian hitting time:  $\mathbb{P}_z(T_y(X) \in dt) = \frac{z-y}{\sqrt{2\pi t^3}} e^{-(z-y)^2/(2t)} dt$ , we obtain that

$$\mathbb{P}_z\left(1 - 2z^2 \leq T_y(X) \leq 1 - \frac{3z^2}{2}\right) \geq C_6 z^3.$$

Assembling these estimates, we get

$$\mathbb{P}\left(G_{y,z}^c\right) \geq C_3 z^2,$$

for some universal constant  $C_3 > 0$ , as desired. ■

### 3 Proof of Theorem 1.1

We begin with the proof of the upper bound:

$$\limsup_{t \rightarrow \infty} \frac{M_j^*(t)}{\sqrt{t \log \log t}} \leq \frac{1}{j\sqrt{2}}, \quad \text{a.s.} \quad (3.1)$$

This follows from Proposition 2.5: Fix an arbitrary constant  $a > \frac{1}{j\sqrt{2}}$ . Let  $n \geq 3$  and  $t_n = e^{n/\log n}$ . We have from Proposition 2.5 that

$$\mathbb{P}\left(\sup_{0 \leq t \leq t_{n+1}} M_j^*(t) > a\sqrt{t_n \log \log t_n}\right) \leq C_1 \exp\left(- (2j^2 a^2 + o(1)) \log \log t_n\right),$$

whose sum over  $n$  converges; this in view of a simple application of Borel-Cantelli lemma yields (3.1).

Now, fix an arbitrary constant  $a < \frac{1}{j\sqrt{2}}$ . It suffices to prove that

$$\limsup_{t \rightarrow \infty} \frac{M_j(t)}{\sqrt{t \log \log t}} \geq a, \quad \text{a.s.} \quad (3.2)$$

To this end, let  $t_n = n^n$  and  $\lambda_n = a\sqrt{t_n \log \log t_n}$ , we consider the event

$$E_n \stackrel{\text{def}}{=} \left\{ M_j(t_n) > \lambda_n \right\},$$

which is  $\mathcal{F}_{t_n} \stackrel{\text{def}}{=}} \sigma\{K(s, u), 0 \leq s \leq 1, 0 \leq u \leq t_n\}$ -measurable. If we can show that

$$\sum_n \mathbb{P}\left(E_n \mid \mathcal{F}_{t_{n-1}}\right) = \infty, \quad \text{a.s.} \quad (3.3)$$

then according to Lévy's version of Borel-Cantelli lemma (cf. [13]), we get  $\mathbb{P}\left(E_n, \text{ i.o. } \right) = 1$  hence (3.2).

Consider the process  $\tilde{K}(s, u) \stackrel{\text{def}}{=} K(s, u + t_{n-1}) - K(s, t_{n-1})$  for  $0 \leq s \leq 1$  and  $u \geq 0$ . The independent increment property says that  $\tilde{K}(\cdot, \cdot)$  is independent of  $\mathcal{F}_{t_{n-1}}$  and has the same law as  $K(\cdot, \cdot)$ . Fix a small  $\epsilon > 0$  such that  $2j^2 a^2(1 + 2\epsilon) \leq (1 - 2\epsilon)$ .

Recall the notation  $D_{\tilde{K}(\cdot, t_n - t_{n-1})}$  in Section 2 for the downcrossings by the process  $\tilde{K}(\cdot, t_n - t_{n-1})$ . Observe that

$$\left\{ D_{\tilde{K}(\cdot, t_n - t_{n-1})}(-\epsilon\lambda_n, (1 + \epsilon)\lambda_n; [0, 1]) \geq j \right\} \cap \left\{ \tilde{M}_1^*(t_{n-1}) < \epsilon\lambda_n \right\} \subset E_n,$$

where  $\tilde{M}_1^*(t_{n-1}) \stackrel{\text{def}}{=} \sup_{0 \leq s \leq 1, 0 \leq u \leq t_{n-1}} |\tilde{K}(s, u)|$ . Therefore, we apply Corollary 2.3 and obtain that for all large  $n$ ,

$$\begin{aligned} \mathbb{P}\left(E_n \mid \mathcal{F}_{t_{n-1}}\right) &\geq \mathbf{1}_{(\tilde{M}_1^*(t_{n-1}) < \epsilon\lambda_n)} \mathbb{P}\left(D_{\tilde{K}(\cdot, t_n - t_{n-1})}(-\epsilon\lambda_n, (1 + \epsilon)\lambda_n; [0, 1]) \geq j\right) \\ &= \mathbf{1}_{(\tilde{M}_1^*(t_{n-1}) < \epsilon\lambda_n)} \mathbb{P}\left(D_p\left(-\epsilon \frac{\lambda_n}{\sqrt{t_n - t_{n-1}}}, (1 + \epsilon) \frac{\lambda_n}{\sqrt{t_n - t_{n-1}}}; [0, 1]\right) \geq j\right) \\ &\geq \mathbf{1}_{(\tilde{M}_1^*(t_{n-1}) < \epsilon\lambda_n)} \exp\left(-2j^2 a^2(1 + 2\epsilon) \log \log t_n\right) \\ &\geq \mathbf{1}_{(\tilde{M}_1^*(t_{n-1}) < \epsilon\lambda_n)} n^{-(1-\epsilon)}, \end{aligned} \quad (3.4)$$

where the above equality is due to the self-similarity:  $\tilde{K}(\cdot, v) \stackrel{\text{law}}{=} \sqrt{v} p(\cdot)$  for any fixed  $v > 0$ , and  $p(\cdot)$  is a standard Brownian bridge. Now, applying (1.1), we obtain that almost surely,  $\tilde{M}_1^*(t_{n-1}) < \epsilon\lambda_n$  for all large  $n$ . This together with (3.4) implies (3.3), completing the proof of Theorem 1.1.  $\blacksquare$

## 4 Proof of Theorem 1.2

### 4.1 Upper bound

It suffices to show that

$$\liminf_{t \rightarrow \infty} \frac{\sqrt{\log t}}{\sqrt{t}} M_2^*(t) = 0, \quad \text{a.s.}$$

According to Lévy's version of Borel-Cantelli's lemma (cf. [13]), the above result follows if we can prove that for any constant  $\epsilon > 0$  and for some sequence  $(t_n \uparrow \infty)$ ,

$$\sum_n \mathbb{P}\left(M_2^*(t_n) < \epsilon \sqrt{\frac{t_n}{\log t_n}} \mid \mathcal{F}_{t_{n-1}}\right) = \infty, \quad \text{a.s.} \quad (4.1)$$

where  $\mathcal{F}_t = \sigma\{K(s, u), 0 \leq s \leq 1, 0 \leq u \leq t\}$ . Let us consider  $t_n = n^{3n}$ . By means of (1.1), we have almost surely for all large  $n$ ,

$$\sup_{0 \leq s \leq 1} |K(s, t_{n-1})| \leq \sqrt{t_{n-1} \log n} \stackrel{\text{def}}{=} \lambda_n. \quad (4.2)$$

Consider large  $n$ . Observe that  $\lambda_n \leq \frac{1}{4}\epsilon \sqrt{\frac{t_n}{\log t_n}} \stackrel{\text{def}}{=} \frac{x_n}{4}$ . By the independent increment property,

$$K(\cdot, t_n) = K(\cdot, t_{n-1}) + \tilde{K}(\cdot, t_n - t_{n-1}),$$

with  $\tilde{K}$  a Kiefer process independent of  $\mathcal{F}_{t_{n-1}}$ . The key observation is that

$$\begin{aligned} & \left\{ M_2^*(t_n) \geq x_n \right\} \cap \left\{ \sup_{0 \leq s \leq 1} |K(s, t_{n-1})| \leq \lambda_n \right\} \\ \subset & \left\{ \exists 0 < a_1 < c_1 < b_1 < a_2 < c_2 < b_2 < 1 : |\tilde{K}(a_i, t_n - t_{n-1})| \leq \lambda_n, \right. \\ & \left. |\tilde{K}(b_i, t_n - t_{n-1})| \leq \lambda_n, |\tilde{K}(c_i, t_n - t_{n-1})| \geq x_n - \lambda_n, i = 1, 2 \right\} \stackrel{\text{def}}{=} \tilde{F}_n, \end{aligned}$$

which implies that

$$\tilde{F}_n^c \cap \left\{ \sup_{0 \leq s \leq 1} |K(s, t_{n-1})| \leq \lambda_n \right\} \subset \left\{ M_2^*(t_n) < x_n \right\}.$$

It follows from the independence of  $\tilde{F}_n^c$  and  $\mathcal{F}_{t_{n-1}}$  that

$$\begin{aligned} \mathbb{P}\left(M_2^*(t_n) < x_n \mid \mathcal{F}_{t_{n-1}}\right) & \geq \mathbf{1}_{(\sup_{0 \leq s \leq 1} |K(s, t_{n-1})| \leq \lambda_n)} \mathbb{P}\left(\tilde{F}_n^c\right) \\ & = \mathbf{1}_{(\sup_{0 \leq s \leq 1} |K(s, t_{n-1})| \leq \lambda_n)} \mathbb{P}\left(G_{y,z}^c\right) \\ & \geq C_3 \mathbf{1}_{(\sup_{0 \leq s \leq 1} |K(s, t_{n-1})| \leq \lambda_n)} z^2 \\ & \geq C_3 \frac{\epsilon^2}{4} \mathbf{1}_{(\sup_{0 \leq s \leq 1} |K(s, t_{n-1})| \leq \lambda_n)} \frac{1}{n \log n}, \end{aligned}$$

where the above equality is due to scaling with  $y = \frac{\lambda_n}{\sqrt{t_n - t_{n-1}}}$ ,  $z = \frac{x_n - \lambda_n}{\sqrt{t_n - t_{n-1}}}$ ,  $G_{y,z}^c$  is the complement event of  $G_{y,z}$  which was defined in (2.8), and the second inequality follows from Lemma 2.10. The above lower bound together with (4.2) implies (4.1).  $\blacksquare$

## 4.2 Lower bound

Fix  $j \geq 2$  and  $\chi > \frac{1}{2}$ . We want to show that almost surely for all large  $t$ :

$$M_j(t) > \sqrt{t} (\log t)^{-\chi}.$$

Consider the two-parameter Ornstein-Uhlenbeck process  $(U(v, t), v \in \mathbb{R}, t \geq 0)$  defined by

$$U\left(\log\left(\frac{s}{1-s}\right), t\right) = \frac{K(s, t)}{\sqrt{s(1-s)}}, \quad 0 < s < 1, t \geq 0.$$

Namely,  $\{U(v, t), v \in \mathbb{R}, t \geq 0\}$  is a centered Gaussian process with covariance

$$\mathbb{E}\left(U(v_1, t_1)U(v_2, t_2)\right) = e^{-\frac{|v_1-v_2|}{2}} t_1 \wedge t_2, \quad v_1, v_2 \in \mathbb{R}, t_1, t_2 \geq 0.$$

Let  $0 < \delta < 1$  be small. First, if there exist some (random) times  $\delta \leq u_1 < v_1 < \dots < u_{j-1} < v_{j-1} < u_j \leq 1-\delta$  such that  $U\left(\log\left(\frac{u_i}{1-u_i}\right), t\right) \geq x$  for  $i = 1, \dots, j$  and  $U\left(\log\left(\frac{v_i}{1-v_i}\right), t\right) = 0$  for  $i = 1, \dots, j-1$ , then  $K(u_i, t) \geq x\sqrt{\delta(1-\delta)}$  and  $K(v_i, t) = 0$ . This implies in particular that  $M_j(t) \geq x\sqrt{\delta(1-\delta)}$ .

Recall (2.4). If we denote by  $D_{U(\cdot, t)}(x, y; [-\log(\frac{1-\delta}{\delta}), \log(\frac{1-\delta}{\delta})])$  the number of downcrossings of  $(x, y)$  by  $U(\cdot, t)$  during the time interval  $[-\log(\frac{1-\delta}{\delta}), \log(\frac{1-\delta}{\delta})]$ , then

$$\left\{D_{U(\cdot, t)}\left(0, x; \left[-\log\left(\frac{1-\delta}{\delta}\right), \log\left(\frac{1-\delta}{\delta}\right)\right]\right) \geq j\right\} \subset \left\{M_j(t) \geq x\sqrt{\delta(1-\delta)}\right\}.$$

Fix a small constant  $c = c(\chi) > 0$  whose value will be determined later. Define  $n_k = \exp(\frac{k}{\log k})$  and let  $\delta_k = (\log n_k)^{-2\chi}$ ,  $I_k = [-\log(\frac{1-\delta_k}{\delta_k}), \log(\frac{1-\delta_k}{\delta_k})]$ ,  $x_k = c\sqrt{n_{k+1}}$  for  $k \geq 3$ . Consider the event

$$F_k \stackrel{\text{def}}{=} \left\{\exists t \in [n_k, n_{k+1}) : D_{U(\cdot, t)}(0, x_k; I_k) \leq j-1\right\}.$$

If we can show that

$$\sum_k \mathbb{P}(F_k) < \infty, \tag{4.3}$$

then the Borel-Cantelli lemma implies that almost surely for all large  $k$ ,  $F_k^c$  realizes; hence for all large  $t$ , we have that  $n_k \leq t < n_{k+1}$ , and  $D_{U(\cdot, t)}(0, x_k; I_k) \geq j$ , which implies that  $M_j(t) \geq x_k\sqrt{\delta_k(1-\delta_k)} \geq \frac{c}{2}\sqrt{t}(\log t)^{-\chi}$ . This yields the convergence part of Theorem 1.2, since  $\chi > \frac{1}{2}$  is arbitrary.

To estimate  $\mathbb{P}(F_k)$ , we consider the following stopping time  $\zeta$  with respect to  $\mathcal{F}_t^U = \sigma\{U(x, s), x \in \mathbb{R}, s \leq t\}$ :

$$\zeta = \inf\{t \geq n_k : D_{U(\cdot, t)}(0, x_k; I_k) \leq j-1\}.$$

We want to estimate  $\mathbb{P}(F_k) = \mathbb{P}(\zeta < n_{k+1})$ . Define  $\tilde{U}(v, t) \stackrel{\text{def}}{=} U(v, t + \zeta) - U(v, \zeta)$  for  $v \in \mathbb{R}$  and  $t \geq 0$ . The independent increments property says that  $\tilde{U}$  is independent of  $\mathcal{F}_\zeta^U$  and has the same law as  $U$ . On  $\{\zeta < n_{k+1}\}$ , we have  $D_{U(\cdot, \zeta)}(0, x_k; I_k) \leq j - 1$ ; Fix a small constant  $\epsilon > 0$ . Consider the event

$$G_k \stackrel{\text{def}}{=} \left\{ \sup_{\delta_k \leq s \leq 1 - \delta_k} \left| \tilde{U}\left(\log\left(\frac{1-s}{s}\right), n_{k+1} - \zeta\right) \right| < \epsilon x_k; \zeta < n_{k+1} \right\} \subset F_k.$$

Using the scaling property:  $\tilde{U}(\cdot, t) \stackrel{\text{law}}{=} \sqrt{t}\tilde{U}(\cdot, 1)$  for any fixed  $t > 0$ , we obtain:

$$\begin{aligned} \mathbb{P}(G_k) &= \int_{[n_k, n_{k+1})} \mathbb{P}(\zeta \in dv) \mathbb{P}\left(\sup_{\delta_k \leq s \leq 1 - \delta_k} \left| \tilde{U}\left(\log\left(\frac{1-s}{s}\right), n_{k+1} - v\right) \right| < \epsilon x_k\right) \\ &= \int_{[n_k, n_{k+1})} \mathbb{P}(\zeta \in dv) \mathbb{P}\left(\sup_{\delta_k \leq s \leq 1 - \delta_k} \left| \tilde{U}\left(\log\left(\frac{1-s}{s}\right), 1\right) \right| < \frac{\epsilon x_k}{\sqrt{n_{k+1} - v}}\right) \\ &\geq \mathbb{P}(\zeta < n_{k+1}) \mathbb{P}\left(\sup_{\delta_k \leq s \leq 1 - \delta_k} \left| \tilde{U}\left(\log\left(\frac{1-s}{s}\right), 1\right) \right| < \frac{\epsilon x_k}{\sqrt{n_{k+1} - n_k}}\right). \end{aligned} \quad (4.4)$$

Observe that on  $G_k$ , the number of downcrossings of  $(-\epsilon x_k, (1 + \epsilon)x_k)$  by  $U(\cdot, n_{k+1})$  during  $I_k = [-\log(\frac{1-\delta_k}{\delta_k}), \log(\frac{1-\delta_k}{\delta_k})]$  can not be larger or equal to  $j$ ; otherwise, we would get  $D_{U(\cdot, \zeta)}(0, x_k; I_k) \geq j$ . In view of this remark, we get

$$\begin{aligned} \mathbb{P}(F_k) &\leq \frac{\mathbb{P}\left(D_{U(\cdot, n_{k+1})}(-\epsilon x_k, (1 + \epsilon)x_k; I_k) \leq j - 1\right)}{\mathbb{P}\left(\sup_{\delta_k \leq s \leq 1 - \delta_k} \left| \tilde{U}\left(\log\left(\frac{1-s}{s}\right), 1\right) \right| < \frac{\epsilon x_k}{\sqrt{n_{k+1} - n_k}}\right)} \\ &= \frac{\mathbb{P}\left(D_{U(\cdot, 1)}(-\epsilon c, (1 + \epsilon)c; I_k) \leq j - 1\right)}{\mathbb{P}\left(\sup_{\delta_k \leq s \leq 1 - \delta_k} \left| U\left(\log\left(\frac{1-s}{s}\right), 1\right) \right| < \frac{\epsilon x_k}{\sqrt{n_{k+1} - n_k}}\right)}, \end{aligned} \quad (4.5)$$

by using the scaling property. We shall bound below the denominator and bound above the numerator in (4.5): the denominator equals

$$\begin{aligned} &\mathbb{P}\left(\sup_{-\log((1-\delta_k)/\delta_k) \leq v \leq \log((1-\delta_k)/\delta_k)} |U(v, 1)| < \frac{\epsilon x_k}{\sqrt{n_{k+1} - n_k}}\right) \\ &= \mathbb{P}\left(\sup_{0 \leq v \leq 2\log((1-\delta_k)/\delta_k)} |U(v, 1)| < \frac{\epsilon x_k}{\sqrt{n_{k+1} - n_k}}\right) \\ &\geq \delta_k^{o(1)}, \quad k \rightarrow \infty, \end{aligned} \quad (4.6)$$

where the above equality follows from the stationarity and the above inequality follows from (2.7) in Fact 2.8 with  $z = \frac{\epsilon x_k}{\sqrt{n_{k+1} - n_k}} \rightarrow \infty$ . On the other hand, we have from Lemma 2.9 that

$$\mathbb{P}\left(D_{U(\cdot, 1)}(-\epsilon c, (1 + \epsilon)c; I_k) \leq j - 1\right) \leq \delta_k^{(2\mu(-\epsilon c, (1 + \epsilon)c) + o(1))}$$



$$= \left( \frac{k}{\log k} \right)^{-(4\chi \mu(-\epsilon c, (1+\epsilon)c) + o(1))}. \quad (4.7)$$

Recall that  $\chi > \frac{1}{2}$ . Since  $\mu(-\epsilon c, (1+\epsilon)c) \rightarrow \frac{1}{2}$  as  $c \rightarrow 0$ , we can choose a sufficiently small constant  $c = c(\chi) > 0$  such that  $4\chi \mu(-\epsilon c, (1+\epsilon)c) > 1$ . Putting (4.6) and (4.7) into (4.5), we obtain some constant  $a > 1$  such that for all large  $k$ ,

$$\mathbb{P}(F_k) \leq k^{-a}$$

proving (4.3), as desired. ■

## Acknowledgements

The authors are grateful to an anonymous referee for her/his suggestions. Cooperation between the authors was supported by the joint French–Hungarian Intergovernmental Grant ”Balaton” (grant no. F-39/00).

## References

- [1] Csáki, E.: Strong limit theorems for empirical processes. *Recent Adv. in Stat. and Prob.* (eds: J. Pérez Vilaplana and M.L. Puri) pp. 247–254, VSP 1994.
- [2] Csáki, E. and Hu, Y.: Asymptotic properties of ranked heights in Brownian excursions. *J. Theoret. Probab.* 14 (2001) no.1, 77–96.
- [3] Csáki, E. and Shi, Z.: Some liminf results for two-parameter processes. *Stoch. Proc. Appl.* 78 (1998) no. 1, 27–46.
- [4] Csörgő, M. and Révész, P.: *Strong Approximations in Probability and Statistics*. Akadémiai Kiadó, Budapest and Academic Press, New York, 1981.
- [5] Freedman, D.: *Brownian Motion and Diffusion*. Holden-Day, San Francisco, 1971.
- [6] Greenwood, P. and Perkins, E.: A conditioned limit theorem for random walk and Brownian local time on square root boundaries. *Ann. Probab.* 11 (1983), no. 2, 227–261.
- [7] Kiefer, J.: Skorohod embedding of multivariate rv’s, and the sample df. *Z. Wahrsch. Verw. Gebiete* 24 (1972) 1–35.
- [8] Mogul’skii, A.A.: Remarks on large deviations of the  $\omega^2$  statistic. (Russian) *Teor. Veroyatnost. i Primenen.* 22 (1977), no. 1, 170–175.

- [9] Perkins, E.: On the Hausdorff dimension of the Brownian slow points. *Z. Wahrsch. Verw. Gebiete* 64 (1983), no. 3, 369–399.
- [10] Pitman, J. and Yor, M.: On the lengths of excursions of some Markov processes. *Séminaire de Probabilités, XXXI* pp. 272–286, Lecture Notes in Math., 1655 Springer, Berlin, 1997.
- [11] Pitman, J. and Yor, M.: Ranked functionals of Brownian excursions. *C. R. Acad. Sci. Paris Sér. I* 326 (1998), 93–97.
- [12] Pitman, J. and Yor, M.: On the distribution of ranked heights of excursions of a Brownian bridge. *Ann. Probab.* 29 (2001) no. 1, 361–384.
- [13] Shiryaev, A.N.: *Probability* (2nd edition), Springer, New York, 1996.
- [14] Uchiyama, K.: Brownian first exit from and sojourn over one-sided moving boundary and application. *Z. Wahrsch. Verw. Gebiete* 54 (1980), no. 1, 75–116.