

On the regularity of the blow-up set for semilinear heat equations

Hatem Zaag

Courant Institute and CNRS ENS

Abstract : We consider $u(x, t)$ a blow-up solution of $u_t = \Delta u + |u|^{p-1}u$ where $u : \mathbb{R}^N \times [0, T) \rightarrow \mathbb{R}$, $p > 1$, $(N - 2)p < N + 2$ and either $u(0) \geq 0$ or $(3N - 4)p < 3N + 8$. The blow-up set $S \subset \mathbb{R}^N$ of u is the set of all blow-up points. Under a non degeneracy condition, we show that if S is continuous, then it is a C^1 manifold.

Résumé : On considère $u(x, t)$ une solution singulière de $u_t = \Delta u + |u|^{p-1}u$ où $u : \mathbb{R}^N \times [0, T) \rightarrow \mathbb{R}$, $p > 1$, $(N - 2)p < N + 2$ et soit $u(0) \geq 0$, soit $(3N - 4)p < 3N + 8$. On définit l'ensemble singulier $S \subset \mathbb{R}^N$ de u comme étant l'ensemble de tous les points d'explosion. Sous une certaine condition de non dégénérescence, on montre que si S est continu, alors c'est une variété de classe C^1 .

AMS Classification : 35K55, 35B40

1 Introduction

We are concerned in this paper with blow-up phenomena arising in the following semilinear problem :

$$\begin{aligned} u_t &= \Delta u + |u|^{p-1}u \\ u(\cdot, 0) &= u_0 \in L^\infty(\mathbb{R}^N), \end{aligned} \tag{1}$$

where $u(t) : x \in \mathbb{R}^N \rightarrow u(x, t) \in \mathbb{R}$ and Δ stands for the Laplacian in \mathbb{R}^N . We assume in addition the exponent $p > 1$ subcritical : if $N \geq 3$ then $1 < p < (N + 2)/(N - 2)$. Moreover, we assume that

$$u_0 \geq 0 \text{ or } (3N - 4)p < 3N + 8. \tag{2}$$

This problem has attracted a lot of attention because it captures features common to a whole range of blow-up problems arising in various physical situations, particularly the role of scaling and self-similarity. Without

pretending to be exhaustive, we would like nonetheless to mention some related equations : the motion by mean curvature (Soner and Souganidis [23]), vortex dynamics in superconductors (Chapman, Hunton and Ockendon [6], Merle and Zaag [18]), surface diffusion (Bernoff, Bertozzi and Witelski [2]) and chemotaxis (Brenner *et al.* [4], Betterton and Brenner [3]). However, equation (1) is simple enough to be tractable in rigorous mathematical terms, unlike other physical equations.

A solution $u(t)$ to (1) blows-up in finite time if its maximal existence time T is finite. In this case,

$$\lim_{t \rightarrow T} \|u(t)\|_{H^1(\mathbb{R}^N)} = \lim_{t \rightarrow T} \|u(t)\|_{L^\infty(\mathbb{R}^N)} = +\infty.$$

Let us consider such a solution. T is called the blow-up time of u . A point $a \in \mathbb{R}^N$ is called a blow-up point if

$$|u(x, t)| \rightarrow +\infty \text{ as } (x, t) \rightarrow (a, T)$$

(this definition is equivalent to the usual local unboundedness definition, thanks to Corollary 2 in [21]). S denotes the blow-up set, that is the set of all blow-up points. From [21], we know that there exists a *blow-up profile* $u^* \in C_{\text{loc}}^2(\mathbb{R}^N \setminus S)$ such that

$$u(x, t) \rightarrow u^*(x) \text{ in } C_{\text{loc}}^2(\mathbb{R}^N \setminus S) \text{ as } t \rightarrow T. \quad (3)$$

The blow-up problem has been addressed in different ways in the literature. A major direction was developed by authors looking for sufficient blow-up conditions on initial data or on the nonlinear term (see Fujita [12], Ball [1], Levine [16] and the review paper by Deng and Levine [7]). The second main direction is about the description of the asymptotic blow-up behavior, *locally* near a given blow-up point \hat{a} (see Giga and Kohn [13], Bricmont and Kupiainen [5], Herrero and Velázquez [14], [24], Merle and Zaag [21]). Given $a \in \mathbb{R}^N$ a blow-up point of y , we know that up to some scalings, u approaches a particular explicit function near the singularity (a, T) (see [24]). Up to replacing u by $-u$, one of the following two cases occurs :

Case 1 : For all $K_0 > 0$,

$$\sup_{|z| \leq K_0} \left| (T-t)^{\frac{1}{p-1}} u \left(a + \tilde{Q}_a z \sqrt{(T-t)|\log(T-t)|}, t \right) - f_{l_a}(z) \right| \rightarrow 0 \quad (4)$$

as $t \rightarrow T$, where \tilde{Q}_a is an orthonormal $N \times N$ matrix, $l_a = 1, \dots, N$ and

$$f_{l_a}(z) = \left(p - 1 + \frac{(p-1)^2}{4p} \sum_{i=1}^{l_a} z_i^2 \right)^{-\frac{1}{p-1}}. \quad (5)$$

Case 2 : For all $K_0 > 0$,

$$\sup_{|z| \leq K_0} \left| (T-t)^{\frac{1}{p-1}} u \left(a + z(T-t)^{\frac{1}{2k}}, t \right) - \left(p-1 + \sum_{|\alpha|=2k} C_\alpha z^\alpha \right)^{-\frac{1}{p-1}} \right| \quad (6)$$

goes to 0 as $t \rightarrow T$, where $k = 2, 3, 4, \dots$, $x^\alpha = x_1^{\alpha_1} \dots x_N^{\alpha_N}$ and $|\alpha| = \alpha_1 + \dots + \alpha_N$ if $\alpha = (\alpha_1, \dots, \alpha_N)$ and $\sum_{|\alpha|=2k} C_\alpha x^\alpha \geq 0$ for all $x \neq 0$.

Remark : Even though the proof of [24] is given in the positive case, it extends to unsigned solutions under (2).

The description of the blow-up set S is a major issue. Examples where S is a set of isolated points or a sphere are known to exist (see [17] and [19] for isolated points and [13] for the sphere). If these solutions are artificially considered as defined on $\mathbb{R}^{N'} \times [0, T)$ where $N' > N$, we obtain examples where S consists in a collection of $(N' - N)$ -dimensional subspaces or spheres. No other geometric configurations are known to occur. In [26], Velázquez proves the following result :

The $(N - 1)$ -dimensional Hausdorff measure of S is bounded on compact sets.

No other regularity result is known.

Our first goal in this paper is to improve this result and obtain partial regularity results on S under some reasonable conditions. Let us consider $\hat{a} \in S$. According to [24] (remark after Theorem 2), if (4) occurs with $l = N$ or (6) occurs with $\sum C_\alpha x^\alpha > 0$ for all $x \neq 0$ (no degenerate directions in the function), then the blow-up point is isolated. The question remains open in the other cases. Even if one assumes that \hat{a} is not isolated, it is unclear whether there is a continuum of blow-up points near \hat{a} or not. This question seems to be very difficult. Whatever the answer is, we don't know how S looks like near \hat{a} , and how the profile u^* is near S (no relevant information on u^* near a non isolated blow-up point was known before). To make our presentation clearer, we restrict to the case $N = 2$ and consider \hat{a} a non isolated point of S such that \hat{a} belongs to a continuous line of blow-up points without being an endpoint. More precisely, we assume that $\hat{a} = a(0) \in \text{Im } a \subset S$ where $a \in C((-1, 1), \mathbb{R}^2)$ and for some α_0 ,

$$\forall \epsilon > 0, a(-\epsilon, \epsilon) \text{ intersects the complimentary of any connected closed cone with vertex at } \hat{a} \text{ and angle } \alpha \in (0, \alpha_0] \quad (7)$$

(this is in a way to insure that \hat{a} is not an endpoint).

Assuming that u behaves according to (4) near the singularity (\hat{a}, T) , we have the following result :

Theorem 1 (Regularity of the blow-up set at a point with the behavior (4) assuming S contains a continuum) *Assume $N = 2$ and consider u a solution of (1) that blows-up at time T on a set S . Consider $\hat{a} = a(0) \in \text{Im } a \subset S$ where $a \in C((-1, 1), \mathbb{R}^2)$ and \hat{a} is not an endpoint (in the sense (7)). If u behaves near (\hat{a}, T) as stated in (4), then there are $\delta > 0$, $\delta_1 > 0$ and $\varphi \in C^1([-\delta_1, \delta_1], \mathbb{R})$ such that*

$$S \cap B(\hat{a}, 2\delta) = \text{graph } \varphi \cap B(\hat{a}, 2\delta) = \text{Im } a \cap B(\hat{a}, 2\delta). \quad (8)$$

In particular, S is a C^1 manifold near the point \hat{a} .

We actually have the following refined C^1 estimate for φ .

Proposition 2 (Refined C^1 estimate for S) *There exists $C_0 > 0$ and h_0 such that for all $|\xi| < \delta_1$ and $|h| < h_0$ such that $|\xi + h| < \delta_1$, we have :*

$$|\varphi(\xi + h) - \varphi(\xi) - h\varphi'(\xi)| \leq C_0|h| \sqrt{\frac{\log |\log |h||}{|\log |h||}}.$$

Remark : Using the techniques of Fermanian-Zaag [9], we show in [27] that φ is actually $C^{1,\alpha}$ for any $\alpha \in (0, \frac{1}{2})$.

Remark : From [24], we know that the limit function at (\hat{a}, T) stated in (4) has a degenerate direction, and that we can not have two curves of blow-up points intersecting transversally at \hat{a} . With our contribution, we eliminate the possibility of two curves meeting tangentially at \hat{a} . In particular, there is no cusp at \hat{a} , and there is no sequence of isolated blow-up points converging to $\hat{a} \in S$.

Remark : The case we are considering does exist indeed. The techniques of [19] hold for the one dimensional equation

$$\partial_t v = \partial_{rr}^2 v + \frac{N-1}{r} \partial_r v + |v|^{p-1} v$$

which is the radial case of (1). Thus, for all $r_0 > 0$, there is a radial solution $u(x, t) = v(|x|, t)$ of (1) such that for all $K_0 > 0$,

$$\sup_{|z| \leq K_0} \left| (T-t)^{\frac{1}{p-1}} v \left(r_0 + z \sqrt{(T-t)|\log(T-t)|} \right) - f(z) \right| \rightarrow 0 \text{ as } t \rightarrow T$$

$$\text{where for all } z \in \mathbb{R}, f(z) = \left(p - 1 + \frac{(p-1)^2}{4p} z^2 \right)^{-\frac{1}{p-1}}. \quad (9)$$

The blow-up set of u is the sphere $r_0 \mathbb{S}^{N-1}$, and near each blow-up point, (4) holds with the degenerate profile f_1 .

The description of the *blow-up profile* u^* defined in (3) near the singularity (\hat{a}, T) is our second concern in this paper. We claim the following :

Theorem 3 (Blow-up behavior and profile near a blow-up point where u behaves as in (4) assuming S contains a continuum) *With the notations of Theorem 1, there exists $t_0 < T$ such that for all $K_0 > 0$, $t \in [t_0, T)$ and $x \in B(\hat{a}, \delta)$ s.t. $d(x, S) \leq K_0 \sqrt{(T-t)|\log(T-t)|}$, we have*

$$\left| (T-t)^{\frac{1}{p-1}} u(x, t) - f \left(\frac{d(x, S)}{\sqrt{(T-t)|\log(T-t)|}} \right) \right| \leq C'_0(K_0) \frac{\log |\log(T-t)|}{|\log(T-t)|} \quad (10)$$

where f is defined in (9). Moreover, $\forall x \in \mathbb{R}^N \setminus S$, $u(x, t) \rightarrow u^*(x)$ as $t \rightarrow T$ with

$$u^*(x) \sim U(d(x, S)) \text{ as } d(x, S) \rightarrow 0 \text{ and } x \in B(\hat{a}, \delta) \quad (11)$$

where $U(z) = \left(\frac{8p}{(p-1)^2} \frac{|\log z|}{z^2} \right)^{\frac{1}{p-1}}$ for $z > 0$.

Remark : This is the first time where the blow-up profile u^* is derived near a non-isolated point. Indeed, in the earlier work of Velázquez, the behavior along the “tangential” direction of S was not derived. (10) shows that in a tubular neighborhood of S , the main term in the blow-up asymptotics is the 1D blow-up profile f , function of only the normal coordinate $\pm d(x, S)$.

Remark : When $p > 3$, we show in [27] that up to a non singular function, u is a superposition of 1D blow-up solutions of (1), organized along the normal directions to the blow-up set.

Theorems 1 and 3 hold in higher dimensions $N \geq 3$. However, the hypotheses should be stated more carefully. We claim the following

Theorem 4 (Regularity of the blow-up set near a point with the behavior (4) assuming S contains a $N-l$ dimensional continuum) *Take $N \geq 2$ and $l \in \{1, \dots, N-1\}$. Consider u a solution of (1) that blows-up at time T on a set S and take $\hat{a} \in S$ where u behaves locally as stated*

in (4). Consider $a \in C((-1, 1)^{N-l}, \mathbb{R}^N)$ such that $\hat{a} = a(0) \in \text{Im } a \subset S$ and $\text{Im } a$ is at least $(N - l)$ dimensional (in the sense (82)). If \hat{a} is not an endpoint (in the sense (83) given below), then there are $\delta > 0$, $\delta_1 > 0$ and $\varphi \in C^1([-\delta_1, \delta_1]^{N-l}, \mathbb{R}^l)$ such that (8) holds and S is a C^1 manifold near \hat{a} . Proposition 2 and Theorem 3 hold as well.

Remark : If $l = N - 1$, then the fact that \hat{a} is not isolated implies that $\text{Im } a$ is at least 1 dimensional near \hat{a} .

Remark : Theorem 4 can be stated without the hypotheses (82) and (83) if we strengthen the assumption on $\text{Im } a$. Indeed, If we already know that $\text{Im } a$ is a $(N - l)$ -dimensional differentiable manifold, then we learn from Theorem 4 that $S \setminus \text{Im } a$ is empty, locally near \hat{a} , and we get the blow-up profile near \hat{a} as stated in Theorem 3.

Up to some complications in the notation, the proof of Theorem 4 remains the same as in the case $N = 2$. We will show in section 6 how to adapt the proof of the case $N = 2$ to the general case.

The paper is organized as follows. In section 2, we recall from previous work the self-similar variables technique and a Liouville Theorem for equation (1). In section 3, we show the stability of the behavior (4) (with $l = 1 < 2 = N$) with respect to the blow-up point in $\text{Im } a$. The regularity of the blow-up set is presented in section 4 where we prove Theorem 1 and Proposition 2. Section 5 is devoted to the blow-up profile of u (Theorem 3). In section 6, we sketch the proof of Theorem 4.

The author wants to thank T. Colding, R. V. Kohn, F. H. Lin and F. Merle for interesting conversations and remarks about the paper. He wants to acknowledge partial support he received from the NSF grant DMS-9631832.

2 Asymptotic behavior in self-similar variables and global estimates for blow-up solutions of (1)

In this section, we introduce the general framework for the study of u near a singularity (a, T) and recall from [21] a uniform (in space and time) comparison property of u with the solution of the associated ODE $u' = u^p$.

2.1 Self-similar variables

Given a a blow-up point of u , we study the behavior of u near the singularity (a, T) through the introduction of the function w_a defined by

$$w_a(y, s) = (T - t)^{\frac{1}{p-1}} u(x, t), \quad y = \frac{x - a}{\sqrt{T - t}}, \quad s = -\log(T - t). \quad (12)$$

From (1), we see that w_a satisfies for all $(y, s) \in \mathbb{R}^N \times [-\log T, \infty)$ the following equation

$$\frac{\partial w}{\partial s} = \Delta w - \frac{1}{2} y \cdot \nabla w - \frac{w}{p-1} + |w|^{p-1} w. \quad (13)$$

We know from [13] that

$$\|w_a\|_{L^\infty(\mathbb{R}^N \times [-\log T, \infty))} \leq M < \infty \quad (14)$$

((12) shows that M is independent of a) and that

$$w_a(y, s) \rightarrow \pm \kappa \equiv \pm(p-1)^{-\frac{1}{p-1}} \text{ as } s \rightarrow \infty \quad (15)$$

in L_ρ^2 where $\rho(y) = e^{-\frac{|y|^2}{4}} / (4\pi)^{N/2}$ and uniformly on compact sets. Assuming that $w_a \rightarrow \kappa$, we define

$$v_a = w_a - \kappa. \quad (16)$$

We know from (15) and (13) that $\|v_a\|_{L_\rho^2} \rightarrow 0$ as $s \rightarrow \infty$ and for all $(y, s) \in \mathbb{R}^N \times [-\log T, \infty)$,

$$\frac{\partial v_a}{\partial s} = \mathcal{L}v_a + f(v_a) \equiv \mathcal{L}v_a + \frac{p}{2\kappa} v_a^2 + g(v_a) \quad (17)$$

where $\mathcal{L} = \Delta - \frac{1}{2} y \cdot \nabla + 1$, $|f(v_a)| \leq C(M)|v_a|^2$ and $|g(v_a)| \leq C(M)|v_a|^3$. Operator \mathcal{L} is self-adjoint on L_ρ^2 , its spectrum is $\text{spec } \mathcal{L} = \{1 - \frac{m}{2} \mid m \in \mathbb{N}\}$. Its eigenfunctions are derived from the Hermite polynomials. If $N = 1$, all the eigenvalues of \mathcal{L} are simple. To $1 - \frac{m}{2}$ corresponds the eigenfunction

$$h_m(y) = \sum_{n=0}^{\lfloor \frac{m}{2} \rfloor} \frac{m!}{n!(m-2n)!} (-1)^n y^{m-2n}.$$

If $N \geq 2$, then the eigenfunctions corresponding to $1 - \frac{m}{2}$ are

$$H_\alpha(y) = h_{\alpha_1}(y_1) \dots h_{\alpha_N}(y_N), \quad \text{with } \alpha = (\alpha_1, \dots, \alpha_N) \text{ and } |\alpha| = m.$$

In particular,

1 is an eigenvalue of multiplicity 1 and its eigenfunction is $H_0(y) = 1$,
 $\frac{1}{2}$ is of multiplicity N and its eigenspace is generated by the orthogonal basis $\{y_i \mid i = 1, \dots, N\}$,
0 is of multiplicity $\frac{N(N+1)}{2}$ and its eigenspace is generated by the orthogonal basis

$$\{y_i y_j \mid i < j\} \cup \{y_i^2 - 2 \mid i = 1, \dots, N\}. \quad (18)$$

Since the eigenfunctions of \mathcal{L} make a total orthonormal family of L_ρ^2 , we expand v_a as follows

$$v_a(y, s) = \sum_{m=0}^2 v_{a,m}(y, s) + v_{a,-}(y, s) \equiv v_{a,2}(y, s) + v_{a,-}(y, s) + v_{a,+}(y, s), \quad (19)$$

where $v_{a,m}(y, s)$ is the orthogonal projection of v_a on the eigenspace of $\lambda = 1 - \frac{m}{2}$, $v_{a,-}(y, s) = P_-(v_a)(y, s)$ and P_- is the projector on the negative subspace of \mathcal{L} . Let us define a $N \times N$ symmetric matrix $A_a(s)$ by

$$A_a(s) = \int_{\mathbb{R}^N} v_a(y, s) M(y) \rho(y) dy \text{ where } M_{i,j}(y) = \frac{1}{4} y_i y_j - \frac{1}{2} \delta_{ij}. \quad (20)$$

Then, from (19), (18) and the orthogonality between eigenfunctions of \mathcal{L} , we have

$$v_{a,2}(y, s) = \frac{1}{2} y^T A_a(s) y - \text{tr } A_a(s). \quad (21)$$

From Filippas and Liu [11] and Velázquez [25], we know that

$$\text{either } v_a \sim v_{a,2} \text{ or } v_a \sim v_{a,-} \text{ in } L_\rho^2 \text{ as } s \rightarrow \infty. \quad (22)$$

In the former case, we know that for some $l_a \in \{1, \dots, N\}$, $\delta_a > 0$ and a $N \times N$ orthogonal matrix \tilde{Q}_a , we have

$$v_a(\tilde{Q}_a y, s) = \frac{\kappa}{2ps} \left(l_a - \frac{1}{2} \sum_{i=1}^{l_a} y_i^2 \right) + O\left(\frac{1}{s^{1+\delta_a}} \right) \text{ as } s \rightarrow \infty \quad (23)$$

in L_ρ^2 and u behaves near (a, T) as stated in (4).

If $l_a = N$, then a is an isolated blow-up point. We proved in [8] with Fermanian and Merle the stability of such a behavior with respect to perturbations in initial data.

In this paper, we consider the case $l_a < N$ and assume that a is not isolated. Although the techniques of [8] imply that this profile is unstable with respect to perturbations in *initial data*, we will show in section 3 its stability with respect to *the blow-up point* (for a fixed solution), in the smaller class of *non* isolated blow-up points.

2.2 A Liouville Theorem and ODE comparison for u

The following rigidity theorem (from [21]) is crucial in the blow-up study of (1). It is a central argument in the proof of our Theorem.

Proposition 2.1 (A Liouville Theorem for equation (1)) *Let u be a solution of (1) defined for all $(x, t) \in \mathbb{R}^N \times (-\infty, T)$ such that for some $C > 0$,*

$$|u(x, t)| \leq \frac{C}{(T - t)^{\frac{1}{p-1}}}.$$

Then, either $u \equiv 0$ or there exist $T_1 \in [T, +\infty)$ and $\omega_0 \in \{-1, +1\}$ such that

$$u(x, t) = \omega_0 \kappa (T_1 - t)^{-\frac{1}{p-1}}.$$

This allows Merle and Zaag [21] to prove for $u_0 \in C^2$ the following localization property which reduces the study of the evolution of $u(b, t)$ for a fixed b to the study of an ODE :

Proposition 2.2 (Uniform ODE comparison of blow-up solutions of (1)) *For all $\epsilon > 0$, there exists $C = C(\epsilon, \|u_0\|_{C^2}, T)$ such that $\forall (x, t) \in \mathbb{R}^N \times [0, T)$,*

$$|\partial_t u - |u|^{p-1} u| \leq \epsilon |u|^p + C.$$

As a consequence, we have the following criterion for regular points (by definition, non blow-up points) :

Proposition 2.3 (Blow-up exclusion criterion) *For all $\epsilon_0 > 0$, there exists $t_0(\epsilon) < T$ such that if $|u(a, t)| \leq (1 - \epsilon_0) \kappa (T - t)^{-1/(p-1)} \equiv (1 - \epsilon_0) v_T(t)$ for some $a \in \mathbb{R}^N$ and $t \in [t_0(\epsilon_0), T)$, then a is not a blow-up point.*

Remark : v_T is the solution of $v'_T = v_T^p$, $v_T(T) = \infty$.

Proof : See Corollary 1 in [20] where the criterion is derived from the ODE comparison (Note that in [20] the criterion holds only for positive data, but since we show in [21] the ODE comparison for unsigned data, the criterion holds in this general case). ■

3 Stability of the blow-up behavior (4) with respect to non isolated blow-up points

From now on, we take $1 = l < N = 2$. We consider \hat{a} a blow-up point of u such that $\hat{a} = a(0)$ where $a \in C((-1, 1), \mathbb{R}^2)$ and \hat{a} is not an endpoint of $\text{Im } a \subset S$ in the sense (7). We assume that u has the behavior (4) near (\hat{a}, T) . From rotation and translation invariance, we assume that $\hat{a} = 0$ and $\tilde{Q}_{\hat{a}} = \text{Id}$. Thus, (4) implies that

$$\sup_{|z| \leq K_0} \left| (T-t)^{\frac{1}{p-1}} u(z\sqrt{(T-t)|\log(T-t)|}, t) - f(z_1) \right| \rightarrow 0 \text{ as } t \rightarrow T \quad (24)$$

where f is defined in (9). Since u has the behavior (24) near $(0, T)$, we know from the previous section (see (22) and (23)) that

$$v_0 \sim v_{0,2} \text{ and } w_0(y, s) - \kappa = v_0(y, s) \sim \frac{\kappa}{2ps} \left(1 - \frac{y_1^2}{2}\right) \text{ as } s \rightarrow \infty \quad (25)$$

in L^2_ρ , where v_0 and w_0 are defined in (12) and (16). In the following, we will write a instead of $a(\sigma)$ and v_a instead of $v_{a(\sigma)}$. A central argument in our proof is the following :

Proposition 3.1 (Stability of the L^2_ρ asymptotic behavior with respect to blow-up points in $\text{Im } a$) *There exist $\sigma_0 > 0$, $C_0 > 0$ and $s_0 \in \mathbb{R}$ such that for all $b \in a(-\sigma_0, \sigma_0)$, there exists Q_b a 2×2 orthogonal matrix such that :*

i) for all $|\sigma| < \sigma_0$ and $s \geq s_0$,

$$\left\| w_a(Q_a y, s) - \left\{ \kappa + \frac{\kappa}{2ps} \left(1 - \frac{y_1^2}{2}\right) \right\} \right\|_{L^2_\rho} \leq C_0 \frac{\log s}{s^2}.$$

ii) $Q_0 = \text{Id}$ and $b \in a(-\sigma_0, \sigma_0) \rightarrow Q_b$ is continuous.

iii) For all $K_0 > 0$, there is $C'_0(K_0) > 0$ such that for all $s \geq s_0$,

$$\sup_{|\sigma| < \sigma_0, |y| \leq K_0 \sqrt{s}} \left| w_a(Q_a y, s) - f\left(\frac{y_1}{\sqrt{s}}\right) \right| \leq C'_0(K_0) \frac{\log s}{s}$$

where f is defined in (9).

Remark : This argument is similar to the result of [8], where we proved the stability of the blow-up behavior (4) with $l = N$ (the isolated blow-up point case), with respect to initial data. Therefore, we will refer to [8] for

the similar steps.

The proof of this Proposition follows from 4 steps.

- In Step 1, we show that the control of v_a near the same asymptotic L^2_ρ behavior as v_0 reduces to the control of its neutral mode $v_{a,2}$, that is the matrix A_a defined in (20) and (21) (this is a finite dimensional problem).

- In Step 2, we show that the eigenvalues of $A_a(s)$ have uniformly the same behavior as those of $A_0(s)$ as $s \rightarrow +\infty$.

- In Step 3, we solve the finite dimensional problem by finding the long time behavior of A_a .

- In Step 4, we give the solution of the infinite dimensional problem (that is the asymptotics of w_a as $s \rightarrow \infty$), which concludes the proof of Proposition 3.1.

Step 1 : Uniform reduction to a finite dimensional problem

In this step, the only relevant information on v_0 we use is that $v_0 \sim v_{0,2}$. We aim at showing that this extends to any $a(\sigma)$ near 0. In particular, the fact that the asymptotic behavior in (25) has a degenerate direction is not relevant here. Thus, this step is not new. It is exactly the same as the analogous one in the proof of the stability of the profile (4) with $l = N$ presented in [8]. Therefore, we just summarize the arguments of the proof in Appendix A. Let us just remark that the Liouville Theorem (Proposition 2.1) is the central argument in getting the uniformity. We claim the following :

Proposition 3.2 (Reduction to a finite dimensional problem) *There exists $\sigma_1 > 0$ such that for all $\epsilon > 0$, there is $s_1(\epsilon)$ such that for all $|\sigma| < \sigma_1$,*

$$\forall s \geq s_1(\epsilon), \begin{cases} \|v_a(s)\|_{L^2_\rho} \leq \epsilon, & \|v_a - v_{a,2}(s)\|_{L^2_\rho} \leq \epsilon \|v_{a,2}(s)\|_{L^2_\rho}, \\ |A'_a(s) - \frac{1}{\beta} A_a(s)| \leq \epsilon |A_a(s)|^2 \end{cases} \quad (26)$$

where $\beta = \frac{\kappa}{2p}$ and $v_{a,2}$ and A_a are defined in (19), (21) and (20).

Proof : See Appendix A. ■

Step 2 : A spectral study of the finite dimensional problem

In Steps 2 and 3, we solve the finite dimensional problem given by Step 1. Since A_a is a symmetric matrix, we can define its eigenvalues as follows :

Lemma 3.3 (Existence of regular eigenvalues for A_a) *There exist 2 real C^1 functions $l_{a,i}(s)$, $i = 1, 2$ eigenvalues of $A_a(s)$. Moreover, the set $\{l_{a,1}(s), l_{a,2}(s)\}$ is continuous in terms of $(a, s) \in S \times [-\log T, \infty)$.*

Proof : From the regularity of w_a , it is clear that for each $a \in \mathbb{R}^N$, the symmetric matrix $A_a(s)$ is a C^1 function of s . Therefore, according to Kato [15], we can define 2 C^1 functions of s , $l_{a,1}(s)$ and $l_{a,2}(s)$, eigenvalues of $A_a(s)$ (see Lemma 3.2 in [11] for a statement). Since $A_a(s)$ is a continuous function of (a, s) and the eigenvalues of a matrix vary continuously with respect to the coefficients, $\{l_{a,1}(s), l_{a,2}(s)\}$ is continuous in terms of (a, s) . ■

Proposition 3.2 and section 2.1 have the following Corollary :

Corollary 3.4

i) **(Non uniform behavior of v_a)** For all $|\sigma| < \sigma_1$, (23) holds with $l_a = 1$. In particular,

$$A_a(s) = -\frac{\beta}{s} \tilde{Q}_a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \tilde{Q}_a^T + O\left(s^{-1-\delta_a}\right) \text{ as } s \rightarrow \infty,$$

and one eigenvalue is equal to $-\frac{\beta}{s} + O\left(s^{-1-\delta_a}\right)$ while the other is equal to $O\left(s^{-1-\delta_a}\right)$ as $s \rightarrow \infty$.

ii) **(Equations on eigenvalues)** For all $\epsilon > 0$, there is $s_1(\epsilon)$ such that for all $i \in \{1, 2\}$, $|\sigma| < \sigma_1$ and $s \geq s_1(\epsilon)$,

$$\left|l'_{a,i}(s) - \frac{1}{\beta} l_{a,i}(s)^2\right| \leq \epsilon(l_{a,1}^2 + l_{a,2}^2).$$

Proof : i) From Proposition 3.2, we have $v_a \sim v_{a,2}$ as $s \rightarrow \infty$ for all $|\sigma| < \sigma_1$, hence (23) holds as stated in section 2.1. Since $\sigma \rightarrow a(\sigma)$ is continuous and $a(0) = 0$ is not an isolated blow-up point (otherwise, (7) can not hold), every $a(\sigma)$ is non isolated in S . Therefore, $1 \leq l_a < N = 2$ in (23), hence $l_a = 1$. (20) then gives the estimate for A_a , which gives the estimate for the eigenvalues.

ii) Since $(l_{a,1}^2 + l_{a,2}^2)^{1/2}$ is a norm for A_a , just evaluate the equation on A_a in Proposition 3.2 at eigenfunctions to get ii). This concludes the proof of Corollary 3.4. ■

At the point $a(0) = 0$, we have from Corollary 3.4,

$$\lambda_0(s) \sim -\frac{\beta}{s} \text{ and } \mu_0(s) = o\left(\frac{1}{s}\right) \text{ as } s \rightarrow \infty. \quad (27)$$

where λ_0 and μ_0 are just $l_{0,1}$ and $l_{0,2}$ renamed. This behavior is in fact stable with respect to σ . In the following proposition, we refine the estimates of Proposition 3.2 and state this stability result.

Proposition 3.5 (Stability of the behavior at infinity of the eigenvalues of $A_a(s)$) *There exists $\sigma_2 > 0$, $s_2 \in \mathbb{R}$ and $C_2 > 0$ such that for all $|\sigma| < \sigma_2$ and $s \geq s_2$,*

$$\begin{aligned} i) \quad & \|v_a(s) - v_{a,2}(s)\|_{L^2_\rho} \leq C_2 s^{-2}, & ii) \quad & |A'_a(s) - \frac{1}{\beta} A_a(s)^2| \leq C_2 s^{-3}, \\ iii) \quad & |\lambda_a(s) + \frac{\beta}{s}| \leq C_2 s^{-2} \log s & \text{and} & \quad |\mu_a(s)| \leq C_2 s^{-2}, \end{aligned}$$

where $\lambda_a = l_{a,\tau_a(1)}$, $\mu_a = l_{a,\tau_a(2)}$ and τ_a is a permutation of $\{1, 2\}$.

Let us first explain our argument for this proposition formally.

Up to the third order term, the eigenvalues satisfy the equation $\lambda' = \frac{1}{\beta} \lambda^2$, which has two orbits going to zero as $s \rightarrow \infty$:

$$\lambda_1(s) = -\frac{\beta}{s + s_0} \text{ and } \lambda_2(s) \equiv 0.$$

It is clear that λ_1 is stable, whereas λ_2 is not. Therefore, the stability of the behavior of λ_0 in (27) comes from the dynamical stability analysis of λ_1 . This argument was enough in [8] where all the eigenvalues were of order $-\frac{\beta}{s}$ (non degenerate profile). However, the stability analysis of λ_2 suggests that μ_a is not stable and does not allow us to derive the stability of its behavior. We need a new argument. λ_2 turns out to be *stable* if s is decreasing from ∞ to some point. Corollary 3.4 implies that one eigenvalue (the degenerate direction) of $A_a(s)$ is $o(\frac{1}{s})$ at infinity, say equal to $\lambda_2(s)$ at infinity, up to the order $o(\frac{1}{s})$. Thus, we recover the stability of the degenerate eigenvalue. We now give the actual proof.

Proof of Proposition 3.5 :

The proof is done in several steps. Let us sketch the main lemmas and derive the proposition first. Thus, we let the lemmas' proof to the end.

Let us fix $\hat{\epsilon} = \min\left(\frac{1}{2}, \frac{1}{100\beta}\right)$ and $s_3 = s_1(\hat{\epsilon})$ defined in Proposition 3.2. From (27) and the continuity of the set of eigenvalues with respect to a , we can find $\sigma_3 \in (0, \sigma_1)$ where σ_1 appears in Corollary 3.4, such that for all $|\sigma| \leq \sigma_3$,

$$|l_{a,\tau_a(1)}(s_3) + \frac{\beta}{s_3}| + |l_{a,\tau_a(2)}(s_3)| \leq \frac{\beta}{100s_3},$$

where τ_a is a permutation of $\{1, 2\}$. Let us rename the eigenvalues such that $\lambda_a = l_{a,\tau_a(1)}$ and $\mu_a = l_{a,\tau_a(2)}$. Therefore,

$$\forall |\sigma| \leq \sigma_3, \quad |\lambda_a(s_3) + \frac{\beta}{s_3}| \leq \frac{\beta}{100s_3} \text{ and } |\mu_a(s_3)| \leq \frac{\beta}{100s_3}. \quad (28)$$

We claim the following :

Lemma 3.6 (Non degeneracy of the decay rate of v_a) *There exists $C_3 > 0$ such that for all $|\sigma| < \sigma_3$ and $s \geq s_3$,*

$$i) N_a(s) \equiv \lambda_a^2 + \mu_a^2 \geq \frac{\beta^2}{16s^2}, \quad ii) \|v_a(s)\|_{L^2_\rho} \geq \frac{C_3}{s}.$$

We then prove the stability for the non degenerate direction.

Lemma 3.7 (Stability of the non degenerate direction of $A_a(s)$) *For all $|\sigma| < \sigma_3$ and $s \geq s_3$,*

$$-\frac{2\beta}{s} \leq \lambda_a(s) \leq -\frac{\beta}{2s} \text{ and } -\frac{2\beta}{s} \leq \mu_a(s) \leq \frac{C}{s}.$$

With this lemma, we can refine the equation satisfied by λ_a and μ_a .

Lemma 3.8 (A refined equation satisfied by $A_a(s)$) *There exists $s_4 \geq s_3$ and $C_4 > 0$ such that for all $|\sigma| < \sigma_3$ and $s \geq s_4$,*

$$\|v_a - v_{a,2}\|_{L^2_\rho} \equiv \left(\|v_{a,+}(s)\|_{L^2_\rho}^2 + \|v_{a,-}(s)\|_{L^2_\rho}^2 \right)^{\frac{1}{2}} \leq C_4 s^{-2} \quad (29)$$

$$|A'_a(s) - \frac{1}{\beta} A_a(s)^2| + |\lambda'_a - \frac{1}{\beta} \lambda_a^2| + |\mu'_a - \frac{1}{\beta} \mu_a^2| \leq C_4 s^{-3}. \quad (30)$$

Lemma 3.7 and Corollary 3.4 imply that for all $|\sigma| < \sigma_3$,

$$\mu_a(s) = O\left(s^{-1-\delta_a}\right) \text{ as } s \rightarrow \infty. \quad (31)$$

Equation (30) propagates this estimate from ∞ to s and improves it. More precisely,

Lemma 3.9 (Stability of the degenerate direction of $A_a(s)$) *There exist $s_5 \geq s_4$ and $C_5 > 0$ such that for all $|\sigma| < \sigma_3$ and $s \geq s_5$,*

$$|\mu_a(s)| \leq C_5 s^{-2},$$

With this information, we can refine the estimate on $\lambda_a(s)$.

Lemma 3.10 (Refinement of the estimate on the non degenerate direction of $A_a(s)$) *There exist $s_6 \geq s_5$, $\sigma_6 < \sigma_3$ and $C_6 > 0$ such that for all $|\sigma| < \sigma_6$ and $s \geq s_6$,*

$$|\lambda_a(s) + \frac{\beta}{s}| \leq C_6 \frac{\log s}{s^2}.$$

It is clear that Lemmas 3.8, 3.9 and 3.10 directly imply Proposition 3.5. Let us now prove the previous Lemmas.

Proof of Lemma 3.6 : Recall that $\hat{\epsilon}$, s_3 and σ_3 are defined just before (28).

i) From Corollary 3.4, we have for all $|\sigma| < \sigma_3$ and $s \geq s_3$,

$$N'_a(s) = 2(\lambda_a \lambda'_a + \mu_a \mu'_a) \geq \frac{2}{\beta}(\lambda_a^3 + \mu_a^3) - 2\hat{\epsilon}(\lambda_a + \mu_a)(\lambda_a^2 + \mu_a^2) \geq -\frac{6}{\beta}N_a^{3/2}$$

(here we used the fact that $\hat{\epsilon} \leq \frac{1}{100\beta}$ and $|\lambda_a^n + \mu_a^n| \leq 2(\lambda_a^2 + \mu_a^2)^{n/2}$).

Since $N_a(s_3) > \frac{\beta^2}{16s_3^2}$ (from (28)) and $\frac{d}{ds} \left(\frac{\beta^2}{16s^2} \right) < -\frac{6}{\beta} \left(\frac{\beta^2}{16s^2} \right)^{3/2}$, straightforward a priori estimates yield i).

ii) Since $\hat{\epsilon} \leq \frac{1}{2}$, Proposition 3.2 implies that $\|v_a\|_{L^2_\rho} \geq \frac{1}{2}\|v_{a,2}\|_{L^2_\rho} \geq C(\lambda_a^2 + \mu_a^2)^{1/2}$ where $C > 0$ (because $(\lambda_a^2 + \mu_a^2)^{1/2}$ is a norm for A_a , hence for $v_{a,2}$ by (21)). Thus, ii) of Lemma 3.6 follows from i). This concludes the proof of Lemma 3.6. \blacksquare

Proof of Lemma 3.7 :

We claim that for all $|\sigma| < \sigma_3$ and $s \geq s_3$,

$$\lambda_a(s) + \mu_a(s) < -\frac{\beta}{50s}. \quad (32)$$

Indeed, from Corollary 3.4, Lemma 3.6 and the fact that $\hat{\epsilon} \leq \frac{1}{100\beta}$, we have

$$\forall |\sigma| < \sigma_3, \forall s \geq s_3, \frac{d}{ds}(\lambda_a + \mu_a) \geq \left(\frac{1}{\beta} - 2\hat{\epsilon} \right) (\lambda_a^2 + \mu_a^2) \geq \frac{1}{2\beta} \frac{\beta^2}{16s^2}.$$

Since $\lambda_a(s) + \mu_a(s) \rightarrow 0$ as $s \rightarrow \infty$ (Corollary 3.4), an integration between s and ∞ gives (32).

(32) shows that Lemma 3.7 follows if we prove that for all $|\sigma| < \sigma_3$ and $s \geq s_3$,

$$-\frac{2\beta}{s} < \lambda_a(s) < -\frac{\beta}{2s} \text{ and } \mu_a(s) > -\frac{2\beta}{s}. \quad (33)$$

We proceed by contradiction. From (28), we consider some $|\sigma| < \sigma_3$ and $s_* > s_3$ such that (33) holds for all $s \in [s_3, s_*)$ with an equality case at s_* . In the following, we rule out those 3 cases of equality. Let us just mention that (33) and (32) yield

$$|\mu_a(s_*)| \leq \frac{2\beta}{s_*}. \quad (34)$$

Case 1 : $\lambda_a(s_*) = -\frac{\beta}{2s_*}$.

On one hand, we have $\lambda_a'(s_*) \geq \frac{d}{ds} \left(-\frac{\beta}{2s} \right) \Big|_{s=s_*} \geq \frac{\beta}{2s_*^2}$.

On the other hand, Corollary 3.4, (33) and (34) imply that

$$\begin{aligned} \lambda_a'(s_*) &\leq \frac{1}{\beta} \lambda_a(s_*)^2 + \hat{\epsilon} (\lambda_a(s_*)^2 + \mu_a(s_*)^2) \\ &\leq \frac{1}{\beta} \left(\frac{\beta}{2s_*} \right)^2 + \hat{\epsilon} \left(\left(\frac{2\beta}{s_*} \right)^2 + \left(\frac{2\beta}{s_*} \right)^2 \right) \leq \frac{\beta}{3s_*^2} \end{aligned}$$

because $\hat{\epsilon} \leq \frac{1}{100\beta}$. Contradiction.

Case 2 or 3 : $\lambda_a(s_*) = -\frac{2\beta}{s_*}$ or $\mu_a(s_*) = -\frac{2\beta}{s_*}$.

Let us handle for instance Case 3. Case 2 is exactly the same.

On one hand, we have $\mu_a'(s_*) \leq \frac{d}{ds} \left(-\frac{2\beta}{s} \right) \Big|_{s=s_*} \leq \frac{2\beta}{s_*^2}$.

On the other hand, Corollary 3.4, (33) and (34) imply that

$$\begin{aligned} \mu_a'(s_*) &\geq \frac{1}{\beta} \mu_a(s_*)^2 - \hat{\epsilon} (\lambda_a(s_*)^2 + \mu_a(s_*)^2) \\ &\geq \frac{1}{\beta} \left(\frac{2\beta}{s_*} \right)^2 - \hat{\epsilon} \left(\left(\frac{2\beta}{s_*} \right)^2 + \left(\frac{2\beta}{s_*} \right)^2 \right) \geq \frac{3\beta}{s_*^2} \end{aligned}$$

because $\hat{\epsilon} \leq \frac{1}{100\beta}$. Contradiction.

Thus, (33) holds for all $|\sigma| < \sigma_3$ and $s \geq s_3$. This concludes the proof of Lemma 3.7. \blacksquare

Proof of Lemma 3.9 :

An iteration argument for $\mu_a(s)$ based on (30) and (31) gives the result. Indeed, these estimates yield $\mu_a' = \beta^{-1} \mu_a^2 + O(s^{-3}) = O(s^{-(2+2\delta_a)}) + O(s^{-3})$ as $s \rightarrow \infty$.

If $2\delta_a \geq 1$, then $\mu_a = O\left(\frac{1}{s^2}\right)$. If $2\delta_a < 1$, then $\mu_a = O\left(\frac{1}{s^{1+2\delta_a}}\right)$. In this case, we repeat the same argument with $2\delta_a$ instead of δ_a until we get

$$\forall |\sigma| < \sigma_3, \quad \mu_a(s) = O\left(\frac{1}{s^2}\right) \text{ as } s \rightarrow \infty. \quad (35)$$

Fix $s_5 \geq s_4$ such that

$$\forall s \geq s_5, \quad \left(C_4 + \frac{1}{\beta}\right) \frac{1}{2s^2} < \frac{1}{s^{7/4}} \quad (36)$$

where s_4 and C_4 are defined in Lemma 3.8. From (35), we can define for all $|\sigma| < \sigma_3$,

$$s_\sigma^* = \min\{s^* \geq s_5 \mid \forall s \geq s^*, |\mu_a(s)| \leq s^{-7/4}\}. \quad (37)$$

Using (30), we have for all $s \in [s_\sigma^*, \infty)$,
 $|\mu_a'(s)| \leq \beta^{-1}|\mu_a(s)|^2 + C_4 s^{-3} \leq (C_4 + \beta^{-1})s^{-3}$.
Therefore,

$$\forall s \in [s_\sigma^*, \infty), \quad |\mu_a(s)| \leq (C_4 + \beta^{-1})s^{-2}/2 < s^{-7/4} \quad (38)$$

since $s_\sigma^* \geq s_5$ (see (36)). (37) then shows that $s_\sigma^* = s_5$ and (38) yields the result. \blacksquare

Proof of Lemma 3.8 : We just follow ideas due to Filippas, Kohn and Liu ([10], [11]). See Appendix B. \blacksquare

Proof of Lemma 3.10 : Let us define

$$Z_a(s) = s^2(\lambda_a(s) + \frac{\beta}{s}). \quad (39)$$

From (30) and Corollary 3.4, we have for all $|\sigma| < \sigma_3$,

$$Z_a(s) = O\left(s^{1-\delta_a}\right) \text{ as } s \rightarrow \infty, \quad \forall s \geq s_4, \quad |Z_a'(s) - \frac{Z_a^2}{\beta s^2}| \leq C_4 s^{-1}. \quad (40)$$

As for Lemma 3.9, we improve the estimate on Z_a iteratively.

From (40), we write $Z_a' = O(s^{-2\delta_a}) + O(s^{-1})$.

If $2\delta_a \geq 1$, then $Z_a = O(\log s)$. If $2\delta_a < 1$, then $Z_a(s) = O(s^{1-2\delta_a})$. We repeat the same argument with $2\delta_a$ instead of δ_a until we get

$$\forall |\sigma| < \sigma_3, \quad Z_a = O(\log s), \text{ hence } \lambda_a = -\frac{\beta}{s} + O\left(\frac{\log s}{s^2}\right)$$

as $s \rightarrow \infty$. We need to prove that this holds uniformly with respect to σ . Let us consider s_7 and $C_7 \geq 2C_4$ such that for all $s \geq s_7$, $|Z_0(s)| \leq C_7 \log s$ and $Z_a(s)$ is continuous in terms of $(a, s) \in S \times [s_7, \infty)$ (for this latter fact, remember from Lemma 3.3 the continuity of $\{\lambda_a(s), \mu_a(s)\}$ in terms of (a, s)). If s_7 is chosen so that $C_5 s_7^{-2} \leq \frac{\beta}{4} s_7^{-1}$, then $\lambda_a(s)$ and $\mu_a(s)$ become apart for $s \geq s_7$ by Lemmas 3.7 and 3.9. Therefore, both are continuous in terms of $(a, s) \in S \times [s_7, \infty)$.

Define $s_6 \geq s_7$ and then $\sigma_6 \leq \sigma_3$ such that

$$\forall s \geq s_6, \quad \frac{16C_7^2 \log^2 s}{\beta s^2} \leq \frac{C_4}{s} \text{ and } \forall |\sigma| \leq \sigma_6, \quad |Z_a(s_6)| \leq 2C_7 \log s_6 \quad (41)$$

We claim that

$$\text{for all } |\sigma| < \sigma_6 \text{ and } s \geq s_6, \quad |Z_a(s)| \leq 4C_7 \log s. \quad (42)$$

Indeed, if for some $|\sigma| < \sigma_6$ and $s \geq s_6$, then we have $|Z_a(s)| > 4C_7 \log s$, we can define from (41) s_σ^* such that

$$\forall s \in [s_6, s_\sigma^*], |Z_a(s)| \leq 4C_7 \log s \text{ and } |Z_a(s_\sigma^*)| = 4C_7 \log s_\sigma^*. \quad (43)$$

Using (40), (41), and the fact that $C_7 \geq 2C_4$, we have

$$\forall s \in [s_6, s_\sigma^*], |Z'_a(s)| \leq \frac{1}{\beta} \frac{Z_a^2}{s^2} + \frac{C_4}{s} \leq \frac{16C_7^2(\log s)^2}{\beta s^2} + \frac{C_4}{s} \leq \frac{2C_4}{s} \leq \frac{C_7}{s}.$$

Therefore, $|Z_a(s_\sigma^*)| \leq |Z_a(s_6)| + C_7(\log s_\sigma^* - \log s_6) \leq 3C_7 \log s_\sigma^*$ by (41). This contradicts (43). Thus, (42) holds. This closes the proof of Lemma 3.10 by (39). Thus Proposition 3.5 is proved. \blacksquare

Step 3 : Solution of the finite dimensional problem

Now, we are ready to solve (26). We claim the following :

Proposition 3.11 (Solution of the finite dimensional problem)

There exists $C_{10} > 0$ such that for all $b \in a(-\sigma_2, \sigma_2)$, there exists a 2×2 orthogonal matrix Q_b such that :

$$\text{for all } s \geq s_2, \left| A_a(s) + \frac{\beta}{s} L_a \right| \leq C_{10} \frac{\log s}{s^2} \quad (44)$$

where

$$L_a = Q_a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} Q_a^T. \quad (45)$$

Moreover, $Q_0 = \text{Id}$ and $b \in a(-\sigma_2, \sigma_2) \rightarrow Q_b$ is continuous.

Proof : It is easy to check from Proposition 3.5 that for all $|\sigma| < \sigma_2$ and $s \geq s_2$,

$$\begin{aligned} |A'_a(s) - \frac{\text{tr} A_a}{\beta} A_a(s) + \det A_a(s) \text{Id}| &\leq C_2 s^{-3}, \\ |A_a(s)| &\leq C s^{-1}, \quad |\text{tr} A_a + \frac{\beta}{s}| \leq C s^{-2} \log s, \quad |\det A_a| \leq C s^{-3}. \end{aligned}$$

Therefore, for all $|\sigma| < \sigma_2$ and $s \geq s_2$,

$$|A'_a + \frac{1}{s} A_a(s)| \leq C \frac{\log s}{s^3}, \text{ hence } \left| \frac{d}{ds} (s A_a(s)) \right| \leq C \frac{\log s}{s^2}.$$

This shows that $-\frac{s}{\beta} A_a(s)$ has a limit as $\rightarrow \infty$. This limit depends only on $a(\sigma)$ and not on σ , for $A_a(s)$ does the same (see (20)). Therefore, we call

this limit $L_{a(\sigma)}$. We define this way a function $b \in a(-\sigma_2, \sigma_2) \rightarrow L_b$. $L_{a(\sigma)}$ is a 2×2 symmetric matrix, such that for all $|\sigma| < \sigma_2$ and $s \geq s_2$,

$$|sA_a(s) + \beta L_a| \leq C \int_s^\infty t^{-2} \log t dt \leq Cs^{-1} \log s. \quad (46)$$

Since the convergence is uniform “with respect to $a(\sigma)$ ” and since for a fixed s , $A_a(s)$ is continuous with respect to a , $b \rightarrow L_b$ is continuous.

Since L_a is symmetric, it has 2 eigenvalues which are the limits as $s \rightarrow \infty$, of $-\frac{s}{\beta}\lambda_a$ and $-\frac{s}{\beta}\mu_a$, say 1 and 0, according to Proposition 3.5. Therefore, since $b \rightarrow L_b$ is continuous and L_b is symmetric with distinct eigenvalues, we can define a 2×2 orthogonal matrix Q_b , continuous in terms of b , such that (45) and then (44) hold (just define continuous eigenvectors). From i) of Corollary 3.4, we can even choose $Q_0 = \tilde{Q}_0$, hence, $Q_0 = \text{Id}$. \blacksquare

Step 4 : Asymptotic behavior of w_a in L_ρ^2

We prove Proposition 3.1 here. We first use the solution of the finite dimensional problem to find the asymptotic behavior of w_a as $s \rightarrow \infty$, in L_ρ^2 or equivalently uniformly on compact sets of \mathbb{R}^N . We then use techniques from [24] to extend the convergence up to sets of the type $\{|y| \leq K_0\sqrt{s}\}$.

Proof of Proposition 3.1 :

i) Take $\sigma_0 = \sigma_2$ and $s_0 = s_2$ where σ_2 and s_2 are defined in Proposition 3.5. Consider $|\sigma| < \sigma_0$ and $s \geq s_0$. With the change of variable $z = Q_a y$ and using (45), we have

$$\begin{aligned} & \left\| w_a(Q_a y, s) - \left\{ \kappa + \frac{\kappa}{2ps} \left(1 - \frac{y_1^2}{2}\right) \right\} \right\|_{L_\rho^2} \\ &= \left\| w_a(z, s) - \left\{ \kappa + \frac{\beta}{s} \left(1 - \frac{(Q_a^T z)_1^2}{2}\right) \right\} \right\|_{L_\rho^2} \quad (\beta = \frac{\kappa}{2p}) \\ &= \left\| v_a(z, s) - \left\{ -\frac{\beta}{2s} z^T L_a z + \frac{\beta}{s} \right\} \right\|_{L_\rho^2} \\ &\leq \|v_a(s) - v_{a,2}(s)\|_{L_\rho^2} + \left\| v_{a,2}(s) - \left\{ -\frac{\beta}{2s} z^T L_a z + \frac{\beta}{s} \right\} \right\|_{L_\rho^2} \equiv E_1 + E_2. \quad (47) \end{aligned}$$

According to Proposition 3.5, we have

$$E_1 = \|v_a(s) - v_{a,2}(s)\|_{L_\rho^2} \leq \frac{C_2}{s^2}. \quad (48)$$

Using (21) and (45), we have

$$E_2 = \left\| \frac{1}{2} z^T A_a(s) z - \text{tr} A_a(s) - \left\{ -\frac{\beta}{2s} z^T L_a z - \text{tr} \left(-\frac{\beta}{s} L_a \right) \right\} \right\|_{L_\rho^2}. \quad (49)$$

Therefore, we have from (44)

$$E_2 \leq C|A_a(s) + \frac{\beta}{s}L_a| \leq CC_{10} \frac{\log s}{s^2}. \quad (50)$$

Combining (47), (48) and (50) gives i) of Proposition 3.1.

ii) See Proposition 3.11.

iii) The derivation of iii) from i) was done by Velázquez in [24] for a fixed blow-up point a . However, in [24], the convergence speed was not given, because the error estimate in the L^2_ρ convergence was not that accurate there. We shall summarize in Appendix C the method of Velázquez, with a special care to the speed of convergence, and of course, to the uniformity with respect to the blow-up point. ■

4 Regularity of the blow-up set near a non isolated point with the behavior (4)

4.1 Continuous differentiability of S

We prove Theorem 1 in this subsection. We proceed in 2 steps :

- In Step 1, we derive from the stability of the blow-up behavior with respect to blow-up points in $\text{Im } a$ a sort of weak differentiability of S at points of $\text{Im } a$ (the cone property).

- In Step 2, we define a C^1 function A whose image is a graph and is equal to S in a neighborhood of the origin.

Step 1 : The cone property for $\text{Im } a$

Let us introduce the cone property first.

Definition 4.1 (Cone property and the weak tangent) *Consider a set $E \subset \mathbb{R}^2$.*

i) E is said to have the cone property at some $a \in E$ if there is $u \in \mathbb{S}^1$ such that for all $\epsilon > 0$, there is $\delta(a, \epsilon) > 0$ such that

$$E \cap B(a, \delta) \subset \Omega_{a,u,\epsilon} \equiv \{x \mid |(x-a) \cdot u| \geq (1-\epsilon)|x-a|\}. \quad (51)$$

$\mathbb{R}u$ is then called the weak tangent of E at a .

ii) E is said to have the uniform cone property at some subset $F \subset E$ if for all $\epsilon > 0$ and $a \in F$, E has the cone property at a with $\delta(a, \epsilon) = \delta(\epsilon)$.

Remark : $\Omega_{a,u,\epsilon}$ is a cone with vertex a . It shrinks to $a + \mathbb{R}u$ as $\epsilon \rightarrow 0$.

Remark : If E is a C^1 curve, then the cone property is equivalent to the differentiability and the weak tangent to the tangent.

Let us explain our argument first. $w_{a(\sigma)}$ defined in (12) describes the local behavior of u , near $a(\sigma)$. From iii) of Proposition 3.1, we see that if we travel along the direction $Q_{a(\sigma)}e_1$ from 0 to $y = \eta\sqrt{s}$ where $\eta > 0$, then we make $w_a(y, s)$ drop down from $f(0) = \kappa$ to $f(\eta) < \kappa$. No change occurs if we travel along $Q_{a(\sigma)}e_2$ (hence, we call it the degenerate direction). In the $u(x, t)$ variable, this means that when we travel along the non degenerate direction $Q_{a(\sigma)}e_1$, from a to $x = a + \eta e^{-\frac{s}{2}}\sqrt{s}$, $u(x, t)$ drops down from $v_T(t) \equiv \kappa(T - t)^{-\frac{1}{p-1}}$ to $(1 - \epsilon_0(\eta))v_T(t)$. Therefore, if s is large enough, all points along this non degenerate direction satisfy the blow-up exclusion criterion of Proposition 2.3. Thus, S is located along the degenerate direction $Q_{a(\sigma)}e_2$. More precisely, we have the following :

Proposition 4.2 (Uniform cone property for S at points of $\text{Im } a$)

- i) S has the uniform cone property at $\text{Im } a_{|\sigma| < \sigma_0}$. The weak tangent at $a(\sigma)$ is $\mathbb{R}Q_{a(\sigma)}e_2$ where $e_2 = (0, 1)$.
- ii) $Q_0 = \text{Id}$ and the weak tangent is continuous as a function of $b \in a(-\sigma_0, \sigma_0)$.

Remark : Velázquez's result in [24] implies that S has the cone property at $a(\sigma)$, but with no uniformity with respect of a .

Proof of Proposition 4.2 :

Note that ii) follows directly from i) of Proposition 4.2 and ii) of Proposition 3.1. Let us prove i). We need to prove that for all $\epsilon > 0$, there is $\delta(\epsilon)$ such that for all $|\sigma| < \sigma_0$, if

$$|x - a| < \delta \text{ and } |(x - a) \cdot Q_a e_2| < (1 - \epsilon)|x - a|, \quad (52)$$

then $x \notin S$. Consider ϵ and let us first introduce $\delta(\epsilon)$ and then show that it is convenient. Define

$$\epsilon_0 = \frac{1}{2} (\kappa - f(\sqrt{\epsilon})) > 0 \text{ and } t_0 = t_0(\epsilon_0) \quad (53)$$

as defined in Proposition 2.3. Consider then $s^*(\epsilon)$ such that

$$\forall s \geq s^*(\epsilon), C'_0(1) \frac{\log s}{s} \leq \epsilon_0 \quad (54)$$

where C'_0 is defined in Proposition 3.1. Define

$$\delta(\epsilon) = e^{-\frac{\tilde{s}}{2}}\sqrt{\tilde{s}} \text{ where } \tilde{s}(\epsilon) = \max(s_0 + 1, s^*(\epsilon), -\log(T - t_0)) \quad (55)$$

where s_0 is introduced in Proposition 3.1. Let us take any $|\sigma| < \sigma_0$ and x as in (52) and show that x is not a blow-up point. We will use the blow-up exclusion criterion of Proposition 2.3. Let us introduce $t_{a,x}$ and similarity variables such that

$$\begin{aligned} |x - a| &= \sqrt{(T - t_{a,x})|\log(T - t_{a,x})|}, \quad s_{a,x} = -\log(T - t_{a,x}), \\ y_{a,x} &= Q_a^{-1} \left(\frac{x-a}{\sqrt{T-t_{a,x}}} \right). \end{aligned} \quad (56)$$

The following lemma allows us to conclude.

Lemma 4.3

$$\begin{aligned} i) \quad s_{a,x} &\geq \max(s^*(\epsilon), -\log(T - t_0), s_0 + 1), & ii) \quad t_{a,x} &\geq t_0, \\ iii) \quad |y_{a,x}| &= \sqrt{s_{a,x}}, & iv) \quad |y_{a,x,2}| &\leq (1 - \epsilon)|y_{a,x}|, \\ v) \quad |y_{a,x,1}| &\geq \sqrt{\epsilon s_{a,x}}, & vi) \quad |u(x, t_{a,x})| &\leq \frac{\kappa - \epsilon_0}{(T - t_{a,x})^{\frac{1}{p-1}}}. \end{aligned}$$

Indeed, according to ii) and vi) of Lemma 4.3 and (53), x satisfies the blow-up exclusion criterion of Proposition 2.3 and is therefore not a blow-up point. Remains to prove Lemma 4.3.

Proof of Lemma 4.3 :

i) From (56), (52) and (55), we have $e^{-\frac{s_{a,x}}{2}}\sqrt{s_{a,x}} = |x - a| \leq \delta = e^{-\frac{\tilde{s}}{2}}\sqrt{\tilde{s}}$. Therefore, $s_{a,x} \geq \tilde{s}$. Use (55) again to conclude.

ii) Since $s_{a,x} = -\log(T - t_{a,x})$, use i) to conclude.

iii) From (56), we have $|y_{a,x}| = |x - a|/\sqrt{T - t_{a,x}} = \sqrt{|\log(T - t_{a,x})|} = \sqrt{s_{a,x}}$.

iv) From (52), we have $|(x - a) \cdot Q_a e_2| \leq (1 - \epsilon)|x - a|$. The conclusion follows since $Q_a y_{a,x} = (x - a)/\sqrt{T - t_{a,x}}$ by (56).

v) We have $y_{a,x,1}^2 = |y_{a,x}|^2 - y_{a,x,2}^2 \geq |y_{a,x}|^2(1 - (1 - \epsilon)^2)$ by iv). Since $\epsilon < 1$, the conclusion follows from iii).

vi) Using (12) and (56), we have $|u(x, t_{a,x})|$

$$= (T - t_{a,x})^{-\frac{1}{p-1}} \left| w_a \left(\frac{x-a}{\sqrt{T-t_{a,x}}}, s_{a,x} \right) \right| = (T - t_{a,x})^{-\frac{1}{p-1}} |w_a(Q_a y_{a,x}, s_{a,x})|.$$

From i), v), the monotonicity of f and Proposition 3.1, we have

$$|u(x, t_{a,x})| \leq (T - t_{a,x})^{-\frac{1}{p-1}} \left[f \left(\frac{y_{a,x,1}}{\sqrt{s_{a,x}}} \right) + C'_0(1) \frac{\log s_{a,x}}{s_{a,x}} \right]$$

$$\leq (T - t_{a,x})^{-\frac{1}{p-1}} [f(\sqrt{\epsilon}) + \epsilon_0] \text{ (use v), i) and (54)}$$

$$= (T - t_{a,x})^{-\frac{1}{p-1}} [\kappa - \epsilon_0] \text{ (use (53))}.$$

This concludes the proof of Lemma 4.3 and the proof of i) of Proposition 4.2. \blacksquare

Step 2 : S as the graph of a C^1 function

At the point $a(0) = 0$, we know from Proposition 4.2 that S is located along the degenerate direction $Q_a(0)e_2 = e_2$. In the following, we will show that $\text{Im } a$ is the graph of φ , function of the degenerate variable x_2 . Since at each point b of this graph, S is located along the degenerate direction $Q_b e_2$ which is continuous in terms of b , $S \setminus \text{graph } \varphi$ is empty, and φ is C^1 . Theorem 1 follows from the following :

Proposition 4.4 (S as the graph of a C^1 function)

i) There exist $\delta_1 > 0$ and $A \in C([-\delta_1, \delta_1], \mathbb{R}^2)$ such that

$$\text{Im } a = \text{Im } a|_{(-\sigma_0, \sigma_0)} \cap \tilde{B}(0, \delta_1) = S \cap \tilde{B}(0, \delta_1), \quad (57)$$

where $\tilde{B}(0, \delta_1) = \{(x_1, x_2) \mid \text{for } i = 1, 2, |x_i| \leq \delta_1\}$ and

$$\forall |x_2| \leq \delta_1, \quad A(x_2) \cdot e_2 = x_2. \quad (58)$$

ii) $A \in C^1([-\delta_1, \delta_1], \mathbb{R}^2)$ and the tangent to S at a point $A(\bar{x}_2)$ is $A(\bar{x}_2) + Q_{A(\bar{x}_2)} e_2 \mathbb{R}$.

Indeed, this implies that locally near $\hat{a}(0) = 0$, S is the graph of a C^1 function φ defined by

$$\forall |x_2| \leq \delta_1, \quad A(x_2) = (\varphi(x_2), x_2). \quad (59)$$

Therefore, (8) follows from (57), which yields the conclusion of Theorem 1. Let us prove Proposition 4.4.

Proof of Proposition 4.4 :

i) Consider $\eta_0 > 0$ such that the angle of Ω_{0, e_2, η_0} is less than $1/100$ and than α_0 where α_0 is defined in (7). From the uniform cone property of S at points of $\text{Im } a$ (Proposition 4.2), there exists $\delta_0 > 0$ such that $\forall |\sigma| < \sigma_0$, $S \cap B(a(\sigma), \delta_0) \subset \Omega_{a(\sigma), Q_{a(\sigma)} e_2, \eta_0}$.

Since $a(\cdot)$ and $Q_{a(\cdot)} e_2$ are continuous, $a(0) = 0$ and $Q_{a(0)} e_2 = e_2$ (see ii) of Proposition 3.1), there exists $\tilde{\sigma}_0 \leq \sigma_0$ such that for all $|\sigma| < \tilde{\sigma}_0$, $a(\sigma) \in B(0, \frac{\delta_0}{2})$, $\Omega_{a(\sigma), Q_{a(\sigma)} e_2, \eta_0} \subset \Omega_{a(\sigma), e_2, 2\eta_0}$ and $Q_{a(\sigma)} e_2 \cdot e_2 \neq 0$. Hence,

$$\forall |\sigma| < \tilde{\sigma}_0, \quad a(\sigma) \in B(0, \frac{\delta_0}{2}) \cap \Omega_{0, e_2, \eta_0}, \quad S \cap B(a(\sigma), \delta_0) \subset \Omega_{a(\sigma), e_2, 2\eta_0} \text{ and } Q_{a(\sigma)} e_2 \cdot e_2 \neq 0. \quad (60)$$

Let δ_- and δ_+ be the infimum and the supremum of $\sigma \rightarrow a(\sigma).e_2$ for $|\sigma| < \tilde{\sigma}_0$. Since 0 is not an endpoint in $\text{Im } a$ (property (7)), we have

$$-\frac{\delta_0}{2} \leq \delta_- < 0 < \delta_+ \leq \frac{\delta_0}{2}. \quad (61)$$

Indeed, $\delta_- \geq -\frac{\delta_0}{2}$ follows from (60). Moreover, if $\delta_- \geq 0$, then for all $|\sigma| < \tilde{\sigma}_0$, $a(\sigma).e_2 \geq 0$, hence $a(-\tilde{\sigma}_0, \tilde{\sigma}_0) \subset \Omega_{0,e_2,\eta_0} \cap \{x_2 \geq 0\}$ by (60). This contradicts (7), since the angle of Ω_{0,e_2,η_0} is less than α_0 . Do the same for δ_+ .

Now, we are ready to define A . If $\delta_1 = \frac{1}{2} \min(\delta_+, -\delta_-) > 0$, then for all $|x_2| \leq \delta_1$, there is $|\sigma^*(x_2)| \leq \tilde{\sigma}_0$ such that $a(\sigma^*(x_2)).e_2 = x_2$, by continuity of $\sigma \rightarrow a(\sigma).e_2$. If we define $A(x_2) = a(\sigma^*(x_2))$, then we readily see that (58) holds and if we use (60), then we get

$$\forall |x_2| \leq \delta_1, A(x_2) = a(\sigma^*(x_2)) \text{ with } |\sigma^*(x_2)| \leq \tilde{\sigma}_0 \text{ and } |A(x_2)| < \frac{\delta_0}{2}. \quad (62)$$

Let us prove that (57) holds. From (62) and (60), we have $\text{Im } A \subset \text{Im } a|_{(-\tilde{\sigma}_0, \tilde{\sigma}_0)} \subset \Omega_{0,e_2,\eta_0}$, whose angle is less than $\frac{1}{100}$. Since A is defined for all $|x_2| \leq \delta_1$, this implies that $\text{Im } A \subset \tilde{B}(0, \delta_1)$. Since $\text{Im } A \subset \text{Im } a|_{(-\tilde{\sigma}_0, \tilde{\sigma}_0)} \subset S$, (57) holds if we just show that $S \cap \tilde{B}(0, \delta_1) \subset \text{Im } A$, or that for all $b \in S \cap \tilde{B}(0, \delta_1)$, $b = A(b.e_2)$. Remark that for such a b , $|b.e_2| \leq \delta_1$ so that $A(b.e_2)$ is well defined. Using (62), we write $|b - A(b.e_2)| \leq |b| + |A(b.e_2)| \leq \sqrt{2}\delta_1 + \frac{\delta_0}{2} \leq \left(\frac{\sqrt{2}}{4} + \frac{1}{2}\right) \delta_0 < \delta_0$. Therefore, $b \in S \cap B(A(b.e_2), \delta_0) \subset \Omega_{A(b.e_2),e_2,2\eta_0}$ by (62) and (60). Since $b.e_2 = A(b.e_2).e_2$ by (58) and the line $x_2 = A(b.e_2).e_2$ intersects $\Omega_{A(b.e_2),e_2,2\eta_0}$ only at $A(b.e_2)$, this implies that $b = A(b.e_2)$. Thus, (57) holds.

Remains to prove that A is continuous. Consider $|x_{2,n}| \leq \delta_1$ such that $x_{2,n} \rightarrow \bar{x}_2 \in [-\delta_1, \delta_1]$ as $n \rightarrow \infty$. Since (62) implies that $A(x_{2,n}) \in S \cap \tilde{B}(0, \frac{\delta_0}{2})$, a compact set, we may assume that $A(x_{2,n}) \rightarrow b \in S \cap \tilde{B}(0, \frac{\delta_0}{2})$ (up to a subsequence). Let us show that $b = A(\bar{x}_2)$. Using (62), we have $|b - A(x_2)| < 2\frac{\delta_0}{2}$. Therefore, by (62) and (60), we have $b \in \Omega_{A(\bar{x}_2),e_2,2\eta_0}$ on one hand. On the other hand, we have from (58) $A(\bar{x}_2).e_2 = \bar{x}_2$ and $A(x_{2,n}).e_2 = x_{2,n}$, hence, $b.e_2 = \bar{x}_2 = A(\bar{x}_2).e_2$. Since the line $x_2 = A(\bar{x}_2).e_2$ intersects $\Omega_{A(\bar{x}_2),e_2,2\eta_0}$ only at $A(\bar{x}_2)$, this implies that $b = A(\bar{x}_2)$ and A is continuous. Thus, i) of Proposition 4.4 holds.

ii) Since A is continuous, we learn from the cone property at $A(\bar{x}_2)$ that we can make $\text{Im } A$ (that is the graph of φ defined in (59)) as close as we want

to the line $A(\bar{x}_2) + Q_{A(\bar{x}_2)}e_2$ by taking x_2 close enough to \bar{x}_2 . Therefore, this line is the tangent to the graph of φ at $A(\bar{x}_2)$ and φ is differentiable at \bar{x}_2 with $\varphi'(\bar{x}_2) = \frac{e_1 \cdot Q_{A(\bar{x}_2)}e_2}{e_2 \cdot Q_{A(\bar{x}_2)}e_2}$ (remember that $e_2 \cdot Q_{A(x_2)}e_2 \neq 0$ by (62) and (60)). Since A and $b \rightarrow Q_b$ are continuous (see ii) of Proposition 4.2), φ is C^1 . This closes the proofs of Proposition 4.4 and Theorem 1. \blacksquare

4.2 A geometric constraint yielding more regularity for S

We prove Proposition 2 in this subsection.

We first rewrite iii) of Proposition 3.1 with the terminology of Proposition 4.4. Thanks to (57), $a(\sigma)$ is viewed as $b \in \text{Im } A$. The variable y_1 is orthogonal to $Q_b e_2$, the tangent direction to S . Therefore, $y_1 = \pm d(y, T_b)$ where T_b is the tangent to S at b . More precisely, we have the following :

Corollary 4.5 *For all $K_0 > 0$, $t \geq t_0 \equiv T - e^{-s_0}$, $b \in \text{Im } A$ and $x \in \mathbb{R}^2$ such that $|x - b| \leq K_0 \sqrt{(T - t)|\log(T - t)|}$, we have*

$$\left| (T - t)^{\frac{1}{p-1}} u(x, t) - f \left(\frac{d(x, T_b)}{\sqrt{(T - t)|\log(T - t)|}} \right) \right| \leq C'_0(K_0) \frac{\log |\log(T - t)|}{|\log(T - t)|}$$

where T_b is the tangent to S at b .

Proof : If we introduce y and s such that

$$s = -\log(T - t) \text{ and } y = Q_b^T \left(\frac{x - b}{\sqrt{T - t}} \right), \quad (63)$$

then we see that $|y| = \frac{|x - b|}{\sqrt{T - t}} \leq K_0 \sqrt{s}$ and $s \geq s_0$. Since $b \in \text{Im } A \subset \text{Im } a_{|(-\sigma_0, \sigma_0)}$ by (57), we obtain from iii) of Proposition 3.1

$$\left| w_b(Q_b y, s) - f \left(\frac{y_1}{\sqrt{s}} \right) \right| \leq C'_0(K_0) \frac{\log s}{s}. \quad (64)$$

Remark that we have from (12)

$$w_b(Q_b y, s) = w_b \left(\frac{x - b}{\sqrt{T - t}}, -\log(T - t) \right) = (T - t)^{\frac{1}{p-1}} u(x, t). \quad (65)$$

From (63), we have $|y_1| = |y \cdot e_1| = \left| \frac{x - b}{\sqrt{T - t}} \cdot \tilde{Q}_b e_1 \right|$. Since $Q_b e_1$ is a normalized normal vector to S (see ii) of Proposition 4.4), we have $|(x - b) \cdot Q_b e_1| = d(x, T_b)$ where T_b is the tangent to S at b . Therefore,

$$|y_1| = \frac{d(x, T_b)}{\sqrt{T - t}}. \quad (66)$$

Combining this with (63), (64) and (65) concludes the Proof of Corollary 4.5. \blacksquare

Proof of Proposition 2 :

Given x near S ($= \text{Im } A$ locally), Corollary 4.5 gives different asymptotic behaviors for $u(x, t)$, depending on the choice of the point $b \in \text{Im } A \cap B(x, K_0 \sqrt{(T-t)|\log(T-t)|})$. All these possible behaviors have to agree, up to the error term in Corollary 4.5. This implies a geometric constraint on S , which gives some more regularity on A (and φ).

We consider some $|x_2| < \delta_1$ and some $h \in \mathbb{R}$ such that $|x_2 + h| < \delta_1$. Since A is C^1 , there is C^* such that

$$|\varphi'(x_2)| \leq C^* \text{ and } |A(x_2 + h) - A(x_2)| \leq C^*|h|. \quad (67)$$

For any time $t > t_0$ such that $|A(x_2) - A(x_2 + h)| \leq \sqrt{(T-t)|\log(T-t)|}$, we can estimate $u(A(x_2 + h), t)$ from Corollary 4.5 in two ways :

- First by taking $x = b = A(x_2 + h)$, which gives

$$\left| (T-t)^{\frac{1}{p-1}} u(A(x_2 + h), t) - f(0) \right| \leq C'_0(1) \frac{\log |\log(T-t)|}{|\log(T-t)|}. \quad (68)$$

- Second, by taking $b = A(x_2)$, $x = A(x_2 + h)$, which gives

$$\left| (T-t)^{\frac{1}{p-1}} u(A(x_2 + h), t) - f\left(\frac{d(A(x_2 + h), T_{A(x_2)})}{\sqrt{(T-t)|\log(T-t)|}}\right) \right| \leq C'_0(1) \frac{\log |\log(T-t)|}{|\log(T-t)|}. \quad (69)$$

Now, if we fix $t = t(x_2, h)$ such that

$$|A(x_2 + h) - A(x_2)| = \sqrt{(T-t(x_2, h))|\log(T-t(x_2, h))|} \quad (70)$$

and take $|h| < h_1(t_0)$ for some $h_1(t_0) > 0$, we see from (67) that $t(x_2, h) \geq t_0$, hence (68) and (69) hold. Therefore,

$$\left| f(0) - f\left(\frac{d(A(x_2 + h), T_{A(x_2)})}{|A(x_2 + h) - A(x_2)|}\right) \right| \leq 2C'_0(1) \frac{\log |\log(T-t(x_2, h))|}{|\log(T-t(x_2, h))|}. \quad (71)$$

Note that since $A(x_2) \in T_{A(x_2)}$, we have $\frac{d(A(x_2 + h), T_{A(x_2)})}{|A(x_2 + h) - A(x_2)|} \leq 1$. Therefore, (9) implies that there is $C > 0$ such that

$$C \left[\frac{d(A(x_2 + h), T_{A(x_2)})}{|A(x_2 + h) - A(x_2)|} \right]^2 \leq \left| f(0) - f\left(\frac{d(A(x_2 + h), T_{A(x_2)})}{|A(x_2 + h) - A(x_2)|}\right) \right| \quad (72)$$

Since $\text{Im } A$ is the graph of φ , we have

$$d(A(x_2 + h), T_{A(x_2)}) = \frac{|\varphi(x_2 + h) - \varphi(x_2) - h\varphi'(x_2)|}{\sqrt{1 + \varphi'(x_2)^2}}. \quad (73)$$

If $\tau(d)$ is given by $d = \sqrt{\tau|\log \tau|}$, then

$$\log \tau \sim 2 \log d \text{ and } \log |\log \tau| \sim \log |\log d| \text{ as } d \rightarrow 0.$$

Therefore, $\frac{\log |\log \tau|}{|\log \tau|} \leq \frac{\log |\log d|}{|\log d|}$ if $|d| \leq d_0$ for some $d_0 > 0$. Combining this with (70) and (67), we have for all $|x_2| < \delta_1$ and $|h| < h_0$ for some $h_0 > 0$ such that $|x_2 + h| \leq \delta_1$,

$$\frac{\log |\log(T - t(x_2, h))|}{|\log(T - t(x_2, h))|} \leq \frac{\log |\log |A(x_2 + h) - A(x_2)||}{|\log |A(x_2 + h) - A(x_2)||} \leq C \frac{\log |\log |h||}{|\log |h||}. \quad (74)$$

Combining (72), (73), (71), (74) and (67) closes the proof of Proposition 2. \blacksquare

5 Blow-up profile at a non isolated blow-up point with the behavior (4)

We prove Theorem 3 in this section.

Step 1 : Asymptotic behavior in self similar variables around the blow-up set

We prove (10) in this step. This follows from Corollary 4.5 by taking $b = P_S(x)$, the orthogonal projection of x on S . Indeed, take $t \geq t_0$ and $x \in B(0, \delta)$ such that $d(x, S) \leq K_0 \sqrt{(T-t)|\log(T-t)|}$. We define $P_S(x)$ as the orthogonal projection of x on S . We claim that $P_S(x) \in B(0, 2\delta)$, hence $P_S(x) \in \text{graph } \varphi = \text{Im } A$ by (8). Indeed, $|x - P_S(x)| \leq |x - 0|$ since $0 \in S$, therefore, $|P_S(x)| \leq |P_S(x) - x| + |x| \leq 2|x| < 2\delta$. Since $|x - P_S(x)| = d(x, S) \leq K_0 \sqrt{(T-t)|\log(T-t)|}$ and $d(x, T_{P_S(x)}) = d(x, S)$, (10) follows directly from Corollary 4.5 applied with $b = P_S(x)$.

Step 2 : Limiting profile in the original set of variables, near the blow-up set

We prove (11) here. This follows from (10) and the uniform ODE comparison property of Proposition 2.2. Define $t(d)$ such that

$$d = \sqrt{(T - t(d))|\log(T - t(d))|}. \quad (75)$$

For all $x \in B(0, \delta) \setminus S$ and $(\xi, \tau) \in \mathbb{R}^N \times [-\frac{t(d(x, S))}{T-t(d(x, S))}, 1)$, we introduce

$$v(x, \xi, \tau) = (T - \tilde{t})^{\frac{1}{p-1}} u \left(x + \xi \sqrt{T - \tilde{t}}, t + \tau(T - \tilde{t}) \right) \quad (76)$$

where $\tilde{t} = t(d(x, S))$. From (10), (76) and Proposition 2.2, we know that there is $\epsilon_0 > 0$ such that for all $\epsilon > 0$, there is C_ϵ such that if $d(x, S) < \epsilon_0$ and $(\xi, \tau) \in \mathbb{R}^N \times [-\frac{t(d(x, S))}{T-t(d(x, S))}, 1)$, then

$$\begin{aligned} |\partial_\tau v - |v|^{p-1}v| &\leq \epsilon |v|^p + C_\epsilon (T - t(d(x, S)))^{\frac{p}{p-1}} \\ |v(x, 0, 0) - f(1)| &\leq C'_0(1) \frac{\log |\log(T - t(d(x, S)))|}{|\log(T - t(d(x, S)))|}. \end{aligned}$$

Therefore, for all $\epsilon > 0$, there is $\eta(\epsilon) > 0$ such that if $d(x, S) < \eta$, then

$$|\partial_\tau v - |v|^{p-1}v| \leq \epsilon (|v|^p + 1) \text{ and } |v(x, 0, 0) - f(1)| \leq \epsilon. \quad (77)$$

This implies that

$$\sup_{\tau \in [0, 1]} |v(x, 0, \tau) - v_0(\tau)| \rightarrow 0 \text{ as } d(x, S) \rightarrow 0 \quad (78)$$

where
$$v_0(\tau) = \left((p-1)(1-\tau) + \frac{(p-1)^2}{4p} \right)^{-\frac{1}{p-1}} \quad (79)$$

is the solution of $v'_0(\tau) = v_0(\tau)^p$, $v_0(0) = f(1)$, defined in particular for all $\tau \in [0, 1]$. Moreover, $\sup_{\tau \in [0, 1]} |\partial_\tau v(x, 0, \tau)| \leq 2 \sup_{\tau \in [0, 1]} |\partial_\tau v_0(\tau)|$ for $d(x, S)$ small. Therefore, for $d(x, S)$ small, $v(x, 0, \tau)$ has a limit as $\tau \rightarrow 1$, hence (76) implies that $u(x, t)$ has a limit $u^*(x)$ as $t \rightarrow T$. Using (78) and (76), we see that

$$u^*(x) \sim (T - t(d(x, S)))^{-\frac{1}{p-1}} v_0(1) \text{ as } d(x, S) \rightarrow 0. \quad (80)$$

We claim that

$$(T - t(d))^{-\frac{1}{p-1}} v_0(1) \sim \left(\frac{d^2}{|\log d|} \frac{(p-1)^2}{8p} \right)^{-\frac{1}{p-1}} \text{ as } d \rightarrow 0. \quad (81)$$

Indeed, $v_0(1) = \left(\frac{(p-1)^2}{4p} \right)^{-\frac{1}{p-1}}$ from (79), $\log(T - t(d)) \sim 2 \log d$ and $T - t(d) \sim \frac{d^2}{2|\log d|}$ as $d \rightarrow 0$ from (75). (11) then follows from (80) and (81). This closes the proof of Theorem 3. \blacksquare

6 The higher dimensional case

We sketch the proof of Theorem 4 here. We need to review the proofs of Theorems 1, 3 and Proposition 2 to adapt them to the new context. We shall stress the most delicate points in the adaptation of Theorem 1 and Proposition 2. Once this is done, Theorem 3 extends in a natural way to higher dimensions, so we don't discuss it here.

If $N \geq 3$, we consider a non isolated blow-up point \hat{a} where u has the behavior (4) with $l < N$. We may take $\hat{a} = 0$ and $\tilde{Q}_{\hat{a}} = \text{Id}$. According to [24], S satisfies near \hat{a} the $(N - l)$ -cone property :

Definition 4.1' (**$(N - l)$ -cone property and weak tangent plane**)
Consider a set $E \subset \mathbb{R}^N$.

i) E is said to have the $(N - l)$ -cone property at some point $a \in E$ if there is π a $(N - l)$ subspace such that for all $\epsilon > 0$, there is $\delta(a, \epsilon) > 0$ such that

$$E \cap B(a, \delta) \subset \Omega_{a, \pi, \epsilon} \equiv \{x \mid |P_{\pi}(x - a)| \geq (1 - \epsilon)|x - a|\}$$

where P_{π} is the orthogonal projection over π . π is then called the weak tangent plane of E at a .

ii) E is said to have the uniform $(N - l)$ -cone property at some subset $F \subset E$ if for all $\epsilon > 0$ and $a \in F$, E has the $(N - l)$ -cone property at a with $\delta(a, \epsilon) = \delta(\epsilon)$.

Remark : $\Omega_{a, \pi, \epsilon}$ is a cone with vertex a . It shrinks to $a + \pi$ as $\epsilon \rightarrow 0$. We have the following consequence of [24] :

Lemma 6.1 (Velázquez, $(N - l_a)$ -cone property for S near a point with the behavior (4)) *If u has the behavior (4) (or (23)) near a , then S satisfies the $(N - l)$ -cone property at a . The weak tangent plane is spanned by $\tilde{Q}_a e_j$, $j = l_a + 1, \dots, N$.*

Proof : See Theorem 2 in [24]. ■

Therefore, S has a weak tangent plane at $\hat{a} = 0$, spanned by e_j , $j = l + 1, \dots, N$. One would expect S to be locally of dimension $N - l$ near \hat{a} . However, we are unable to prove that S is a continuum near \hat{a} . Therefore, we assume there is $a \in C((-1, 1)^{N-l}, \mathbb{R}^N)$ such that $a(0) = \hat{a} = 0$ and $\text{Im } a \subset S$ where $\text{Im } a$ is at least $(N - l)$ dimensional in the sense that

$$\forall b \in \text{Im } a, \text{ there are } (N - l) \text{ independent vectors } v_1, \dots, v_{N-l} \text{ in } \mathbb{R}^N \text{ and } a_1, \dots, a_{N-l} \text{ functions in } C^1([0, 1], S) \text{ such that } a_i(0) = b \text{ and } a_i'(0) = v_i. \quad (82)$$

This hypothesis means that b is actually non isolated in $(N-l)$ independent directions. We also assume that $\hat{a} = 0$ is not an endpoint in $\text{Im } a$ in the sense that

$$\forall \epsilon > 0, \text{ the projection of } a((-\epsilon, \epsilon)^{N-l}) \text{ on the weak tangent plane} \quad (83)$$

at $\hat{a} = 0$ contains an open ball with center $\hat{a} = 0$.

Let us first show the stability result of section 3 in the case $N \geq 3$.

6.1 Stability of the behavior (4)

Since we have taken $\hat{a} = 0$ and $\tilde{Q}_{\hat{a}} = \text{Id}$, (4) implies that

$$\sup_{|z| \leq K_0} \left| (T-t)^{\frac{1}{p-1}} u(z\sqrt{(T-t)|\log(T-t)|}, t) - f(|z'|) \right| \rightarrow 0 \text{ as } t \rightarrow T$$

where f is defined in (9) and $z = (z_1, z_2, \dots, z_N) \equiv (z', z_{l+1}, \dots, z_N)$. Section 2 then implies that

$$v_0 \sim v_{0,2} \text{ and } w_0(y, s) - \kappa = v_0(y, s) \sim \frac{\kappa}{2ps} \left(l - \frac{|y'|^2}{2} \right) \text{ as } s \rightarrow \infty \quad (84)$$

where $y' = (y_1, \dots, y_l)$. We claim that Proposition 3.1 holds here (with the obvious changes $a(-\sigma_0, \sigma_0) \rightarrow a((-\sigma_0, \sigma_0)^{N-l})$, $2 \times 2 \rightarrow N \times N$ and

$$y_1^2 \rightarrow \sum_{i=1}^l y_i^2).$$

Proof of Proposition 3.1 in higher dimensions :

Let us follow the 4 steps of the proof given in section 3.

Step 1 : Uniform reduction to a finite dimensional problem

This step holds as it is in section 2.

Step 2 : A spectral study of the finite dimensional problem

We should define N eigenvalues of $A_a(s)$, $l_{a,i}(s)$, C^1 as functions of s . The multi-function $\{l_{a,i}(s), i = 1, \dots, N\}$ is continuous in terms of $(a, s) \in S \times [-\log T, \infty)$. Corollary 3.4 has to be changed formally. The following Proposition crucially uses the $(N-l)$ dimensionality property of $\text{Im } a$.

Corollary 3.4' (Higher dimensional version of Corollary 3.4)

i) (Non uniform behavior of v_a) For all $|\sigma| < \sigma_1$, (23) holds with $l_a \leq l$. In particular, $A_a(s) = -\frac{\beta}{s} \tilde{Q}_a I_l \tilde{Q}_a^T + O(s^{-1-\delta_a})$ where I_l is a $N \times N$ diagonal matrix with $I_{l,ii} = 1$ if $i \leq l_a$ and $I_{l,ii} = 0$ if $i \geq l_a + 1$. Moreover, $A_a(s)$ has $(N-l_a)$ (hence, at least $(N-l)$) eigenvalues equal to $O(s^{-1-\delta_a})$, while the

others are equal to $-\frac{\beta}{s} + O(s^{-1-\delta_a})$.

ii) **(Equations on eigenvalues)** For all $\epsilon > 0$, there is $s_1(\epsilon)$ such that for all $|\sigma| < \sigma_1$, $i = 1, \dots, N$ and $s \geq s_1(\epsilon)$,

$$|l'_{a,i}(s) - \frac{1}{\beta} l_{a,i}(s)^2| \leq \epsilon \sum_{j=1}^N l_{a,j}(s)^2$$

where $l_{a,j}(s)$ are the eigenvalues of $A_a(s)$.

Proof: i) From Proposition 3.1, we have $v_a \sim v_{a,2}$ as $s \rightarrow \infty$ for all $|\sigma| < \sigma_1$, hence (23) holds as stated in section 2.1. Lemma 6.1 then implies that u satisfies the $(N - l_a)$ -cone property at a and therefore, S has a $(N - l_a)$ dimensional weak tangent plane at a . This plane contains the $(N - l)$ independent directions in property (82). Therefore, $l_a \leq l$. The estimate for A_a then follows from (20), and gives the estimate for eigenvalues.

ii) See the proof of ii) of Corollary 3.4. ■

At the point $a(0) = 0$, we know from (84) that $l_0 = l$. Using Corollary 3.4', we can rename the eigenvalues of $A_0(s)$ such that

$$\forall i = 1, \dots, l, \lambda_{0,i}(s) \sim -\frac{\beta}{s} \text{ and } \forall j = 1, \dots, N - l, \mu_{0,j}(s) = o\left(\frac{1}{s}\right) \text{ as } s \rightarrow \infty. \quad (85)$$

We claim then that Proposition 3.5 holds with eigenvalues $\lambda_{a,i}$ for $i = 1, \dots, l$ and $\mu_{a,j}$ for $j = 1, \dots, N - l$ instead of (λ_a, μ_a) .

Proof of Proposition 3.5 in higher dimensions : We should fix a new $\hat{\epsilon} > 0$ and take $s_3 = s_1(\hat{\epsilon})$ defined in Proposition 3.2. From (85) and the continuity of eigenvalues with respect to a , we can find $\sigma_3 \in (0, \sigma_1)$ where σ_1 appears in Corollary 3.4, such that

$$\forall |\sigma| \leq \sigma_3, \quad \begin{aligned} \forall i = 1, \dots, l, \quad & |l_{a,\tau_a(i)}(s_3) + \frac{\beta}{s_3}| \leq \frac{\beta}{100s_3}, \\ \forall i = l + 1, \dots, N, \quad & |l_{a,\tau_a(i)}(s_3)| \leq \frac{\beta}{100s_3} \end{aligned}$$

where τ_a is a permutation of $\{1, \dots, N\}$. If we rename the eigenvalues such that $\lambda_{a,i}(s) = l_{a,\tau_a(i)}$ for all $i = 1, \dots, l$ and $\mu_{a,j} = l_{a,\tau_a(l+j)}$ for all $j = 1, \dots, N - l$, then we get

$$\forall |\sigma| \leq \sigma_3, \quad \begin{aligned} \forall i = 1, \dots, l, \quad & |\lambda_{a,i}(s_3) + \frac{\beta}{s_3}| \leq \frac{\beta}{100s_3}, \\ \forall j = 1, \dots, N - l, \quad & |\mu_{a,j}(s_3)| \leq \frac{\beta}{100s_3}. \end{aligned}$$

Lemma 3.6 and its proof hold with the change $N_a(s) = \sum_{i=1}^l \lambda_{a,i}^2 + \sum_{j=1}^{N-l} \mu_{a,j}^2$.

Lemmas 3.7 and 3.8 then hold, with (λ_a, μ_a) replaced by $(\lambda_{a,i}, \mu_{a,j})$ and different constants. Here comes a delicate point before Lemma 3.9 : we need to prove that (31) holds for all $\mu_{a,j}$, for $j = 1, \dots, N-l$. This comes from two arguments. On one hand, we know from Corollary 3.4' i) that $A_a(s)$ has at least $N-l$ degenerate eigenvalues (that is eigenvalues satisfying (31)). On the other hand, we see from Lemma 3.7 that all degenerate eigenvalues must be in the set of all $\mu_{a,j}$, $j = 1, \dots, N-l$, which contains $(N-l)$ elements. Therefore, all $\mu_{a,j}$ are degenerate and satisfy (31).

One can easily see that Lemmas 3.9 and 3.10 extend naturally to $N \geq 3$ and hold for each $\mu_{a,j}$ and $\lambda_{a,i}$. ■

Step 3 : Solution of the finite dimensional problem

Proposition 3.11 holds with “ $N \times N$ ” instead of “ 2×2 ” and $L_a = Q_a I_l Q_a^T$ where I_l is a $N \times N$ diagonal matrix with $I_{l,ii} = 1$ if $i \leq l$ and $I_{l,ii} = 0$ if $i \geq l+1$.

Proof of Proposition 3.11 for $N \geq 3$: If $N \geq 3$, one needs to find the asymptotic behavior of all principal minors of order k of the matrix $A_a(s)$ with $k = N, \dots, 1$, before getting the asymptotic behavior of the coefficients of $A_a(s)$ (property (46)). This has been done by Filippas and Liu [11], Proposition 5.1. The continuity of $b \in a((-\sigma_0, \sigma_0)^{N-l}) \rightarrow L_b$ follows from (46) as in the case $N = 2$. Since L_a is symmetric, it has N eigenvalues which are the limits as $s \rightarrow \infty$ of $-\frac{s}{\beta} \lambda_{a,i}$ ($i = 1, \dots, l$) and $-\frac{s}{\beta} \mu_{a,j}$ ($j = 1, \dots, N-l$), say 1 (multiplicity l) and 0 (multiplicity $N-l$), according to Proposition 3.5. Since $b \rightarrow L_b$ is continuous and the multiplicities of eigenvalues 0 and 1 are independent of b , we can define continuous eigenvectors for L_b (see Oustry and Overton [22], corollary 2.5). Therefore, we can define a $N \times N$ orthogonal matrix Q_b , continuous in terms of b , diagonalizing L_b . This concludes the proof of Propositions 3.11 in the higher dimensional case. ■

Step 4 : Asymptotic behavior of w_a in L_ρ^2

This step extends naturally to $N \geq 3$, which concludes the proof of Proposition 3.1. ■

6.2 Regularity of the blow-up set

We prove here the part of Theorem 4 equivalent to Theorem 1. For this, we adapt section 4 to the case $N \geq 3$.

Step 1 : The $(N-l)$ -cone property for $\text{Im } a$

We claim the following :

Proposition 4.2' (Uniform $(N - l)$ -cone property for S at points of $\text{Im } a$)

i) S has the uniform $(N - l)$ -cone property at $a \left((-\sigma_0, \sigma_0)^{N-l} \right)$. The weak tangent plane $\pi_{a(\sigma)}$ at $a(\sigma)$ is spanned by $Q_{a(\sigma)} e_j$, $j = l + 1, \dots, N$, where e_j is the j -th vector of the canonical basis of \mathbb{R}^N .

ii) $Q_0 = \text{Id}$ and the weak tangent plane has an orthogonal basis $(Q_b e_j, j = l + 1, \dots, N)$ continuous as a function of $b \in a \left((-\sigma_0, \sigma_0)^{N-l} \right)$.

Remark : If $j \geq l + 1$, then $Q_b e_j$ is a degenerate direction in the asymptotic expansion of iii) in Proposition 3.1. If $j \leq l$, then $Q_b e_j$ is a non degenerate direction.

Proof : ii) follows directly from i) by ii) of Proposition 3.1. The proof of i) in the case $N = 2$ extends naturally to the case $N \geq 3$. Just note that $y_{a,x,1}^2$

and $y_{a,x,2}^2$ should be replaced respectively by $\sum_{i=1}^l y_{a,x,i}^2$ and $\sum_{j=l+1}^N y_{a,x,j}^2$. ■

Step 2 : S as the graph of a C^1 function.

The part of Theorem 4 equivalent to Theorem 1 follows from the following :

Proposition 4.4' (S as the graph of a C^1 function)

i) There exist $\delta_1 > 0$ and $A \in C([- \delta_1, \delta_1]^{N-l}, \mathbb{R}^N)$ such that

$$\text{Im } A = a \left((-\sigma_0, \sigma_0)^{N-l} \right) \cap \tilde{B}(0, \delta_1) = S \cap \tilde{B}(0, \delta_1), \quad (86)$$

where $\tilde{B}(0, \delta_1) = \{x \mid \forall i = 1, \dots, N, |x_i| \leq \delta_1\}$ and

$$\forall (x_{l+1}, \dots, x_N) \in [-\delta_1, \delta_1]^{N-l}, \forall j = l + 1, \dots, N, A(x_{l+1}, \dots, x_N) \cdot e_j = x_j \quad (87)$$

ii) $A \in C^1([- \delta_1, \delta_1]^{N-l}, \mathbb{R}^N)$ and the tangent plane to S at a point $b \in S \cap \tilde{B}(0, \delta_1)$ is π_b , spanned by $Q_b e_j$, $j = l + 1, \dots, N$.

Proof : The “no-end-point” property of $\hat{a} = 0$ in $\text{Im } a$ stated in (83) is apparently different from (7) stated for $N = 2$. That is why we should carefully define A here. Once this is done, one should follow the case $N = 2$ to finish the proof.

From the uniform cone property of S at points of $\text{Im } a$ (Proposition 4.2'), there exists $\delta_0 > 0$ such that $\forall |\sigma| < \sigma_0$, $S \cap B(a(\sigma), \delta_0) \subset \Omega_{a(\sigma), \pi_{a(\sigma)} e_2, 1/4}$. Since $a(\cdot)$ and $Q_{a(\cdot)}$ are continuous, $a(0) = 0$ and $Q_{a(0)} = \text{Id}$ (see ii) of Proposition 3.1), there exists $\tilde{\sigma}_0 \leq \sigma_0$ such that for all $|\sigma| < \tilde{\sigma}_0$, $a(\sigma) \in$

$B(0, \frac{\delta_0}{N})$, $\Omega_{a(\sigma), Q_{a(\sigma)} e_2, 1/4} \subset \Omega_{a(\sigma), e_2, 1/3}$ and $\det_{(e_k, k=l+1, \dots, N)}(Q_{a(\sigma)} e_j, j = l+1, \dots, N) \neq 0$. Hence, $\forall |\sigma| < \tilde{\sigma}_0$,

$$\begin{aligned} a(\sigma) \in B(0, \frac{\delta_0}{N}) \cap \Omega_{0, e_2, 1/4}, \quad S \cap B(a(\sigma), \delta_0) \subset \Omega_{a(\sigma), e_2, 1/3} \quad \text{and} \\ \det_{(e_k, k=l+1, \dots, N)}(Q_{a(\sigma)} e_j, j = l+1, \dots, N) \neq 0. \end{aligned} \quad (88)$$

Since $Q_{a(0)} = \text{Id}$, we learn from Proposition 4.2' that the weak tangent plane at $a(0) = 0$ is π_0 , spanned by e_j , $j = l+1, \dots, N$. Using property (83), we find $\delta_1(\tilde{\sigma}_0) > 0$ such that $\tilde{B}_{\pi_0}(0, \delta_1) \subset P_{\pi_0}(a[-\tilde{\sigma}_0, \tilde{\sigma}_0]^{N-l})$, where

$$\tilde{B}_{\pi_0}(0, \delta_1) = \{x \in \pi_0 \mid |x_j| \leq \delta_1, \text{ for } j = l+1, \dots, N\}.$$

This implies that A can be defined on $\tilde{B}_{\pi_0}(0, \delta_1)$, so that (87) holds. One can then finish the proof of Proposition 4.4' by just following the proof of Proposition 4.4. \blacksquare

6.3 A geometric constraint yielding some more regularity

This subsection is dedicated to the higher dimensional version of Proposition 2. Note that from Proposition 4.4', $\text{Im } A$ is the graph of $\varphi \in C^1([- \delta_1, \delta_1]^{N-l}, \mathbb{R}^l)$ such that $\forall (x_{l+1}, \dots, x_N) \in [- \delta_1, \delta_1]^{N-l}$, $A(x_{l+1}, \dots, x_N) = (\varphi_1, \dots, \varphi_l, x_{l+1}, \dots, x_N)$ where $\varphi_i = \varphi_i(x_{l+1}, \dots, x_N)$. Proposition 2 holds with $\varphi(x_2)$ replaced by $\varphi_i(x_{l+1}, \dots, x_N)$ and φ' by $\nabla \varphi_i$.

Proof of Proposition 2 in higher dimensions : Just follow the case $N = 2$ with $A(x_2)$ replaced by $A(\tilde{x})$ where $\tilde{x} = (x_{l+1}, \dots, x_N)$. The only delicate point is the equivalent of (73) which is : for all $i = 1, \dots, l$,

$$d\left(A(\tilde{x} + \tilde{h}), \pi_{A(\tilde{x})}\right) \geq \frac{\left| \varphi_i(\tilde{x} + \tilde{h}) - \varphi_i(\tilde{x}) - \tilde{h} \cdot \nabla \varphi_i(\tilde{x}) \right|}{\sqrt{1 + |\nabla \varphi_i(\tilde{x})|^2}}. \quad (89)$$

Note that we just need this inequality in our argument.

To prove (89), just note that $\text{Im } A \subset S_i$, the surface of equation $x_i = \varphi_i(x_{l+1}, \dots, x_N)$. Hence, $\pi_{A(\tilde{x})} \subset \pi_{i, A(\tilde{x})}$, the $(N-1)$ dimensional tangent plane to S_i at $A(\tilde{x})$. Therefore, $d\left(A(\tilde{x} + \tilde{h}), \pi_{A(\tilde{x})}\right) \geq d\left(A(\tilde{x} + \tilde{h}), \pi_{i, A(\tilde{x})}\right)$, equal to the right hand side of (89) (standard calculations).

A Reduction of the asymptotic blow-up behavior to a finite dimensional problem

We prove Proposition 3.2 here. The use of the Liouville Theorem is crucial here. The proof is the same as in the proof of Proposition 1.11 and 1.12 in

[8]. All is about stability, with respect to *initial data* in [8], and with respect to *the blow-up point* here. For this reason, we just sketch the proof and refer to [8] for the proofs.

We know from (25) that $w_0(y, s) \rightarrow \kappa$ as $s \rightarrow \infty$ in L_ρ^2 . If a is a blow-up point near 0, we know from (15) that $w_a \rightarrow \pm\kappa$ as $s \rightarrow \infty$. Only $+\kappa$ is selected because of the following local constant sign property for u , which is a direct consequence of the positivity of u for $x = 0$, and the ODE comparison of Proposition 2.2.

Lemma A.1 (Constant sign property of u for (x, t) close to $(0, T)$)

There exists $\delta > 0$ such that

$$\forall t \in [T - \delta, T), \forall x \in B(0, \delta), u(x, t) \geq 0.$$

Proof : See corollary 1.8 in [8]. ■

The Liouville Theorem of Proposition 2.1 allows us to show that $v_a \equiv w_a - \kappa \rightarrow 0$ as $s \rightarrow \infty$ in L_ρ^2 , uniformly for a in a neighborhood of 0. This uniformity is a central argument in our proof.

Lemma A.2 (Uniform smallness of v_a) *There exists $\tilde{\sigma}_2 > 0$ such that*

$$i) \sup_{|\sigma| < \tilde{\sigma}_2} \|v_a(s)\|_{L_\rho^2} \rightarrow 0 \text{ as } s \rightarrow \infty,$$

$$ii) \text{ For all } R > 0, \sup_{|\sigma| < \tilde{\sigma}_2} \left(\sup_{|y| < R} |v_a(y, s)| \right) \rightarrow 0 \text{ as } s \rightarrow \infty.$$

Proof : See Proposition 1.10 in [8]. ■

Note that this Lemma yields the first estimate of Proposition 3.2. We know from (22) that for each a , either $v_a \sim v_{a,2}$ or $v_a \sim v_{a,-}$ as $s \rightarrow \infty$. (25) shows that $v_0 \sim v_{0,2}$, its neutral mode, as $s \rightarrow \infty$. The predominance of the neutral mode turns out to be a stable behavior for solutions of (17). Indeed :

Lemma A.3 (Uniform stability of the dynamics where v_2 is pre-dominant) *There exists $\tilde{\sigma}_4$ with the following property :*

i) There exists s^ such that for all $|\sigma| < \tilde{\sigma}_4$ and $s_0 \geq s^*$,*

$$\text{if } X_a(s_0) \geq Y_a(s_0) + Z_a(s_0), \text{ then } \forall s \geq s_0, X_a(s) \geq \frac{1}{2} (Y_a(s) + Z_a(s))$$

where

$$X_a(s) = \|v_{a,2}(s)\|_{L_\rho^2}, Y_a(s) = \|v_{a,-}(s)\|_{L_\rho^2} + \| |y|^{\frac{k}{2}} v_a \|_2^2, Z_a(s) = \|v_{a,+}(s)\|_{L_\rho^2}$$

and $k = k(M) > 0$ (where M is defined in (14)) is fixed.

ii) For all $\epsilon > 0$, there exists $s_3(\epsilon)$ such that for all $|\sigma| < \tilde{\sigma}_4$ and $s \geq s_3(\epsilon)$,

$$Y_a(s) + Z_a(s) \leq \epsilon X_a(s).$$

Proof: See Proposition 1.11 in [8]. ■

This lemma yields the second estimate of Proposition 3.2, namely the fact that $v_a \sim v_{a,2}$ as $s \rightarrow \infty$, uniformly in $a(\sigma)$. Therefore, the study of (17) reduces to the study of its projection over the neutral mode, $v_{a,2}$, where, of course, one should take into account the quadratic term. This leads to the equation stated in ii) of Proposition 3.2 (A_a is the matrix of the components of $v_{a,2}$, thanks to (21)). See Proposition 1.12 in [8] for a proof.

This closes the proof of Proposition 3.2. ■

B A refined equation on the neutral mode of (17)

We prove Lemma 3.8 here. We first prove (29). We claim the following :

Claim B.1 (Control of $v_a(s)$ in L_ρ^r) For all $r > 1$, there exists $C(r) > 0$ and $s_0^*(r) > 0$ such that for all $|\sigma| < \sigma_3$ and $s \geq s_3 + s_0^*(r)$,

$$\left(\int |v_a(y, s)|^r \rho(y) dy \right)^{\frac{1}{r}} \leq \frac{C(r)}{s}.$$

Proof :

If $r = 2$, we use Lemma 3.7 to get $|A_a(s)| \leq Cs^{-1}$, hence $\|v_{a,2}(s)\|_{L_\rho^2} \leq Cs^{-1}$ by (21). Since $\|v_a(s)\|_{L_\rho^2} \sim \|v_{a,2}(s)\|_{L_\rho^2}$ as $s \rightarrow \infty$ (uniformly for $|\sigma| < \sigma_3$), we get the estimate with some $s_0^*(2)$.

If $r \neq 2$, we combine the case $r = 2$ with the following regularizing effect of the operator \mathcal{L} .

Claim B.2 (Herrero-Velázquez) If v_a satisfies (17), then for all $r > 1$, there exists $\tilde{s}_0(r)$ and $C(r, M)$ such that

$$\left(\int |v_a(y, s)|^r \rho(y) dy \right)^{1/r} \leq C \left(\int |v_a(y, s - \tilde{s}_0)|^2 \rho(y) dy \right)^{1/2}.$$

Proof : See Lemma 2.3 in [14]. This closes the proof of Claim B.1. ■

(29) follows if we prove that for all $|\sigma| < \sigma_3$ and $s \geq s'_3$ for some $s'_3 \geq s_3$,

$$z'_a(s) \geq \frac{1}{2}z_a(s) - \frac{C}{s^2} \tag{90}$$

$$y'_a(s) \leq -\frac{1}{2}y_a(s) + \frac{C}{s^2} \tag{91}$$

where $z_a(s) = \|v_{a,+}(s)\|_{L^2_\rho}$ and $y_a(s) = \|v_{a,-}(s)\|_{L^2_\rho}$. Indeed, integrating (90) between s and $+\infty$ and (91) between s'_3 and s , we obtain for all $|\sigma| < \sigma_3$ and $s \geq s'_3$,

$$\begin{aligned} z_a(s) &\leq C e^{s/2} \int_s^\infty e^{-t/2} t^{-2} dt \leq C' s^{-2} \\ y_a(s) &\leq e^{-(s-s'_3)/2} y_a(s'_3) + C \int_{s'_3}^s e^{-(s-t)/2} t^{-2} dt < C' s^{-2} \end{aligned}$$

(Note that $y_a(s) + z_a(s) \leq \|v_a(s)\|_{L^2_\rho} \leq \|v_a(s)\|_{L^\infty} \leq M + \kappa$ by (14)). Now we prove (90). (91) follows in the same way.

Projecting (17) on the positive eigenspace of \mathcal{L} , we get

$$\partial_s v_{a,+} = \mathcal{L} v_{a,+} + P_+(f(v_a))$$

where P_+ is the corresponding L^2_ρ orthogonal projector. Multiplying this by $v_{a,+}\rho$ and integrating over \mathbb{R}^N , we get

$$\begin{aligned} \frac{1}{2} \frac{d}{ds} \int v_{a,+}^2 \rho &= \int \mathcal{L} v_{a,+} v_{a,+} \rho + \int P_+(f(v_a)) v_{a,+} \rho \\ &\geq \frac{1}{2} \int v_{a,+}^2 \rho - \left(\int v_{a,+}^2 \rho \right)^{1/2} \left(\int P_+(f(v_a))^2 \rho \right)^{1/2} \end{aligned}$$

where we used Cauchy-Schwartz's inequality.

Since P_+ is a L^2_ρ projector and f is quadratic, we have

$$\left(\int [P_+(f(v_a))]^2 \rho \right)^{1/2} \leq \left(\int (f(v_a))^2 \rho \right)^{1/2} \leq C \left(\int (v_a)^4 \rho \right)^{1/2}.$$

Using claim B.1, we have for all $|\sigma| < \sigma_3$ and $s \geq s_3 + s_0^*(4)$, $(\int v_a^4 \rho)^{1/2} \leq C(4)s^{-2}$. Thus, (90) follows with $s'_3 = s_3 + s_0^*(4)$. This closes the proof of (29).

Now we prove (30). The estimate on eigenvalues is obtained by evaluating the estimate on A_a at eigenfunctions. Therefore, we just focus on A_a . With (29) and Claim B.1, we are ready to get a refined equation on $A_a(s)$. Using (20) and (17), we write for all $|\sigma| < \sigma_3$ and $s \geq s_3$,

$$A'_a(s) = E_1 + E_2 + E_3 \tag{92}$$

where

$$\begin{aligned} E_1 &= \frac{p}{2\kappa} \int v_{a,2}(y, s)^2 M(y) \rho(y) dy, \\ E_2 &= \frac{p}{2\kappa} \int (v_a^2 - v_{a,2}^2) M(y) \rho(y) dy, \quad E_3 = \int g(v_a) M(y) \rho(y) dy. \end{aligned}$$

By straightforward calculations, we get from (21)

$$E_1 = \frac{1}{\beta} A_a(s)^2 \text{ where } \beta = \frac{\kappa}{2p}. \quad (93)$$

Recalling that $v = v_{a,+} + v_{a,2} + v_{a,-}$ (see (19)), we write

$$\begin{aligned} |E_2| &\leq C \int |v_a^2 - v_{a,2}^2| |M(y)| \rho(y) dy \\ &= C \int |v_{a,+} + v_{a,-}| |v_a + v_{a,2}| |M(y)| \rho(y) dy \\ &\leq C \left(\int |v_{a,+} + v_{a,-}|^2 \rho \right)^{1/2} \left(\int |v_a + v_{a,2}|^4 \rho \right)^{1/4} \left(\int |M(y)|^4 \rho \right)^{1/4} \end{aligned}$$

where we used Hölder's inequality.

Using (29), Claim B.1, and the norm equivalence in the finite dimensional space (18) where $v_{a,2}$ lays $(\int |v_{a,2}|^4 \rho \leq C (\int |v_{a,2}|^2 \rho)^2 \leq C (\int |v_a|^2 \rho)^2$ by Proposition 3.1), we end-up with

$$|E_2| \leq C s^{-3} \text{ for all } s \geq s_3'' \quad (94)$$

for some $s_3'' > s_3$.

We finally estimate E_3 . From (20), (17) and Cauchy-Schwartz inequality, we write $|E_3| \leq \int |g(v_a)| |M(y)| \rho(y) dy \leq C (\int |v_a|^6 \rho)^{1/2} (\int |M(y)|^2 \rho(y) dy) = C (\int |v_a|^6 \rho)^{1/2}$. Using Claim B.1, we end-up with

$$|E_3| \leq C s^{-3} \text{ for all } s \geq s_3 + s_0^*(6) \quad (95)$$

Combining (92), (93), (94) and (95), we obtain the first inequality in (30), for all $|\sigma| < \sigma_3$ and $s \geq \tilde{s}_3$ for some $\tilde{s}_3 \geq s_3$. This closes the proof of Lemma 3.8.

C Asymptotic behavior in the $\frac{y}{\sqrt{s}}$ variable

We prove iii) of Proposition 3.1 here. We follow Velázquez's work in [24] (Proposition 2.3), with a special care to the convergence speed and to the uniformity with respect to blow-up points.

Define

$$\varphi(y, s) = \left(p - 1 + \frac{(p-1)^2 y_1^2}{4p s} \right)^{-\frac{1}{p-1}} + \frac{\kappa}{2ps} \text{ and } q_a = w_a - \varphi.$$

Straightforward calculations based on (13) and i) of Proposition 3.1 yield for all $|\sigma| < \sigma_0$ and $s \geq s_0$,

$$\begin{cases} \forall y \in \mathbb{R}^N, \partial_s q_a = (\mathcal{L} + V)q_a + B(q_a) + R(y, s), \\ \|q_a(s)\|_{L_\rho^2} \leq C_0 \frac{\log s}{s^2} \end{cases} \quad (96)$$

where

$$\begin{aligned} \mathcal{L} &= \Delta - \frac{1}{2}y \cdot \nabla + 1, \quad V(y, s) = p\varphi^{p-1} - \frac{p}{p-1} \\ B(q) &= |\varphi + q|^{p-1}(\varphi + q) - \varphi^p - p\varphi^{p-1}q, \\ R(y, s) &= -\partial_s \varphi + \Delta \varphi - \frac{1}{2}y \cdot \nabla \varphi - \frac{\varphi}{p-1} + \varphi^p. \end{aligned} \quad (97)$$

Let us introduce the following norm

$$L_\rho^{2,r}(\psi) = \sup_{|\xi| < r} \left(\int |\psi(y)|^2 \rho(y - \xi) dy \right)^{1/2}.$$

Velázquez's idea in [24] is to make estimates on solutions of (96) in the $L_\rho^{2,r(\tau)}$ norm where $r(\tau) = K_0 e^{\frac{\tau-s_0}{2}} \leq K_0 \sqrt{\tau}$. The following is proved in [24] :

Lemma C.1 *For all $C_3 > 0$ and $K_3 > 0$, there is $s_3(C_3, K_3) > 0$ such that if $Z(y, s)$ satisfies*

$$\begin{cases} \partial_s Z \leq (\mathcal{L} + \frac{C_3}{s})Z + C_3 \left(Z^2 + \frac{1+|y|^2}{s^2} + 1_{\{|y| > 2K_3 \sqrt{s}\}} \right), \\ 0 \leq Z(y, s) \leq C_3, \end{cases} \quad (98)$$

for all $(y, s) \in \mathbb{R}^N \times [s_3, \infty)$, then, for all $s' \geq s_3$ and $\tau \in [s', s]$, we have

$$\begin{aligned} g(\tau) &\leq C_4(C_3) \left[e^{\tau-s'} \left(\|Z(s')\|_{L_\rho^2} + \frac{\log s'}{s'^2} \right) \right. \\ &\quad \left. + \int_{s'}^{(\tau-2K_3)_+} \frac{e^{\tau-t-2K_3} g(t)^2}{(1 - e^{-(\tau-t-2K_3)})^{1/20}} dt \right] \end{aligned}$$

where $g(\tau) = L_\rho^{2,r(K_3, \tau, s')}(Z(\tau))$, $r(K_3, \tau, s') = K_3 e^{\frac{\tau-s'}{2}}$, $e^{\frac{s-s'}{2}} = \sqrt{s}$ and $h_+ = \max(h, 0)$.

Proof : See Proposition 2.3 in [24] (in particular, pages 1575-1581). \blacksquare

Let us apply this to our case (96). It is readily seen from (97) that for all $K_0 > 0$, there is $C_2(K_0, M)$ (M is defined in (14)) such that

$$\begin{aligned} V(y, s) &\leq \frac{C_2}{s}, \quad |B(q)| \leq C_2 \left(|q|^2 + 1_{\{|y| \geq 2K_0\sqrt{s}\}} \right). \\ |R(y, s)| &\leq C_2 \left(\frac{1+|y|^2}{s^2} + 1_{\{|y| \geq 2K_0\sqrt{s}\}} \right) \end{aligned} \quad (99)$$

(one may consider first the case $|y| \geq 2K_0\sqrt{s}$ and then $|y| \leq 2K_0\sqrt{s}$ and make a Taylor expansion for $\xi = \frac{y}{\sqrt{s}}$ bounded).

If $Z_a = |q_a|$, then we use Kato's inequality $\Delta g \cdot \text{sgn}(g) \leq \Delta(|g|)$ to derive from (96) and (99) the following :

For all K_0 , there is $C_2(K_0, M) > 0$ such that for all $|\sigma| < \sigma_0$ and $s \geq s_0$,

$$\begin{cases} \forall y \in \mathbb{R}^N, \quad \partial_s Z_a \leq (\mathcal{L} + \frac{C_2}{s})Z_a + C_2 \left(Z_a^2 + \frac{1+|y|^2}{s^2} + 1_{\{|y| \geq 2K_0\sqrt{s}\}} \right), \\ \|Z_a(s)\|_{L_\rho^2} \leq C_0 \frac{\log s}{s^2}. \end{cases} \quad (100)$$

We claim the following :

Claim C.2 *For all $K_0 > 0$, there exists $C_6 > 0$ and s_6 such that for all $|\sigma| < \sigma_0$ and $s \geq s_6$,*

$$L_\rho^{2, K_0\sqrt{s}}(Z_a(s)) \leq C_6 \frac{\log s}{s}.$$

Proof : Consider $K_0 > 0$. Fix $s_5(K_0) \geq 2 \max(s_3(C_2(K_0, M), K_0), s_0)$ such that for all $s \geq s_5$, $\frac{s}{2} - \log s \geq 0$ where C_2 , s_3 and M are defined in (100),

(14) and Lemma C.1. If $s \geq s_5$ and s' is defined by $e^{\frac{s-s'}{2}} = \sqrt{s}$, then it is readily seen that $s' = s - \log s \geq \frac{s}{2} \geq \max(s_3, s_0)$. Therefore, we have from (96), (100) and lemma C.1 : for all $|\sigma| < \sigma_0$, for all $\tau \in [s', s]$,

$$g(\tau) \leq C'_5(C_0, K_0, M) \left[e^{\tau-s'} \frac{\log s'}{s'^2} + \int_{s'}^{(\tau-2K_0)_+} \frac{e^{\tau-t-2K_0} g(t)^2}{(1 - e^{-(\tau-t-2K_0)})^{1/20}} dt \right]$$

where $g(\tau) = L_\rho^{2, r(K_0, \tau, s')} (Z_a(\tau))$ and $r(K_0, \tau, s') = K_0 e^{\frac{\tau-s'}{2}}$.

By a standard Gronwall estimate (see Lemma 2.2 in [24]), there is $s_6(C_0, K_0) \geq s_5$ and $C_5(C_0, K_0) > C'_5$ such that if $s \geq s_6$, then for all $\tau \in [s', s]$, $g(\tau) \leq C_5 e^{\tau-s'} \frac{\log s'}{s'^2}$.

If $\tau = s$, then we get $L_\rho^{2, K_0\sqrt{s}}(Z_a(s)) \leq C_5 e^{s-s'} \frac{\log s'}{s'^2} \leq 2C_5 \frac{\log s}{s}$. This finishes the proof of Claim C.2. \blacksquare

To conclude the proof, we use the following result from [24] :

Claim C.3 Assume that Z satisfies (98), then

$$\sup_{|y| \leq \frac{K_3}{2} \sqrt{s}} |Z_a(y, s)| \leq C(C_3, K_3) L_p^{2, K_3 \sqrt{s}}(Z_a(s)).$$

Proof : See Proposition 2.3 in [24] (in particular, page 1581). ■

Thus, for all $|\sigma| < \sigma_0$ and $s \geq s_6$,

$$\sup_{|y| \leq \frac{K_0}{2} \sqrt{s}} |Z_a(y, s)| \leq C(K_0, C_0, M) \frac{\log s}{s}.$$

Since $|w_a(y, s) - f_1(\frac{y}{\sqrt{s}})| \leq Z_a + \frac{\kappa}{2ps}$, this concludes the proof of iii) of Proposition 3.1. ■

References

- [1] J. M. Ball. Remarks on blow-up and nonexistence theorems for nonlinear evolution equations. *Quart. J. Math. Oxford Ser. (2)*, 28(112):473–486, 1977.
- [2] A. J. Bernoff, A. L. Bertozzi, and T. P. Witelski. Axisymmetric surface diffusion: dynamics and stability of self-similar pinchoff. *J. Statist. Phys.*, 93(3-4):725–776, 1998.
- [3] M. D. Betterton and M. P. Brenner. Collapsing bacterial cylinders. *Phys. Rev. E*, 64(061904), 2001.
- [4] M. P. Brenner, P. Constantin, L. P. Kadanoff, A. Schenkel, and S. C. Venkataramani. Diffusion, attraction and collapse. *Nonlinearity*, 12(4):1071–1098, 1999.
- [5] J. Bricmont and A. Kupiainen. Universality in blow-up for nonlinear heat equations. *Nonlinearity*, 7(2):539–575, 1994.
- [6] S. J. Chapman, B. J. Hunton, and J. R. Ockendon. Vortices and boundaries. *Quart. Appl. Math.*, 56(3):507–519, 1998.
- [7] K. Deng and H. A. Levine. The role of critical exponents in blow-up theorems : the sequel. *J. Math. Anal. Appl.*, 2000.
- [8] C. Fermanian Kammerer, F. Merle, and H. Zaag. Stability of the blow-up profile of non-linear heat equations from the dynamical system point of view. *Math. Annalen*, 317(2):195–237, 2000.

- [9] C. Fermanian Kammerer and H. Zaag. Boundedness up to blow-up of the difference between two solutions to a semilinear heat equation. *Nonlinearity*, 13(4):1189–1216, 2000.
- [10] S. Filippas and R. V. Kohn. Refined asymptotics for the blowup of $u_t - \Delta u = u^p$. *Comm. Pure Appl. Math.*, 45(7):821–869, 1992.
- [11] S. Filippas and W. X. Liu. On the blowup of multidimensional semilinear heat equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 10(3):313–344, 1993.
- [12] H. Fujita. On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$. *J. Fac. Sci. Univ. Tokyo Sect. I*, 13:109–124, 1966.
- [13] Y. Giga and R. V. Kohn. Nondegeneracy of blowup for semilinear heat equations. *Comm. Pure Appl. Math.*, 42(6):845–884, 1989.
- [14] M. A. Herrero and J. J. L. Velázquez. Blow-up behaviour of one-dimensional semilinear parabolic equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 10(2):131–189, 1993.
- [15] T. Kato. *Perturbation theory for linear operators*. Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition.
- [16] H. A. Levine. Some nonexistence and instability theorems for solutions of formally parabolic equations of the form $Pu_t = -Au + F(u)$. *Arch. Rational Mech. Anal.*, 51:371–386, 1973.
- [17] F. Merle. Solution of a nonlinear heat equation with arbitrarily given blow-up points. *Comm. Pure Appl. Math.*, 45(3):263–300, 1992.
- [18] F. Merle and H. Zaag. Reconnection of vortex with the boundary and finite time quenching. *Nonlinearity*, 10(6):1497–1550, 1997.
- [19] F. Merle and H. Zaag. Stability of the blow-up profile for equations of the type $u_t = \Delta u + |u|^{p-1}u$. *Duke Math. J.*, 86(1):143–195, 1997.
- [20] F. Merle and H. Zaag. Optimal estimates for blowup rate and behavior for nonlinear heat equations. *Comm. Pure Appl. Math.*, 51(2):139–196, 1998.
- [21] F. Merle and H. Zaag. A Liouville theorem for vector-valued nonlinear heat equations and applications. *Math. Annalen*, 316(1):103–137, 2000.

- [22] F. Oustry and Overton M. L. Variational analysis of the total projection for symmetric matrices. 2000.
- [23] H. M. Soner and P. E. Souganidis. Singularities and uniqueness of cylindrically symmetric surfaces moving by mean curvature. *Comm. Partial Differential Equations*, 18(5-6):859–894, 1993.
- [24] J. J. L. Velázquez. Higher-dimensional blow up for semilinear parabolic equations. *Comm. Partial Differential Equations*, 17(9-10):1567–1596, 1992.
- [25] J. J. L. Velázquez. Classification of singularities for blowing up solutions in higher dimensions. *Trans. Amer. Math. Soc.*, 338(1):441–464, 1993.
- [26] J. J. L. Velázquez. Estimates on the $(n - 1)$ -dimensional Hausdorff measure of the blow-up set for a semilinear heat equation. *Indiana Univ. Math. J.*, 42(2):445–476, 1993.
- [27] H. Zaag. One dimensional behavior of singular N dimensional solutions of semilinear heat equations. *Comm. Math. Phys.*, 2002. to appear.

Address:

Courant Institute, NYU, 251 Mercer Street, NY 10012, New York.

Current address:

École Normale Supérieure, Département de mathématiques et applications,
45 rue d’Ulm, 75005 Paris.

e-mail : Hatem.Zaag@ens.fr