

Stable time-parallel integration of advection dominated problems using Parareal with space coarsening.


Julien Bodart – *ISAE-SUPAERO*
Serge Gratton – *Toulouse-INP-IRIT*
Xavier Vasseur – *ISAE-SUPAERO*
Thibaut Lunet – *University of Geneva*

May 4, 2018



Larger and larger problems for research and industrial applications with Computational Fluid Dynamics

- ▶ Higher complexity
→ Turbulence, Acoustics, Combustion ...
- ▶ High fidelity simulation
→ High Order Discretization, LES, DNS, ...



From right to left: RANS, LES, DNS

Massively parallel supercomputer for tomorrow

- ▶ Supercomputer speed rather based on **number of cores** than *processor speed*
- ▶ Largest one today:
 - ▶ $\sim 10 \times 10^6$ cores
 - ▶ ~ 100 PetaFlop/s
- ▶ Highlights the limits of exclusive space-parallelization



Sunway TaihuLight (2016) ©www.dailymail.co.uk

⇒ **Space-time parallelism could be an interesting alternative to enhance efficiency on exascale supercomputers**

Actual solutions for time-parallelization

- ▶ Space-Time Multigrid - The first born
- ▶ Parareal - The famous cadet
- ▶ PFASST - When complexity serves efficiency
- ▶ MGRIT - Toward an universal solution
- ▶ And many others ...

How to convince the HPC-CFD community ?

- ▶ Proof of concept on representative test-cases
 1. Accuracy of the time-parallel integration
 2. Efficiency gain compared to exclusive space-parallelization
- ▶ Solution that can be easily integrated into (huge) pre-existent CFD codes
 - ▶ Explicit time-stepping solvers
 - ▶ Temporal evolution of variables (e.g. pressure sensor for acoustics simulation)
 - ▶ ...

⇒ **First step : investigations of Parareal^{R1} with space coarsening^{R2}**

[R1] Lions et al., "A "Parareal" in time discretization of PDE's" (2001)

[R2] Fischer et al., "A Parareal in time semi-implicit approximation of the Navier-Stokes equations" (2005)

What was done so far

PhD Thesis - "Space-time parallel strategies for the numerical simulation of turbulent flows"
(Defended January 9, 2018)

- ▶ What could be the best solution from today's algorithms ? (Chap. 2)
- ▶ Can we understand theoretically the behavior of explicit forms of PARAREAL? (Chap. 3)
- ▶ What about large scale turbulent flow problems ? (Chap. 4)
 - ▶ Space-time parallel efficiency ?
 - ▶ Accuracy on two representative test case
(Homogeneous Isotropic Turbulence, Turbulent Channel Flow)

Part of the work was accepted for publication ^{R1}

But there was a major issue at the beginning ...

[R1] Lunet et al., "Time-parallel simulation of the decay of homogeneous turbulence using Parareal with spatial coarsening" (2017)

Parareal VS Advective Problems

Many studies underlined the difficulties of Parareal on

$$\frac{\partial U}{\partial t} + c \frac{\partial U}{\partial x} = 0$$

- ▶ Numerical instabilities^{R1} and slow convergence for some setting^{R2}
- ▶ PARAREAL loses its contraction factor on periodic domains (cf. M. Gander's talk)

Difficulty to prove with such problem if it would work on CFD problems

1. PARAREAL does not define a unique algorithm
2. Molecular viscosity and Reynolds number
 - ▶ *"The convergence of Parareal deteriorates as the viscosity parameter becomes smaller and the flow becomes more and more dominated by convection."*^{R3}
 - ▶ **But : the Reynolds number does not have a unique definition !**
Low influence of the Re_λ number increase compared to other parameters for Homogeneous Isotropic Turbulence (cf. PhD manuscript)
3. In most CFD problem, space resolution and Reynolds number increase simultaneously
4. Space coarsening implies to choose an interpolation method
(Linear, High Order, Fourier,)

[R1] Ruprecht and Krause, "Explicit parallel-in-time integration of a linear acoustic-advection system" (2012)

[R2] Gander, "Analysis of the Parareal algorithm applied to hyperbolic problems using characteristics" (2008)

[R3] Steiner et al., "Convergence of Parareal for the Navier-Stokes equations depending on the Reynolds number" (2015)

Main object of this talk

- ▶ Starts from the 1D linear advection problem with low diffusion
- ▶ Focus on one particular PARAREAL form
 1. Space coarsening for \mathcal{G} (one point out of two)
 2. High order explicit time-integration (RK4)
 3. Highly accurate space discretization (Centered 6th order)
- ▶ Change several parameters that can influence PARAREAL convergence (Reynolds, space resolution, interpolation method, ...)
- ▶ Increase problem complexity (non-linearity, ...)
- ▶ Try to answer the following questions :

What are the most influent parameters for this version of PARAREAL ?

How to set them to enhance convergence for a more complex case ?

Definition of a baseline test case


- ▶ Advection with small diffusion

$$\frac{\partial u}{\partial t} = -c \frac{\partial u}{\partial x} + \nu \frac{\partial^2 u}{\partial x^2}, \quad \nu \ll c$$

- ▶ Periodic 1D mesh with $x \in [0, 1]$
- ▶ Gaussian initial solution with varying width

$$u_0(x) = e^{-\frac{(x - 1/2)^2}{\sigma^2}}$$

- ▶ $CFL = 1$ for both fine and coarse solvers
- ▶ Final time $T = 64\delta_t$ ($\sim T_{period}/7$)
- ▶ Time domain decomposition in 4 time-slices



Error criterion based on fine solution comparison

$$E_{T,L_2}^k = \frac{\|U_{\mathcal{P}}^k(T) - U_{\mathcal{F}}(T)\|_2}{\|U_{\mathcal{F}}(T)\|_2}, \quad \text{for } k \in \{0, 1, 2, 3\}$$

Definition of a baseline test case


- ▶ Advection with small diffusion

$$\frac{\partial u}{\partial t} = -c \frac{\partial u}{\partial x} + \nu \frac{\partial^2 u}{\partial x^2}, \quad \nu \ll c$$

- ▶ Periodic 1D mesh with $x \in [0, 1]$
- ▶ Gaussian initial solution with varying width

$$u_0(x) = e^{-\frac{(x - 1/2)^2}{\sigma^2}}$$

- ▶ $CFL = 1$ for both fine and coarse solvers
- ▶ Final time $T = 64\delta_t$ ($\sim T_{period}/7$)
- ▶ Time domain decomposition in 4 time-slices



Error criterion based on fine solution comparison

$$E_{T, L_2}^k = \frac{\|U_{\mathcal{P}}^k(T) - U_{\mathcal{F}}(T)\|_2}{\|U_{\mathcal{F}}(T)\|_2}, \quad \text{for } k \in \{0, 1, 2, 3\}$$

What will vary in the next graphs

Main parameters


- ▶ Interpolation method
 1. Linear ($\sigma_I = 1$, blue-circle)
 2. Cubic ($\sigma_I = 3$, orange-square)
 3. 7th order ($\sigma_I = 7$, green-triangle)
- ▶ Space mesh resolution
 1. Fine (left side)
 2. Coarse (right side)

Secondary parameters (lines - dashes - dots)


1. Reynolds number
2. Time slice length
3. Regularity of the solution
4. Non-linearity of the advection term

Linear case - influence of the Reynolds number

$Re = c/\nu$: from low to high
2000 (line) \rightarrow 10000 (dashes) \rightarrow 20000 (dots)



High space resolution ($N_x = 500$)




Low space resolution ($N_x = 100$)

Main observations


- ▶ Staggered benefit of interpolation order increase (first on \mathcal{G} , then on Parareal convergence)
- ▶ Few influence of Re for the 1st iteration with low order interpolation or low space resolution

Linear case - influence of the time-slice length

Number of δ_t per time-slice: from large to small
 64 (line) \rightarrow 32 (dashes) \rightarrow 16 (dots)



High space resolution ($N_x = 500$)




Low space resolution ($N_x = 100$)

Main observations


- ▶ Small impact on the convergence
- ▶ Effect is "inverted" when going to high order interpolation

Linear case - influence of the solution regularity

Width of the initial Gaussian: from large to small
 $\sigma = 8$ (line) $\rightarrow \sigma = 16$ (dashes) $\rightarrow \sigma = 32$ (dots)



High space resolution ($N_x = 500$)



Low space resolution ($N_x = 100$)


Main observations

- ▶ Mainly influence the coarse solver error, less the convergence
- ▶ A too low space resolution cancels the beneficial impact of high order interpolation

The new problem

- ▶ Non-linear advection term


$$\frac{\partial u}{\partial t} = -u \frac{\partial u}{\partial x} + \nu \frac{\partial^2 u}{\partial x^2}, \quad \nu \ll \max_x(u_0)$$




- ▶ Centered scheme applied to $\frac{1}{2} \frac{\partial u^2}{\partial x}$

Non linear case - influence of the Reynolds number

$Re = \max(u_0)/\nu$: from low to high
2000 (line) \rightarrow 10000 (dashes) \rightarrow 20000 (dots)



High space resolution ($N_x = 500$)




Low space resolution ($N_x = 250$)

Main observations


- ▶ Similar behavior as the linear case, except for deterioration of the coarse solver accuracy
- ▶ Bad space resolution quickly cancels high order interpolation benefits

Non linear case - influence of the time-slice length

Number of δ_t per time-slice: from large to small
 128 (line) \rightarrow 64 (dashes) \rightarrow 32 (dots)



High space resolution ($N_x = 500$)



Low space resolution ($N_x = 250$)


Main observation

- Increasing the time-slice length enhances the convergence (for each resolutions)


\neq linear case

Non linear case - influence of the solution regularity

Width of the initial Gaussian: from large to small
 $\sigma = 8$ (line) $\rightarrow \sigma = 16$ (dashes) $\rightarrow \sigma = 32$ (dots)



High space resolution ($N_x = 500$)



Low space resolution ($N_x = 250$)

Main observation

- Increasing sharpness of the solution \simeq increasing the Reynolds number

Conclusions from this study

General conclusion for PARAREAL with space coarsening on advection problem

- ▶ Reasonably good convergence obtained for some cases
- ▶ Advection is not the only player to blame, there is also
 1. Low order interpolation (**PLEASE do not use linear interpolation !**)
 2. Space mesh resolution not adapted to a sharp initial solution
 3. ...
- ▶ Non-linearity can change everything
 1. Increasing the time-slice can enhance the convergence
 2. More sensitivity to the tuple: (mesh resolution, solution form)

Perspectives

- ▶ Numerical experiments done with the CASPER PYTHON code
 1. Not open-source yet but can be shared at demand
 2. Could be used to conduct many other tests
- ▶ Theoretical Fourier analysis of the algorithm to understand its main behavior (DD25 + draft)
- ▶ Complete convergence theory for the advection-diffusion problem (contraction factor, ...)

Thanks a lot for your attention,

I would be glad to answer if you have
Any questions ?