

DENJOY CONSTRUCTIONS FOR FIBRED HOMEOMORPHISMS OF THE TORUS

François Béguin, Sylvain Crovisier, Tobias Jäger and Frédéric Le Roux

July 27, 2007

Abstract

We construct different types of quasiperiodically forced circle homeomorphisms with transitive but non-minimal dynamics. Concerning the recent Poincaré-like classification for this class of maps of [1], we demonstrate that transitive but non-minimal behaviour can occur in each of the different cases. This closes one of the last gaps in the topological classification.

Actually, we are able to get some transitive quasiperiodically forced circle homeomorphisms with rather complicated minimal sets. For example, we show that, in some of the examples we construct, the unique minimal set is a Cantor set and its intersection with each vertical fibre is uncountable and nowhere dense (but may contain isolated points).

We also prove that minimal sets of the later kind cannot occur when the dynamics are given by the projective action of a quasiperiodic $SL(2, \mathbb{R})$ -cocycle. More precisely, we show that, for a quasiperiodic $SL(2, \mathbb{R})$ -cocycle, any minimal strict subset of the torus either is a union of finitely many continuous curves, or contains at most two points on generic fibres.

1 Introduction

We study quasiperiodically forced (qpf) circle homeomorphisms, that is homeomorphisms of the torus which are homotopic to the identity and have the form

$$f : \mathbb{T}^2 \rightarrow \mathbb{T}^2, \quad (\theta, x) \mapsto (\theta + \omega, f_\theta(x)), \quad (1.1)$$

with $\omega \in \mathbb{R} \setminus \mathbb{Q}$. The class of such maps will be denoted by \mathcal{F} .

Skew products like this occur in various situations in physics. One well-known example is the so-called Harper map, which appears in the study of quasi-crystals and the corresponding Schrödinger operators (see, for example, [2, 3]). Another one is the qpf Arnold circle map, which is used as a simple model for oscillators forced with two or more incommensurate frequencies [4].

The interest in transitive but non-minimal dynamics in this kind of maps is motivated by a recent classification result in [1], which we briefly want to discuss in order to motivate the problem.

Given any lift $F : \mathbb{T}^1 \times \mathbb{R} \rightarrow \mathbb{T}^1 \times \mathbb{R}$, $\theta \in \mathbb{T}^1$ and $n \in \mathbb{N}$, let

$$F_\theta^n := F_{\theta+(n-1)\omega} \circ \dots \circ F_\theta .$$

Then the limit

$$\rho(F) = \lim_{n \rightarrow \infty} (F_\theta^n(x) - x)/n \quad (1.2)$$

exists and does not depend on $(\theta, x) \in \mathbb{T}^1 \times \mathbb{R}$. Furthermore, the convergence in (1.2) is uniform [5]. $\rho(f) = \rho(F) \bmod 1$ is called the (*fibred*) *rotation number* of f . However, unlike the one-dimensional case, the *deviations from the average rotation*, given by

$$D_n(\theta, x) := F_\theta^n(x) - x - n\rho(F) , \quad (1.3)$$

need not be uniformly bounded in n, θ, x anymore.¹ This gives rise to a basic dichotomy: a homeomorphism $f \in \mathcal{F}$ is called ρ -*bounded* if $\sup_{n, \theta, x} |D_n(\theta, x)| < \infty$ and ρ -*unbounded* otherwise.

Another concept which turned out to be fundamental in this context are (p, q)-*invariant strips*. These are compact invariant sets which intersect each vertical fibre $\{\theta\} \times \mathbb{T}^1$ in exactly pq compact intervals, have an internal p -periodic structure and certain additional regularity properties. Since the precise formulation is somewhat technical and we will not use it later, we refrain from stating it here and refer to [1] or [7] for the definition. Among (p, q)-invariant strips are the *continuous (p, q)-invariant graphs*. These are the minimal invariant subsets of \mathbb{T}^2 such that the restriction of $p_1 : (\theta, x) \mapsto \theta$ is a pq -fold covering with p connected components. In order to have a rough idea, the reader should just think of an invariant strip as a generalisation of a continuous (p, q)-invariant graph, where each point of the graph is possibly replaced by a vertical segment. In particular, the existence of such an object forces the rotation number $\rho(f)$ to be rationally related to ω and the deviations (1.3) to be bounded [7].

It turns out that in the ρ -bounded case a direct analogue to the Poincaré Classification Theorem (e.g. [8]) holds, with invariant strips playing the role of periodic orbits in the unforced case:

Theorem 1.1 (theorems 3.1 and 4.1 in [1]).

- (a) If $f \in \mathcal{F}$ is ρ -bounded, then either there exists a (p, q)-invariant strip and $\rho(f)$, ω and 1 are rationally dependent or f is semi-conjugate to the irrational torus translation $(\theta, x) \mapsto (\theta + \omega, x + \rho(f))$ by a continuous semi-conjugacy h which is fibre-respecting (i.e. $p_1 \circ h = p_1$).
- (b) If $f \in \mathcal{F}$ is ρ -unbounded, then it is topologically transitive.

Since all known examples of ρ -unbounded behaviour are either minimal (all ρ -unbounded skew rotations are minimal, see [8, proposition 4.2.6]; other examples are given in [9]) or their topological dynamics have not yet been clarified, this immediately raises the question whether transitive but non-minimal dynamics can occur in the ρ -unbounded case. Similarly, it is not known whether

¹In the case of an unforced circle homeomorphism, the uniform bound is 1.

this is possible when f is semi-conjugate to an irrational rotation - which could be interpreted as “a Denjoy counter-example without wandering sets”.²

The positive answer to these questions is provided by the following theorem, which is the main result of this paper:

Theorem 1.2.

- *There exists a transitive non minimal qpf circle homeomorphism which is ρ -unbounded;*
- *There exists a transitive non minimal qpf circle homeomorphism which is semi-conjugate to an irrational rotation.*

This theorem will follow from a general construction.

Theorem 1.3. *Suppose R is a minimal qpf circle homeomorphism. Then there exist a continuous and surjective map $\pi : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ and a qpf circle homeomorphism f such that $\pi \circ f = R \circ \pi$ and f is topologically transitive, but not minimal. In addition, if R is a diffeomorphism,³ then f can be chosen such that all fibre maps f_θ are circle diffeomorphisms and $\partial_x f_\theta$ depends continuously on (θ, x) .*

This can be interpreted as follows. If R is a minimal qpf circle homeomorphism with certain additional properties, and if these properties are preserved by topological extension, then there exists $f \in \mathcal{F}$ with the same properties, but transitive and non-minimal dynamics. In particular, this is true for the properties ‘*semi-conjugated to an irrational rotation*’ and ‘*unbounded deviations*’. Another such property, related to the structure of the ergodic invariant measures, will be discussed in section 4.3 .

The proof of theorem 1.3 will be given via propositions 2.2 and 3.1 below and their addenda, which immediately imply the above statement. The construction we carry out is very similar to Denjoy’s construction of circle homeomorphisms with wandering sets. The main idea is to start with a continuous curve Γ and to ‘blow up’ this curve and all its images to small annuli, just as the points of an orbit are blown up to wandering intervals in Denjoy’s construction. However, instead of requiring that Γ is disjoint from all its images, which would lead to wandering sets, we choose the curve Γ such that there are ‘many’ intersections, and this fact is then used to establish the transitivity of f . Further, it turns out that in order to make the construction work, the initial curve must have another, rather surprising property: whenever it intersects any of its images this must happen over a whole interval - in other words the connected components of the intersection must not be singletons (see definition 2.1 and proposition 2.2). This property will turn out to be crucial in order to ensure the continuity of the semi-conjugacy π during the construction.

The construction of such a curve Γ is first carried out in the case where R is real-analytic, since this allows to avoid some technical problems and renders the main ideas more visible.

²In this context, we would also like to mention the constructions by Mary Rees in [13] and [14] (see also [15]). These equally produce Denjoy-like examples without wandering sets on the two-torus, but in the fibred case the dynamics will always remain minimal.

³In fact, it suffices that all fibre maps R_θ are circle diffeomorphisms and $\partial_x R_\theta$ depends continuously on (θ, x) .

It should be mentioned that there exist well-known examples of qpf circle homeomorphism with transitive but non-minimal dynamics, which are due to Shnirelman [6] (see also, for example, [8, section 12.6(b)]). However, in these examples there always exists an invariant curve, which is just a special case of an invariant strip. Consequently, the resulting minimal set (the invariant curve) has have a very simple structure. In contrast to this, it is known that minimal strict subsets of \mathbb{T}^2 in the absence of invariant strips must be much more complicated (see proposition 4.1 taken from [7, theorem 4.5 and lemma 4.6]). In particular, we obtain the following result (see subsection 4.2).

Proposition 1.4. *There exists a transitive non minimal qpf circle homeomorphism whose unique minimal set is a Cantor set, and whose intersection with each fibre $\{\theta\} \times \mathbb{T}^1$ is uncountable.*

Apart from the Denjoy-like constructions, we collect some general properties of minimal sets of qpf circle homeomorphism. In particular, we prove the following uniqueness result (see subsection 4.1 for more results).

Proposition 1.5. *Suppose $f \in \mathcal{F}$ has no invariant strip. Then it has a unique minimal set.*

Proposition 1.4 shows that general qpf circle homeomorphisms may possess quite complicated minimal sets. In the particular case of quasiperiodic $SL(2, \mathbb{R})$ -cocycles, which has received a lot of attention in the recent years (see, for example, [11] and references therein), we prove that such “complicated” minimal sets cannot occur. More precisely, we obtain the following.

Proposition 1.6 (see propositions 4.6, 4.7, 4.8 for more detailed statements). *Suppose f is given by the projective action of a quasiperiodic $SL(2, \mathbb{R})$ -cocycle. Then any minimal set of f*

1. *is the whole torus,*
2. *or is a continuous (p, q) -invariant graph,*
3. *or intersects generic fibres in one point,*
4. *or intersects generic fibres in two points.*

If f is in addition ρ -unbounded, then any minimal set either is the whole torus, or intersects generic fibres in only one point.

This proposition seems to improve some recent results of Bjerklöv and Johnson ([16]), by showing that one of the five possible cases of the classification obtained by these author never occurs.

Finally, we want to mention another result which had originally been a motivation for the presented work.

Theorem 1.7 ([7, theorem 4.4]). *Suppose $f \in \mathcal{F}$ is \mathcal{C}^2 and has no invariant strips. Then f is topologically transitive.*

Obvioulsy, this raises again the question whether transitive but non-minimal behaviour is possible in the absence of invariant strips. However, it must be said that our results are not directly related to theorem 1.7, since we do not obtain examples with this type of regularity. As stated, we are only able to

choose the fibre maps \mathcal{C}^1 , and by some slight modifications one might push this to $\mathcal{C}^{1+\alpha}$ (see section 3.2.3). (Of course, the fact that a Denjoy-like construction produces this type of regularity is by no means surprising.) Hence, the question whether the assertion of theorem 1.7 can be improved to minimality is still open. In fact, this is not even known under much stronger assumptions, for example if f is real-analytic and ω is Diophantine, or if f is induced by the projective action of a quasiperiodic $\mathrm{SL}(2, \mathbb{R})$ -cocycle.

2 Construction of the graph Γ

2.1 Graphs with flat intersections

Let R be a quasiperiodically forced circle homeomorphism over some circle irrational rotation $\theta \mapsto \theta + \omega$. We will consider the graphs Γ of continuous maps $\gamma : \mathbb{T}^1 \rightarrow \mathbb{T}^1$. For $I \subseteq \mathbb{T}^1$ we define $\Gamma|_I := \Gamma \cap (I \times \mathbb{T}^1)$. The \mathcal{C}^0 -distance between continuous maps induces a distance d between graphs. By $p_i : (\theta, x) \mapsto \theta$ we denote the canonical projection to the i -th coordinate.

Definition 2.1. *Suppose $\gamma, \gamma' : \mathbb{T}^1 \rightarrow \mathbb{T}^1$ are two continuous maps with graphs Γ, Γ' .*

- (a) *We say Γ and Γ' have flat intersections if $p_1(\Gamma \cap \Gamma')$ consists of a finite union of disjoint intervals, none of which is reduced to a single point.*
- (b) *We say Γ and Γ' cross over some interval $I \subset \mathbb{T}^1$ if there exists an interval $I' \subset I$ and an open interval $O \subsetneq \mathbb{T}^1$ such that $\Gamma|_{I'}, \Gamma'|_{I'} \subset I' \times O$ and $\Gamma'|_{I'}$ meets both connected components of $(I' \times O) \setminus \Gamma$.*

Proposition 2.2. *Let R be a quasiperiodically forced circle homeomorphism. Assume that R does not admit any continuous (p, q) -invariant graph.*

Then there exists a continuous graph Γ which has flat intersections with all its iterates $R^n(\Gamma)$ ($n \in \mathbb{Z}$).

In the situation of theorem 1.3, the non-existence of continuous (p, q) -invariant graphs follows immediately from the minimality of R . However, the fact that the above proposition holds under this weaker assumption will be useful in the later sections, and the proof is identical in both cases.

Addendum 2.3. *Assume that R is topologically transitive. Then the graph Γ may be required to satisfy the following property:*

- (T) *For all non-trivial intervals $I, J \subset \mathbb{T}^1$, there exists some $n \geq 0$ such that Γ and $R^n(\Gamma)$ cross over $I \cap (J + n\omega)$.*

The graph Γ required by proposition 2.2 will be the limit of a sequence (Γ_n) , obtained by induction using proposition 2.4 below. The section is organised as follows. We first state proposition 2.4 and show that it entails proposition 2.2. Sections 2.2, 2.3 and 2.4 are mostly devoted to the proof of proposition 2.4. The addendum is proved at the end of section 2.4.

We say a graph $\tilde{\Gamma}$ is an ε -modification of a graph Γ over $I \subseteq \mathbb{T}^1$ if $\tilde{\Gamma}|_{I^c} = \Gamma|_{I^c}$ and $d(\tilde{\Gamma}, \Gamma) < \varepsilon$, where d denotes the \mathcal{C}^0 -distance.

Proposition 2.4. *Let R be a quasiperiodically forced circle homeomorphism. Assume that R does not admit any continuous (p, q) -invariant graph. Let Γ be a continuous graph that has flat intersections with $R^k(\Gamma)$ for $1 \leq k \leq n-1$.*

Let $\varepsilon > 0$. Then there exists a continuous graph $\tilde{\Gamma}$ such that

1. $\tilde{\Gamma}$ is an ε -modification of Γ over a set of measure less than ε ;
2. for $1 \leq k \leq n-1$, the graph $\tilde{\Gamma}$ has flat intersections with $R^k(\tilde{\Gamma})$, moreover the set $p_1(\tilde{\Gamma} \cap R^k \tilde{\Gamma})$ contains $p_1(\Gamma \cap R^k \Gamma)$ and has the same number of connected components;
3. the graph $\tilde{\Gamma}$ has flat intersections with $R^n(\tilde{\Gamma})$.

Proof of proposition 2.2 using proposition 2.4. We begin the induction by choosing Γ_1 to be any continuous graph. Let $n \geq 1$, and assume inductively that there exists a continuous graph Γ_{n-1} which has flat intersection with its iterates $R^k(\Gamma_{n-1})$ for all k such that $1 \leq k \leq n-1$. We apply proposition 2.4, to get a continuous graph $\Gamma_n = \tilde{\Gamma}_{n-1}$ which has flat intersection with its iterates $R^k(\Gamma_n)$ for all k such that $1 \leq k \leq n$. Furthermore, we can demand that the graph Γ_n is a ε_n -modification of the graph Γ_{n-1} , with $\varepsilon_n \leq \frac{1}{2^n}$ (the choice of ε_n will be made more precise below).

Thus we get a sequence $(\Gamma_n)_{n \geq 1}$ of continuous graphs, which is a Cauchy sequence for the \mathcal{C}^0 -distance. Let $\tilde{\Gamma}$ be the limit map. Let k be a fixed positive integer. The sequence $(p_1(R^k(\Gamma_n) \cap \Gamma_n))_{n \geq 1}$ is an increasing sequence of subsets of \mathbb{T}^1 , denote its limit by I^k ,

$$I^k = \text{Clos} \left(\bigcup_{n \geq 1} p_1(R^k(\Gamma_n) \cap \Gamma_n) \right).$$

Note that according to property 2 of proposition 2.4, every set in this sequence has the same number a_k of connected components, so that I^k is again the disjoint union of at most a_k non trivial compact intervals.

This set I^k is included in $p_1(R^k(\Gamma) \cap \Gamma)$. To get the reverse inclusion we have to make a more careful choice of the sequence (ε_n) . For a fixed $n \geq k$, denote by $\mathcal{O}_{n,k}$ the set of continuous graphs Δ such that $p_1(R^k(\Delta) \cap \Delta)$ is included in the $\frac{1}{n}$ -neighbourhood of $p_1(R^k(\Gamma_n) \cap \Gamma_n)$ (which we denote by $V_{\frac{1}{n}}(p_1(R^k(\Gamma_n) \cap \Gamma_n))$). This set is open for the \mathcal{C}^0 distance. Thus we may have chosen the sequence $(\varepsilon_n)_{n \in \mathbb{N}}$ so small that for every $n \geq k$, $\Gamma \in \mathcal{O}_{n,k}$. This entails, for every $k \leq n$,

$$p_1(R^k(\Gamma_n) \cap \Gamma_n) \subset p_1(R^k(\Gamma) \cap \Gamma) \subset V_{\frac{1}{n}}(p_1(R^k(\Gamma_n) \cap \Gamma_n)).$$

Letting n tends towards infinity (with fixed k) gives the required equality $I^k = p_1(R^k(\Gamma) \cap \Gamma)$. Thus we get that Γ has flat intersection with $R^k(\Gamma)$. \square

2.2 Perturbation boxes

In this section, we introduce the tools required by the proof of proposition 2.4. We consider a continuous graph Γ and some positive integer n and make the following *escaping hypothesis*.

Every point has an iterate outside $\Gamma \cup R(\Gamma) \cup \dots \cup R^n(\Gamma)$.

In the case where R is minimal this is obviously true, in lemma 2.11 below we show that it also holds under the weaker hypothesis of proposition 2.2 .

Returning dynamics on Γ

We introduce the *first return map* T on Γ in time less or equal to n : let D be the set points z in Γ such that there exists some iterate $R^q(z)$ in Γ with $1 \leq q \leq n$; we then define the map T on D by $T(z) = R^q(z)$ where q is the least such integer.

Due to the escaping hypothesis, the orbit of any point of Γ by T is finite. This allows the following definitions.

Notations and definitions. To any point $z \in \Gamma$ is associated a unique finite set, $N(z) = \{q_{-r} < \dots < q_0 = 0 < \dots < q_s\} \subset \mathbb{Z}$, called the *itinerary* of z , such that

- for any integer k such that $q_{-r} - n \leq k \leq q_s + n$, the point $R^k(z)$ belongs to Γ if and only if $k = q_i$ for some $-r \leq i \leq s$,
- $|q_i - q_{i+1}| \leq n$ for each $-r \leq i < s$.

We thus have $T^i(z) = R^{q_i}(z)$, and the sequence $(R^{q_i}(z))_{i=-r, \dots, s}$ is the T -orbit of z . We will denote by $\ell(z) = s + r$ the length of this orbit. When we wish to emphasize the dependence on the point z , we will write $r(z)$, $s(z)$, and so on. Let us note that the T -orbit of z is reduced to $\{z\}$ if and only if all the iterates $f^k(z)$ are outside Γ for $0 < |k| \leq n$.

More generally, for any sufficiently small interval $I \subset \mathbb{T}^1$, there exists a finite set $\{q_i, i = -r, \dots, s\}$, again called the *itinerary* of I , such that $0 < q_{i+1} - q_i \leq n$ and for all k such that $q_{-r} - n \leq k \leq q_s + n$ we have

$$R^k(\Gamma|_I) \cap \Gamma \neq \emptyset \Leftrightarrow k \in \{q_i, i = -r, \dots, s\}.$$

This follows from compactness together with the openness with respect to z of the property $R^k(z) \notin \Gamma$ for a given k .

Definition and existence of perturbation boxes

Definition 2.5. A rectangle $B = I \times J$ is a perturbation box if

1. I has a finite itinerary $\{q_i, i = -r, \dots, s\}$ and has pairwise disjoint iterates $I + k\omega$ with $q_{-r} - n \leq k \leq q_s + n$;
2. one of the two endpoints of $\Gamma|_I$ has the same itinerary as I ;
3. for all k such that $q_{-r} - n \leq k \leq q_s + n$, the graph $(R^{-k}(\Gamma))|_I$
 - is contained in B if $k = q_i$ for some $-r \leq i \leq s$,
 - is disjoint from B otherwise.

We define the itinerary of the box to be the itinerary of I . Next we prove that every points of Γ belong to a perturbation box.

Lemma 2.6. If $z = (\theta, x) \in \Gamma$, then there exists δ, η arbitrarily small such that $B = [\theta, \theta + \delta] \times [x - \eta, x + \eta]$ is a perturbation box whose itinerary coincides with the itinerary of z .

Proof of the lemma. For δ and η small enough, the rectangle B is disjoint from the graphs $R^{-k}(\Gamma)$ with $q_{-r} - n \leq k \leq q_s + n$ such that k does not belong to the itinerary of z . For $k = q_i$, the graph $R^{-k}(\Gamma)$ contains the point z ; hence, if the interval δ is chosen after η and small enough, then $(R^{-k}(\Gamma))|_{[\theta, \theta + \delta]}$ is contained in B . \square

Perturbation lemma

If one assumes that the intersections $\Gamma \cap R^k(\Gamma)$ are controled for any k up to $n - 1$, the perturbation boxes can be used to build a perturbed graph Γ' whose intersections $\Gamma' \cap R^k(\Gamma')$ are flat for k up to n . These perturbations are given by the following lemma.

Lemma 2.7 (Perturbation lemma). *Let $B = I \times J$ be a perturbation box. Denote by $\{q_i, i = -r, \dots, s\}$ the itinerary of I . Then there exists a perturbation Γ' of Γ such that*

1. *the perturbation is supported in $\bigcup_{i=-r}^s R^{q_i}(B)$, and in particular the graphs $(R^{-q_i}(\Gamma'))|_I$ are still included in B .*
2. *Let $1 \leq k \leq n$ and define the sets $X_k = p_1(\Gamma \cap R^k \Gamma)$ and $X'_k = p_1(\Gamma' \cap R^k \Gamma')$. Then X'_k is the union of X_k and a finite number of intervals meeting X_k . More precisely, for every i such that X_k meets $I + q_i \omega$, there exists a non-trivial interval $J_{i,k}$ satisfying $X_k \cap (I + q_i \omega) \subset J_{i,k} \subset I + q_i \omega$, and we have the equality*

$$X'_k = X_k \cup \bigcup_{X_k \cap (I + q_i \omega) \neq \emptyset} J_{i,k}.$$

Furthermore, if R is real-analytic and Γ is piecewise real-analytic, then Γ' can be chosen piecewise real-analytic.

Proof of the perturbation lemma 2.7. We want to construct the new graph Γ' by modifying the graph Γ above each interval $I + q_i \omega$. Since all the intervals $I + q_i \omega$ are pairwise disjoint (item 1 of the definition of perturbation box), this amounts to modifying each graph $R^{-q_i}(\Gamma)$ above I .

We denote by z the endpoint of $\Gamma|_I$ that has the same itinerary as I (item 2 of the definition of perturbation box). To fix ideas, we assume that z is the left endpoint (the proof is entirely similar if z is the right endpoint). The definition of the itinerary entails that z also belongs to $R^{-q_i}(\Gamma)$ for all i . Denote $I = [\theta, \theta + \delta]$. Choose some $\lambda \in (0, \delta)$ with the property that for any given $i, j \in \{-r, \dots, s\}$ the graphs $R^{-q_i}(\Gamma)|_{[\theta + \lambda, \theta + \delta]}$ and $R^{-q_j}(\Gamma)|_{[\theta + \lambda, \theta + \delta]}$ either coincide at $\theta + \delta$ or are disjoint. Above I , we replace each graph $R^{-q_i}(\Gamma)$ by the union of two straight segments:

- above $[\theta, \theta + \lambda]$, the new graph is a horizontal segment (starting at z) ;
- above $[\theta + \lambda, \theta + \delta]$, the new segment is forced by continuity (it connects the right endpoint of the first segment to the point of $R^{-q_i}(\Gamma)$ above $\theta + \delta$).

Let us check the new graph Γ' has the announced properties. The first one concerning the support is a consequence of item 3 of the definition of a perturbation box. Let us turn to the second one. For any i, j between $-r$ and s , let us define the set

$$I_{i,j} = p_1(R^{-q_i}(\Gamma') \cap R^{-q_j}(\Gamma')) \cap I.$$

This set is either equal to $[\theta, \theta + \lambda]$ or to I . In case $p_1(R^{-q_i}(\Gamma) \cap R^{-q_j}(\Gamma)) \cap I = I$ then we still have $I_{i,j} = I$. By the choice of λ , in the opposite cases, $[\theta, \theta + \lambda]$

contains $p_1(R^{-q_i}(\Gamma) \cap R^{-q_j}(\Gamma)) \cap I$, and then so does $I_{i,j}$. Note that in any case we have the following property:

For all $i, j \in \{-r, \dots, s\}$, the interval $I_{i,j}$ is a non-trivial interval containing $p_1(R^{-q_i}(\Gamma) \cap R^{-q_j}(\Gamma)) \cap I$. (*)

Claim 2.8. *For every $k = 1, \dots, n$ and every $i = -r, \dots, s$,*

$$X_k \cap (I + q_i\omega) \neq \emptyset \Leftrightarrow \exists j, k = q_i - q_j.$$

When these equivalent properties hold we define $J_{i,k} = I_{i,j} + q_i\omega$. Note that according to property (*), $J_{i,k}$ contains $X_k \cap (I + q_i\omega)$. Obviously $J_{i,k}$ is contained in $I + q_i\omega$.

Proof of the claim.

$$\begin{aligned} \exists j, k = q_i - q_j &\Leftrightarrow R^{k-q_i}(\Gamma) \cap B \neq \emptyset && \text{(item 3 of definition 2.5)} \\ &\Leftrightarrow z \in R^{k-q_i}(\Gamma) && \text{(choice of } z) \\ &\Leftrightarrow R^{k-q_i}(\Gamma) \cap (R^{-q_i}(\Gamma))|_I \neq \emptyset \\ &\Leftrightarrow X_k \cap (I + q_i\omega) \neq \emptyset && \text{(apply } R^{q_i}). \end{aligned}$$

□

Now for getting property 2 it only remains to checking the following equality:

$$X'_k = X_k \cup \bigcup_{q_i - q_j = k} (I_{i,j} + q_i\omega).$$

Let us define the set $J = \bigcup_{i=-r}^s I + q_i\omega$. In order to check the above equality we partition \mathbb{T}^1 into the four sets

$$J \cap (J + k\omega), \quad J \setminus (J + k\omega), \quad (J + k\omega) \setminus J, \quad \mathbb{T}^1 \setminus (J \cup (J + k\omega)).$$

Let us examine the first set $J \cap (J + k\omega)$. According to item 1 of the definition of the perturbation boxes, we have

$$J \cap (J + k\omega) = \bigcup_{q_i - q_j = k} (I + q_i\omega).$$

Let i, j be such that $q_i - q_j = k$. Restricted to $I + q_i\omega$, we have $X'_k = I_{i,j} + q_i\omega$, and, according to property (*), this set contains the restriction of X_k : in other words,

$$X'_k \cap (I + q_i\omega) = [X_k \cup (I_{i,j} + q_i\omega)] \cap (I + q_i\omega).$$

It remains now to check that outside $J \cap (J + k\omega)$, the sets X'_k and X_k coincides. The second set $J \setminus (J + k\omega)$ is the union of the intervals $I + q_i\omega$ for those i such that $q_j \neq q_i - k$ for every j . Choose such an i . According to item 3 of the definition of perturbation boxes,

$$\Gamma|_{I+q_i\omega} \subset R^{q_i}(B) \text{ and } (R^k(\Gamma))|_{I+q_i\omega} \cap R^{q_i}(B) = \emptyset,$$

and in particular $X_k \cap (I + q_i\omega)$ is empty. According to item 1 of the lemma, the same relations hold when Γ is replaced by Γ' , thus $X'_k \cap (I + q_i\omega)$ is also empty. Thus X_k and X'_k coincide in restriction to $J \setminus (J + k\omega)$.

Restricting to the third set $(J + k\omega) \setminus J$, we prove symmetrically that both sets X_k and X'_k are empty, and so they also coincide.

On the last set $\mathbb{T}^1 \setminus (J \cup (J + k\omega))$, according to item 1 of the lemma, we have $\Gamma' = \Gamma$ and $R^k(\Gamma') = R^k(\Gamma)$ and thus $X'_k = X_k$. □

Remark 2.9. *In the proof of the perturbation lemma 2.7, we chose to replace the graph Γ by the simplest possible curve, i. e. the concatenation of two segments. But of course we could have used more complicated curves, for example the concatenation of a finite number of segments. Thus, if we are given some point z within the interior of the perturbation box B , this modification allows us to force the perturbed curve Γ' to contain the point z . The same holds for any finite number of points in $\text{Int}(B)$ (obviously having distinct first coordinate).*

2.3 Construction of the graph Γ : the real-analytic minimal case

The construction of the graph Γ is easier if R is a minimal rotation, or more generally if R is minimal and real-analytic. To explain this easy case, we state a simpler version of proposition 2.4.

Proposition 2.10. *Suppose that the assumptions of proposition 2.4 hold and in addition:*

- R is real-analytic and minimal,
- Γ is piecewise analytic.

Then there exists a curve $\tilde{\Gamma}$ which satisfies the assertions proposition 2.4 and is piecewise analytic.

Replacing proposition 2.4 by the preceding one (which is much easier to show), the proof of proposition 2.2 given at the beginning of section 2 can be easily adapted to prove the statement in the case when R is minimal and real-analytic.

Proof of proposition 2.10. Using that R is real-analytic and Γ is piecewise real-analytic, we know that the intersection $\Gamma \cap R^n\Gamma$ has finitely many connected components: these are isolated points and non-trivial curves. We will explain how to build a modification Γ' that is piecewise real-analytic, satisfies items 1 and 2 of the proposition 2.4 and moreover

3-bis. the number of connected components of $\Gamma' \cap R^n\Gamma'$ that are reduced to a point is strictly less than the corresponding number for $\Gamma \cap R^n\Gamma$.

By repeating this construction finitely many times, one obtains a modification $\tilde{\Gamma}$ of Γ that now satisfies the item 3 of the proposition.

Since R is supposed to be minimal, the escaping property is satisfied, and we can apply section 2.2. Let us consider an isolated point $z \in \Gamma \cap R^n(\Gamma)$. By lemma 2.6 there exists an arbitrarily small perturbation box B containing z in its boundary and having the same itinerary as z . We now apply the perturbation lemma 2.7. Since B is arbitrarily small, the perturbation Γ' is small: item 1 of proposition 2.4 is satisfied. One can also assume that the width $p_1(B)$ is smaller than half of the distance between any two connected

components of $\Gamma \cap R^k(\Gamma)$ for any $1 \leq k \leq n$. Item 2 of lemma 2.7 implies that $\Gamma' \cap R^k(\Gamma')$ contains $\Gamma \cap R^k(\Gamma)$ and has the same number of connected components. When $k < n$, one gets item 2 of the proposition. One also obtains item 3-bis by noting that the component $\{z\}$ of $\Gamma \cap R^n(\Gamma)$ has been replaced by a non-trivial interval. This completes the construction of Γ in the real-analytic case. \square

2.4 Construction of the graph Γ : the general case

In this section we prove proposition 2.4 and the addendum 2.3 to proposition 2.2 .

Escaping hypothesis

We start by checking the escaping hypothesis under which points have finite itineraries (see section 2.2).

Lemma 2.11. *Assume R does not admits any continuous (p, q) -invariant graph. Then for every continuous graph Γ and every $n > 0$, the escaping hypothesis is satisfied.*

Proof. If the escaping hypothesis is not satisfied, then there exists a invariant compact set K included in the union \hat{K} of a finite number of iterates of some graph Γ . Let us choose a minimal such K . Then the compact set K is a continuous (p, q) invariant graph contradicting the assumption. Indeed, since K is compact and invariant, $p_1(K)$ is the whole circle ; thus there is some k such that the projection of $R^k(\Gamma) \cap K$ has non-empty interior (Baire theorem), so that K contains some graph α over some interval. By taking a smaller open interval, we find a graph over an open interval which is an open set of K . By minimality, K is a one-dimensional topological manifold, thus a union of simple curves. The same argument shows that p_1 is a local homeomorphism on K , and thus a covering map (by compactness). \square

Proof of proposition 2.4

The proposition is obtained by applying inductively the following lemma. This lemma roughly says that if $R^n(\Gamma)$ has flat intersections with Γ outside some closed subset F , then we can construct $\tilde{\Gamma}$ such that $R^n(\tilde{\Gamma})$ has flat intersections with $\tilde{\Gamma}$ outside a closed subset F' , where F' is substantially smaller than F . Remember that the map T and the function ℓ have been defined on section 2.2; we will denote by T' and ℓ' the corresponding objects with respect to the graph Γ' .

Lemma 2.12. *Let Γ be a continuous graph that has flat intersections with $R^k(\Gamma)$ for $1 \leq k \leq n - 1$. Let F be a non-empty closed set which is a union of T -orbits. Suppose that $(\Gamma \cap R^n(\Gamma)) \setminus F$ has a finite number of connected components, none of which is a single point.*

Let $\varepsilon_0 > 0$. Then there exists an ε_0 -perturbation Γ' of Γ supported on an arbitrarily small neighbourhood of F , and there exists a non-empty closed set $F' \subset F$ which is a union of T -orbits such that

1. $(\Gamma' \cap R^n(\Gamma')) \setminus F'$ has a finite number of connected components, none of which is a single point.
2. For $1 \leq k \leq n - 1$, the graph Γ' has flat intersections with $R^k(\Gamma')$, moreover the set $p_1(\Gamma' \cap R^k \Gamma')$ contains $p_1(\Gamma \cap R^k \Gamma)$ and has the same number of connected components.
3. Either the set F' is empty, or the supremum of the function ℓ' on F' is strictly less than the supremum of ℓ on F .

Proof of the lemma. We assume the hypotheses of the lemma. If $\ell(z) = 0$ for every $z \in F$, then $R^n(\Gamma) \cap F = \emptyset$, and the graph $\Gamma' = \Gamma$ together with its closed subset $F' = \emptyset$ satisfies the conclusion of the lemma. From now on we assume that $\sup_F \ell > 0$.

Let

$$M = \{z \in F, \quad \ell(z) = \sup_F \ell\}.$$

The set M is obviously a non-empty union of T -orbits.

Claim 2.13. *The set M is closed.*

To prove the claim we consider the map $z \mapsto N(z)$ which associates to each point of Γ its itinerary (see section 2.2). Let $z \in \Gamma$; by definition of the itinerary, the points $f^k(z)$ are outside Γ for every $k \in \{q_{-r(z)} - n, \dots, q_{s(z)} + n\} \setminus N(z)$. We note that these conditions are open, so we have $N(z') \subset N(z)$ for all z' in a neighbourhood of z (in other words, the map $z \rightarrow N(z)$ is semi-continuous). The claim follows.

We consider the finite partition \mathcal{P} of M induced by N : two points y and z are in the same element P of \mathcal{P} if and only if $N(y) = N(z)$. Since the functions N and ℓ are constant on P we use the notation $N(P)$ and $\ell(P)$. The semi-continuity of N and the maximality of ℓ on M also entail that the elements of this partition are closed sets. Let \mathcal{P}_0 be the family of $P \in \mathcal{P}$ whose points z satisfy $r(z) = 0$. Then the partition \mathcal{P} can be written as

$$\mathcal{P} = \{T^k(P), P \in \mathcal{P}_0 \text{ and } 0 \leq k \leq \ell(P)\} = \{R^q(P), P \in \mathcal{P}_0 \text{ and } q \in N(P)\}.$$

Claim 2.14. *Let $P \in \mathcal{P}_0$. There exists a finite collection $\mathcal{B}(P)$ of arbitrarily small perturbation boxes such that*

1. each box B has the same itinerary $N(B)$ as the points of P ,
2. the projections $p_1(B)$ of the boxes have pairwise disjoint interior,
3. $p_1(P)$ is contained in the interior of the union of all the $p_1(B)$.

Proof of the claim. Let z be any point of P . By lemma 2.6, we can find two arbitrarily small perturbation boxes $B^-(z) = I^-(z) \times J^-(z)$, $B^+(z) = I^+(z) \times J^+(z)$ whose itineraries coincide with the itinerary of z , and such that $p_1(z)$ is the right end-point of $I^-(z)$ and the left end-point of $I^+(z)$. Since $p(P)$ is compact it is covered by the interior of finite number of intervals $I^-(z) \cup I^+(z)$. Thus we find a finite collection $\mathcal{B}^*(P)$ of perturbation boxes having properties 1 and 3 of the claim but maybe not property 2. Then property 2 will be achieved by replacing some of the intervals $I^\pm(z)$ by smaller subintervals. For this we first make the following remark. Let $z = (\theta_0, x) \in P$, and $\theta_1 \in \text{Int}(I^-(z)) = (\theta_0 - \delta, \theta_0)$. Then

- $[\theta_1, \theta_0] \times J^-(z)$ is a perturbation box,
- if $\theta_1 \in p_1(P)$ then $[\theta_0 - \delta, \theta_1] \times J^-(z)$ is a perturbation box,

and in both cases the new box has the same itinerary as the old one. Now consider any couple of perturbation boxes $B_1, B_2 \in \mathcal{B}^*(P)$ whose interiors are not disjoint. If the p_1 -projection of one box contains the other one then we can just eliminate the smallest one. In the opposite case, the above remark allows us to replace one of the two boxes, say B_1 , by a smaller perturbation box B'_1 whose projection by p_1 is disjoint from B_2 and such that the p_1 -projection of $B'_1 \cup B_2$ equals the p_1 -projection of $B_1 \cup B_2$. Thus, by considering one by one all the couples of boxes in $\mathcal{B}^*(P)$, we can construct a new collection $\mathcal{B}(P)$ having the wanted properties. \square

Let $\mathcal{B} = \cup \mathcal{B}(P)$ be the family of all the constructed boxes. Since the partition \mathcal{P} of M consists of disjoint closed sets, each $P \in \mathcal{P}_0$ is contained in an open set $U(P)$ such that the collection

$$\{R^q(U(P)), P \in \mathcal{P}_0 \text{ and } q \in N(P)\}$$

still consists of pairwise disjoint sets. Thus items 2 and 3 of the claim allows us to choose the families $\mathcal{B}(P)$ of perturbation boxes such that the elements of the following family have disjoint interior:

$$\{p_1(B) + q\omega, B \in \mathcal{B} \text{ and } q \in N(B)\}.$$

Thus we can apply the perturbation lemma 2.7 independently on each box of the family \mathcal{B} , and denote by Γ' the resulting graph. The graph Γ hence has been modified only in the domain

$$Z = \bigcup_{\substack{B \in \mathcal{B} \\ q \in N(B)}} p_1(R^q(B)) \times \mathbb{T}^1.$$

We define the set F' by $F' = F \setminus \text{Int}(Z)$ and we now check the properties.

The set F' is clearly a closed subset of F . For any $B \in \mathcal{B}$, and any $z \in B$, the itinerary of z is a subset of the itinerary of B . Thus the full T -orbit of z is included in the union $\bigcup_{q \in N(B)} R^q(B)$. Also note that the maps T and T' coincide outside $\text{Int}(Z)$, and thus on F' . Consequently, F' is a union of non-trivial T' -orbits.

Item 2 is obvious (cf property 2 of the perturbation lemma 2.7).

Let us check item 1. We first analyze the set $(\Gamma' \cap R^n(\Gamma')) \setminus F'$ in restriction to the complementary set of Z . We have

$$((\Gamma' \cap R^n(\Gamma')) \setminus F') \setminus Z = ((\Gamma \cap R^n(\Gamma)) \setminus F) \setminus Z.$$

The right-hand set is the restriction of $(\Gamma \cap R^n(\Gamma)) \setminus F$ (the union of finitely many non-trivial closed graphs) above the union of finitely many open intervals; thus it is the union of a finite number of non trivial intervals. Then we analyze the set $(\Gamma' \cap R^n(\Gamma')) \setminus F'$ in restriction to Z . According to lemma 2.7, $\Gamma' \cap R^n(\Gamma') \cap Z$ has a finite number of connected components, all of them non-trivial. Furthermore, by definition of F' , the set $F' \cap Z$ is included in the boundary of Z and is finite. Thus $((\Gamma' \cap R^n(\Gamma')) \setminus F') \cap Z$ is again the union of a finite number of non trivial intervals. Putting everything together, we get item 1.

Finally for item 3 we note that $M \subset \text{Int}(Z)$ and thus F' does not meet M . Since the maps T and T' coincide on F' , and according to the definition of the set M , we have

$$\sup_{F'} \ell' = \sup_{F'} \ell < \sup_F \ell .$$

This completes the proof. \square

Proof of proposition 2.4 from lemma 2.12. Let Γ and ε be as in the hypotheses of the proposition. We fix once and for all the value $\varepsilon_0 = \frac{\varepsilon}{\sup_{\Gamma} \ell + 1}$. We set $\Gamma_0 = \Gamma$ and $F_0 = \Gamma$. Then Γ_0 and F_0 satisfy the hypotheses of the lemma. Applying the lemma provides a new graph $\Gamma_1 = \Gamma'$ with a closed subset $F_1 = F'$. If F' is empty, then we define $\tilde{\Gamma} = \Gamma_1$ and note that $\tilde{\Gamma}$ satisfies the conclusion of proposition 2.4.

In the opposite case, Γ_1 and F_1 again satisfy the hypotheses of the lemma. Applying the lemma recursively provides sequences (Γ_p) and (F_p) . Since the inequality $\sup_{F_{p+1}} \ell_{p+1} < \sup_{F_p} \ell_p$ holds, there exists some p with $F_p = \emptyset$, and then we can define the graph $\tilde{\Gamma} = \Gamma_p$. Note that $p\varepsilon_0 < \varepsilon$, so that points 1 and 2 of the proposition concerning the size of the perturbation hold. This completes the proof of the proposition. \square

Proof of addendum 2.3

Addendum 2.3 requires the following additional property for the graph Γ : for any small pieces $\Gamma|_I, \Gamma|_J$, there exists some positive iterate of the first one that crosses the second one. In order to get this additional property, we will refine the construction of the sequence (Γ_n) , by inserting between two successive steps of the construction another small modification. The modification between steps n and $n+1$ will achieve the wanted property concerning two specific intervals I_n, J_n , while keeping the previously obtained properties of the graph Γ_n . We will get the modification by applying the following proposition.

Proposition 2.15. *Assume that R is topologically transitive. Let Γ be a continuous graph that has flat intersections with $R^k(\Gamma)$ for $1 \leq k \leq n-1$.*

Let $\varepsilon > 0$, and $I, J \subset \mathbb{T}^1$ be two non-trivial intervals. Then there exists a Graph $\tilde{\Gamma}$ such that

1. $\tilde{\Gamma}$ is an ε modification of Γ over a set of measure less than ε ;
2. there exists some integer $m > 0$ such that Γ and $R^m(\Gamma)$ crosses over $I \cap (J + m\omega)$;
3. for $1 \leq k \leq n-1$, the graph $\tilde{\Gamma}$ has flat intersections with $R^k(\tilde{\Gamma})$, moreover the set $p_1(\tilde{\Gamma} \cap R^k \tilde{\Gamma})$ contains $p_1(\Gamma \cap R^k \Gamma)$ and has the same number of connected components.

Proof of proposition 2.15. We first note that, up to replacing I and J by subintervals, we can assume the following additional properties:

- I and J have finite itineraries \mathcal{I} and \mathcal{J} ;
- there exists vertical intervals $I', J' \subset \mathbb{T}^1$ such that the rectangles $B_I = I \times I'$ and $B_J = J \times J'$ are perturbation boxes;
- any two intervals $I + k\omega, k \in \mathcal{I}$ and $J + \ell\omega, \ell \in \mathcal{J}$ are disjoint.

Since R is topologically transitive, there exists a point $z \in \text{Int}(B)$ with a dense forward orbit. Now we apply the perturbation lemma 2.7 with the perturbation box B_I to construct a first ε -modification $\bar{\Gamma}$ of Γ that contains z (this is possible thanks to remark 2.9). Note that the perturbation is supported on the iterates of B_I corresponding to the itinerary of B_I , and thus $\bar{\Gamma} \cap B_J = \Gamma \cap B_J$.

By hypothesis on z there is an iterate $R^m(z)$ with positive m belonging to $\text{Int}(B_J)$. Let z_1, z_2 be two points of $\text{Int}(B_J \cap ((I+m\omega) \times \mathbb{T}^1))$ that are separated in $B_J \cap ((I+m\omega) \times \mathbb{T}^1)$ by $R^m(\bar{\Gamma})$. Now we perform a second modification of Γ , this time using the box B_J , to construct a new graph $\tilde{\Gamma}$ that contains both points z_1, z_2 . Since $\tilde{\Gamma} \cap B_I = \bar{\Gamma} \cap B_I$, it follows that $R^m(\tilde{\Gamma})_{(I+m\omega)} = R^m(\bar{\Gamma})_{(I+m\omega)}$: thus z_1 and z_2 are still separated in $B_J \cap ((I+m\omega) \times \mathbb{T}^1)$ by $R^m(\tilde{\Gamma})$, that is, $\tilde{\Gamma}$ and $R^m(\tilde{\Gamma})$ cross over $(I+m\omega) \cap J$. \square

Proof of addendum 2.3 using proposition 2.15. We explain how to modify the inductive proof of proposition 2.4 given at the beginning of section 2 so that the addendum is satisfied.

Let (I_i) be a countable basis for the topology of \mathbb{T}^1 which consists of intervals. Let $(i_n, j_n)_{n \geq 1}$ be an enumeration of \mathbb{N}^2 , so that $(I_{i_n} \times I_{j_n})_{n \geq 1}$ is a basis for $\mathbb{T}^1 \times \mathbb{T}^1$.

Let $n \geq 1$, and assume inductively that there exists a continuous graph Γ_{n-1} which has flat intersections with its iterates $R(\Gamma_{n-1}), \dots, R^{n-1}(\Gamma_{n-1})$. We first mimic the construction of proposition 2.4 to get a continuous graph which we denote by Γ'_n ; in particular, Γ'_n is an ε_n -modification of Γ_{n-1} and has flat intersection with its n first iterates.

Thus we can apply proposition 2.15 to this graph Γ'_n , with intervals $I = I_{i_n}$ and $J = I_{j_n}$. We get again a continuous graph Γ_n which is an ε'_n -modification of Γ'_n (property 1). This new graph still has flat intersections with its n first iterates (property 3). Furthermore, it satisfies the following additional property (property 2): there exists some integer $m > 0$ such that Γ_n and $R^m(\Gamma_n)$ crosses over the interval $I_{i_n} \cap (I_{j_n} + m\omega)$.

We note that this last property is an open property among continuous graphs for the C^0 distance. Thus, by choosing the sequences (ε_n) and (ε'_n) to decrease sufficiently fast, we can ensure that the limit graph Γ also satisfies this property: for each integer $n \geq 1$ there exists an integer $m > 0$ such that Γ and $R^m(\Gamma)$ crosses over the interval $I_{i_n} \cap (I_{j_n} + m\omega)$. In particular, Γ meets the property required by the addendum. Furthermore, the argument of proposition 2.4 showing the flat intersection of Γ with $R^n(\Gamma)$ for all n is still valid. This completes the proof of the addendum. \square

3 Blowing up the orbit of Γ

In this section, we consider a quasiperiodically forced circle homeomorphism R and a continuous graph Γ which has flat intersections with all its iterates $R^n(\Gamma)$ ($n \in \mathbb{Z}$). We denote the fibres of p_1 by $\mathbb{T}_\theta^1 = \{\theta\} \times \mathbb{T}^1$.

Let us first recall that every probability measure μ on \mathbb{T}^2 can be disintegrated with respect to fibres of the projection $p_1 : (\theta, x) \mapsto \theta$. More precisely, one can find a family $(\mu_\theta)_{\theta \in \mathbb{T}^1}$ of probability measures on the circle \mathbb{T}^1 such that, for

every measurable set A ,

$$\mu(A) = \int_{\mathbb{T}^1} \mu_\theta(A_\theta) d\theta . \quad (3.1)$$

Here we use the notation

$$A_\theta = \{x \in \mathbb{T}^1 \mid (\theta, x) \in A\} ,$$

where A is any subset of \mathbb{T}^2 . The purpose of this section is to prove the following.

Proposition 3.1. *Let R be a qpf circle homeomorphism and suppose there exists a curve Γ which satisfies the assertions of proposition 2.2. Define*

$$\Xi := \bigcup_{n \in \mathbb{Z}} R^n(\Gamma) . \quad (3.2)$$

Then there exists a continuous onto map π and a homeomorphism f ,

$$\begin{array}{ccc} \pi : \mathbb{T}^2 & \longrightarrow & \mathbb{T}^2 \\ (\theta, x) & \longmapsto & (\theta, \pi_\theta(x)) \end{array} \quad \text{and} \quad \begin{array}{ccc} f : \mathbb{T}^2 & \longrightarrow & \mathbb{T}^2 \\ (\theta, x) & \longmapsto & (\theta + \omega, f_\theta(x)) \end{array}$$

with the following properties:

- (i) *for all $\theta \in \mathbb{T}^1$ the map π_θ is increasing;*
- (ii) *if $(\theta, x) \in \Xi$ then $\pi^{-1}(\theta, x)$ is a non-trivial interval in the circle \mathbb{T}_θ^1 ;*
- (iii) *if $(\theta, x) \notin \Xi$ then $\pi^{-1}(\theta, x)$ is a single point;*
- (iv) $\pi \circ f = R \circ \pi$.
- (v) $\pi^{-1}(\Xi)$ has non-empty interior.

Addendum 3.2. *Suppose R is transitive, Γ has the additional property provided by addendum 2.3 and in addition the set Ξ is dense in \mathbb{T}^2 . Then f is topologically transitive.*

Note that if R is minimal, then the fact that Ξ is dense is obvious. If R is only transitive, one may construct Γ with this property, see remark 3.10 . We remark that the proof of the addendum is short and does not depend on the proof of the proposition (see section 3.4).

Theorem 1.3 now follows immediately from propositions 3.1 and 2.2 and their addenda. Note that the non-minimality of f follows from property (v) in the proposition, since this implies that $\overline{\pi^{-1}(\Xi)^c}$ is a compact invariant strict subset of \mathbb{T}^2 .

Idea of the proof of proposition 3.1 Remember that we see proposition 3.1 as a generalisation of the classical Denjoy example on the circle. Here is one way to construct the Denjoy example. First choose an orbit O for an irrational rotation R on \mathbb{T}^1 , and let μ be a probability measure which has an atom at each point of O and no other atom. There is an (essentially unique) increasing map $\pi : \mathbb{T}^1 \rightarrow \mathbb{T}^1$ which sends the Lebesgue measure onto the measure μ . Then one looks for a circle homeomorphism f such that $\pi \circ f = R \circ \pi$: this equality determines f outside $\pi^{-1}(O)$; then one completes the construction by

choosing one way to extend f from the interval $\pi^{-1}(x)$ to $\pi^{-1}(R(x))$ for all $x \in O$.

We will adapt this construction to our setting, with the following modifications. The role of the orbit O is now played by the orbit of the curve Γ . The new measure μ is essentially a sum of one-dimensional measures along the iterates of Γ (μ_θ has an atom at x if and only if (θ, x) belongs to some iterate of Γ). As a consequence of the flat intersection hypothesis, there exists as before a map π sending the Lebesgue measure of \mathbb{T}^2 to μ . Then we construct a “nice” measure ν on \mathbb{T}^2 which satisfies $\pi_*\nu = R_*\mu$. This is the difficult part of the proof, the difficulty being linked to the fact that ν is not uniquely determined by this equality. Then f is defined as the (essentially unique) map sending the Lebesgue measure onto ν . The equality $\pi \circ f = R \circ \pi$ will follow automatically. The construction is summed up by the following commutative diagram.

$$\begin{array}{ccc} \mathbb{T}^2, \text{Leb} & \xrightarrow{f} & \mathbb{T}^2, \nu \\ \downarrow \pi & & \downarrow \pi \\ \mathbb{T}^2, \mu & \xrightarrow{R} & \mathbb{T}^2, R_*\mu \end{array}$$

3.1 The semi-conjugacy π

In this section, we consider any sequence $(\Gamma_n)_{n \in \mathbb{Z}}$ of curves in \mathbb{T}^2 such that, for every $i, j \in \mathbb{Z}$, the curves Γ_i and Γ_j have flat intersections. We denote $\Xi := \bigcup_{n \in \mathbb{Z}} \Gamma_n$.

3.1.1 Construction of the measure μ

For any graph Γ , we denote by δ_Γ the probability measure on \mathbb{T}^2 whose conditional $\delta_{\Gamma, \theta}$ is the Dirac mass at the point x such that $(\theta, x) \in \Gamma$. We choose a sequence of non-negative real numbers $(a_n)_{n \in \mathbb{Z}}$ such that $\beta := 1 - \sum_{n \in \mathbb{Z}} a_n$ is positive. Then we define a measure μ as follows:

$$\mu := \beta \text{Leb} + \sum_{n \in \mathbb{Z}} a_n \delta_{\Gamma_n}. \quad (3.3)$$

3.1.2 Definition of π

The flat intersections hypothesis plays a crucial role in the addendum of the following proposition.

Proposition 3.3. *For any probability measure μ as defined by (3.3), there exists a continuous onto map $\pi: \mathbb{T}^2 \rightarrow \mathbb{T}^2$ of the form $(\theta, x) \mapsto (\theta, \pi_\theta(x))$, such that, for every θ , the map $\pi_\theta: \mathbb{T}^1 \rightarrow \mathbb{T}^1$ is increasing and maps the Lebesgue measure onto μ_θ .*

In particular, π satisfies the properties (i), (ii) and (iii) stated in proposition 3.1.

Addendum 3.4. *For any given $n_0 \in \mathbb{Z}$, it is possible to choose π in proposition 3.3 such that the set $B_{n_0} = \pi^{-1}(\Gamma_{n_0})$ contains the annulus $S^1 \times [0, a_{n_0}]$.*

Proof of proposition 3.3 and addendum 3.4. Fix an integer $n_0 \in \mathbb{Z}$. Let $\Phi : (\theta, x) \mapsto (\theta, x + \alpha(\theta))$ be the skew rotation sending the graph Γ_{n_0} to the zero section $\mathbb{T}^1 \times \{0\}$. If we construct a map π satisfying the statement of the proposition and its addendum with the graphs Γ_n replaced by $\Phi(\Gamma_n)$ and the measure μ replaced by $\Phi_*\mu$, then the map $\pi' = \Phi^{-1} \circ \pi$ will satisfy the statements for the original objects. Thus we may change coordinates under Φ and assume that Γ_{n_0} is the zero section $\mathbb{T}^1 \times \{0\}$. In order to define the map $\pi : \mathbb{T}^2 \rightarrow \mathbb{T}^2$, we will first construct a map $\hat{\pi} : \mathbb{T}^1 \times [0, 1] \rightarrow \mathbb{T}^1 \times [0, 1]$. The following lemma is easy but crucial.

Lemma 3.5. *Denote by $P : \mathbb{T}^1 \times [0, 1] \rightarrow \mathbb{T}^2$ the natural projection. Then there exists a continuous lift $\hat{\mu}$ of μ , that is, $\hat{\mu}$ is a probability measure on $\mathbb{T}^1 \times [0, 1]$ such that $P_*\hat{\mu} = \mu$, and such that $\hat{\mu}$ depends continuously on θ (with respect to the weak topology on the space of probability measures).*

Proof of lemma 3.5. Clearly, the Lebesgue measure on $\mathbb{T}^1 \times [0, 1]$ is a continuous lift of the Lebesgue measure on \mathbb{T}^2 . Seemingly, since Γ_{n_0} is the null section, the measure $\delta_{\mathbb{T}^1 \times \{0\}}$ in $\mathbb{T}^1 \times [0, 1]$ is a lift of $\delta_{\Gamma_{n_0}}$ in \mathbb{T}^2 . It remains to prove that any measure $m = \delta_\Gamma$ on a graph Γ having a flat intersection with the null section Γ_{n_0} has a continuous lift \hat{m} . Now let θ be such that the points of Γ and Γ_{n_0} on \mathbb{T}_θ^1 are distinct. Then we choose \hat{m}_θ to be the only measure that projects down to m_θ , that is, the Dirac mass on $P^{-1}(\Gamma_\theta)$. By continuity this determines the value of $\hat{m}_\theta \in \{\delta_0, \delta_1\}$ when θ is an endpoint of a (non trivial) interval I where Γ coincides with the null section. Then we extend the construction on such an interval I by continuously (e. g. linearly) interpolating the Dirac masses δ_0 and δ_1 . \square

Next, we define the map $\hat{\pi} : \mathbb{T}^1 \times [0, 1] \rightarrow \mathbb{T}^1 \times [0, 1]$ by

$$\hat{\pi}(\theta, x) := (\theta, \hat{\pi}_\theta(x))$$

and

$$\hat{\pi}_\theta(x) = \min\{y \in [0, 1] \mid \hat{\mu}_\theta([0, y]) \geq x\}. \quad (3.4)$$

Observe that the map $\hat{\pi}$ is continuous; indeed:

- the map $\hat{\pi}_\theta : [0, 1] \rightarrow [0, 1]$ is continuous for every $\theta \in \mathbb{T}^1$ since the measure $\hat{\mu}_\theta$ gives a positive mass to every open set in $\{\theta\} \times [0, 1]$ (recall that $\beta > 0$);
- the map $\hat{\pi}_\theta$ depends continuously on θ since the measure $\hat{\mu}_\theta$ depends continuously on θ .

Moreover, by construction, for every θ the map $\hat{\pi}_\theta$ is increasing and maps the Lebesgue measure of $[0, 1]$ onto $\hat{\mu}_\theta$.

Clearly, the map $\hat{\pi} : \mathbb{T}^1 \times [0, 1] \rightarrow \mathbb{T}^1 \times [0, 1]$ induces a continuous map $\pi : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ having the wanted properties. Note that this map also satisfies the property required by addendum 3.4. \square

3.1.3 Uniqueness of π

Proposition 3.6. *Let us consider a measure μ as defined in (3.3) on \mathbb{T}^2 and two maps π, π' which both satisfy the assertions of proposition 3.3. Then, there*

exists a continuous skew rotation $A : (\theta, x) \mapsto (\theta, x + \alpha(\theta))$ of \mathbb{T}^2 such that $\pi = \pi' \circ A$.

Proof. For every $\theta \in \mathbb{T}^1$, both π_θ and π'_θ are circle maps which map the Lebesgue measure of \mathbb{T}_θ^1 onto μ_θ . It follows that, for every $\theta \in \mathbb{T}^1$, the circle maps π_θ and π'_θ coincide modulo a rotation: for every $\theta \in \mathbb{T}^1$, there exists $\alpha(\theta) \in \mathbb{T}^1$ such that $\pi_\theta(y) = \pi'_\theta(y + \alpha(\theta))$. This provides a skew rotation $A : (\theta, x) \mapsto (\theta, x + \alpha(\theta))$ of \mathbb{T}^2 such that $\pi = \pi' \circ A$. We are left to show that A is continuous, i.e. that $\alpha(\theta)$ depends continuously on θ .

Fix $\theta_0 \in \mathbb{T}^1$, and $\epsilon > 0$. Then choose $x_0 \in \mathbb{T}_\theta^1$ such that $\mu_{\theta_0}(\{x_0\}) = 0$. For every θ , we consider the intervals $I_\theta := \pi_\theta^{-1}(\{x_0\})$ and $I'_\theta := (\pi'_\theta)^{-1}(\{x_0\})$. Since $\mu_{\theta_0}(\{x_0\}) = 0$, both intervals I_{θ_0} and I'_{θ_0} consist of a single point. Moreover, the continuity of π implies that there exists $\delta > 0$ such that, for $\theta \in [\theta_0 - \delta, \theta_0 + \delta]$, the interval I_θ (resp. I'_θ) is included in an ϵ -neighbourhood of the interval I_{θ_0} (resp. I'_{θ_0}). Now, for every θ , the rotation $y \mapsto y + \alpha(\theta)$ maps the interval I_θ to the interval I'_θ . Consequently α is continuous at θ_0 , and as θ_0 was arbitrary it follows that α is continuous on the whole of \mathbb{T}^1 . \square

3.1.4 Description of the sets $\pi^{-1}(\Gamma_n)$

In order to construct the measure ν in the next section, we will need some more details about the geometry of the sets $B_n := \pi^{-1}(\Gamma_n)$. Note that the following proposition will only be used in the construction of ν , and it may be a good idea to skip the proof for a first reading.

Proposition 3.7. *There exists a sequence of open sets (U_n) , such that for all $n \in \mathbb{Z}$ there holds:*

- (i) $U_n \subseteq \text{int}(B_n)$;
- (ii) $U_n \cap U_m = \emptyset \ \forall m \neq n$;
- (iii) $\text{Leb}(U_{n,\theta}) = a_n$;
- (iv) for every θ and every n , $U_{n,\theta}$ is the union of at most $2|n|+1$ open intervals.

Remark 3.8.

- (a) Item (i) of this proposition implies property (v) of proposition 3.1.
- (b) For every θ , the union over n of the sets $U_{n,\theta}$ is contained in $(\text{Int}(\pi^{-1}(\Xi)))_\theta$. Property (iii) implies that this union has full measure in $(\pi^{-1}(\Xi))_\theta$. Therefore, for any $(\theta, x) \in \Xi$, the intersection $(\text{Int}(\pi^{-1}(\Xi)))_\theta \cap \pi_\theta^{-1}(\{x\})$ has full measure in $\pi_\theta^{-1}(\{x\})$, and therefore is dense in $\pi_\theta^{-1}(\{x\})$. This observation will be used in the construction of specific examples in section 4.

Addendum 3.9. *For any $\epsilon > 0$ and any $n \in \mathbb{Z}$, there exists a compact set $V_n \subset U_n$ that satisfies $\text{Leb}(V_{n,\theta}) \geq (1 - \epsilon)a_n \ \forall \theta \in \mathbb{T}^1$.*

Proof of proposition 3.7. Let us first give an idea of the proof, by explaining the construction of U_0 and U_1 . According to addendum 3.4, we may assume that $\pi^{-1}(\Gamma_0)$ contains the annulus $A_0 := \mathbb{T}^1 \times [0, a_0]$. We define U_0 to be the interior of this annulus. To construct U_1 , we first note that the map π can be factorised as $\pi = \pi_0 \circ P_0$ where P_0 consists in collapsing the annulus U_0 . Furthermore, the map π_0 is very similar to the map π : the results concerning π will also apply to π_0 , and in particular we will find an annulus A_1 included in

$\pi_0^{-1}(\Gamma_1)$ having the desired width. Taking the complement of the curve $P_0(U_0)$ in the interior of A_1 , and bringing this open set back under P_0 , will provide the open set U_1 ; note that in each fibre U_1 consists of at most two intervals, as required.

By relabelling the sequence Γ_n , we can assume that it is indexed over \mathbb{N} instead of \mathbb{Z} , which is more suitable for a proof by induction. Formally, every n in the following proof should read $\varphi(n)$ where φ is some bijection between \mathbb{N} and \mathbb{Z} . Note in particular that this changes property (iv), that now reads:

(iv-bis) for every θ and every n , $U_{n,\theta}$ is the union of at most $n+1$ open intervals.

The construction will be done by induction on $n \in \mathbb{N}$ by assuming the following additional hypotheses.

- (v) 1. There exists a fibre respecting monotonic map $P_n : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ having the following property: the restriction of the Lebesgue measure to the complement of $U_{0,\theta} \cup \dots \cup U_{n,\theta}$ is sent to the measure $\alpha_n \text{Leb}$ where $\alpha_n = 1 - a_0 - \dots - a_n$.
2. The projections of the open sets U_0, \dots, U_n by P_n are $n+1$ continuous graphs.
3. There exists a fibre respecting monotonic map $\pi_n : \mathbb{T}^2 \rightarrow \mathbb{T}^2$, such that for every θ , the map $\pi_{n,\theta} : \mathbb{T}^1 \rightarrow \mathbb{T}^1$ sends the Lebesgue measure on the fibre measure $\mu'(n)_\theta$ of the measure

$$\mu'(n) = \frac{1}{\alpha_n} \left(\beta \text{Leb} + \sum_{i>n} a_i \mu^i \right).$$

4. We have $\pi_n \circ P_n = \pi$.

Note that the crucial hypothesis here is the fact that the simultaneous collapsing of the sets U_0, \dots, U_n yields $n+1$ continuous graphs (hypothesis (v).2 above). Hypothesis (v) is illustrated by the following commutative diagram, showing the correspondence between the different measures.

$$\begin{array}{ccc}
 \text{Leb} & & \\
 \downarrow \pi & \searrow P_n & \\
 & & \alpha_n \text{Leb} + \sum_{i=0}^n a_i \delta_{P_n(U_i)} \\
 & \swarrow \pi_n & \\
 \mu = \alpha_n \mu'(n) + \sum_{i=0}^n a_i \delta_{\Gamma_i} & &
 \end{array}$$

Assume we have constructed the sets U_0, \dots, U_n and the maps P_n, π_n satisfying the above hypotheses.

Let $q : \mathbb{T}^2 \mapsto \mathbb{T}^2$ be the fibre respecting increasing map that sends the annulus $\bar{A} := \mathbb{T}^1 \times [0, \frac{a_{n+1}}{\alpha_n}]$ on the null section $\mathbb{T}^1 \times \{0\}$, and sends the restriction of the Lebesgue measure to the complement of $[0, \frac{a_{n+1}}{\alpha_n}]$ on the measure $\left(1 - \frac{a_{n+1}}{\alpha_n}\right) \text{Leb}$ (see the diagram below). Let $\pi_{n+1} : \mathbb{T}^2 \mapsto \mathbb{T}^2$ be a fibre respecting increasing map that sends the nul section on the curve Γ_{n+1} , and sends the Lebesgue measure in each fibre to the measure $\mu'(n+1) =$

$\alpha_{n+1}^{-1} (\beta \text{Leb} + \sum_{i>n+1} a_i \mu^i)$, where $\alpha_{n+1} = 1 - a_0 - \dots - a_{n+1}$. The existence of such a map π_{n+1} is guaranteed by proposition 3.3 and its addendum, applied with μ replaced by $\mu'(n+1)$, and choosing $n_0 = n+1$. In particular, this implies that property (v).3 is satisfied for $n+1$.

$$\begin{array}{ccc}
 \text{Leb} & \xrightarrow{\quad \Omega \quad} & \text{Leb} = \text{Leb}_{\bar{A}^c} + \text{Leb}_{\bar{A}} \\
 \downarrow \pi_n & & \downarrow q \\
 & & \frac{\alpha_{n+1}}{\alpha_n} \text{Leb} + \frac{a_{n+1}}{\alpha_n} \delta_{\mathbb{T}^1 \times \{0\}} \\
 & \swarrow \pi_{n+1} & \\
 \mu'(n) = \frac{\alpha_{n+1}}{\alpha_n} \mu'(n+1) + \frac{a_{n+1}}{\alpha_n} \delta_{\Gamma_{n+1}} & &
 \end{array}$$

Now the map $\pi_{n+1} \circ q$ is fibre-respecting and increasing, and it is easy to see that it sends the Lebesgue measure on the measure $\mu'(n)$. The map π_n shares the same properties. According to proposition 3.6, there exists a fibred rotation Ω such that $\pi_n = \pi_{n+1} \circ q \circ \Omega$. We take $P_{n+1} = q \circ \Omega \circ P_n$. Note that $\pi_{n+1} \circ P_{n+1} = \pi_n \circ P_n = \pi$, so that property (v).4 is satisfied for $n+1$. We have the following commutative diagram.

$$\begin{array}{ccccc}
 \mathbb{T}^2 & & & & \\
 \downarrow \pi & \searrow P_n & & & \downarrow P_{n+1} \\
 & \mathbb{T}^2 & \xrightarrow{\Omega} & \mathbb{T}^2 & \\
 & \downarrow \pi_n & & \downarrow q & \\
 & & & \mathbb{T}^2 & \\
 & \swarrow \pi_{n+1} & & & \\
 \mathbb{T}^2 & & & &
 \end{array}$$

Let A be the preimage by Ω of the annulus $\bar{A} = \mathbb{T}^1 \times [0, \frac{a_{n+1}}{\alpha_n}]$. Since Ω is a fibred rotation, by definition of \bar{A} , there exists a continuous map $\sigma : \mathbb{T}^1 \rightarrow \mathbb{T}^1$ such that

$$A = \left\{ (\theta, x), \sigma(\theta) \leq x \leq \sigma(\theta) + \frac{a_{n+1}}{\alpha_n} \right\}.$$

By the definitions of the maps Ω, q, π_{n+1} , the annulus A is contained in $\pi_n^{-1}(\Gamma_{n+1})$. Let $\Delta_0, \dots, \Delta_n$ be the $n+1$ continuous graphs $P_n(U_0), \dots, P_n(U_n)$, provided by property (v).2 for n . Let

$$U_{n+1} = P_n^{-1}(\text{Inte}(A) \setminus (\Delta_0 \cup \dots \cup \Delta_n)).$$

This is clearly an open set that is disjoint from U_0, \dots, U_n . Thus (ii) is satisfied for $n+1$. We now check the remaining properties at step $n+1$. By construction, we have $\pi_n(A) = \Gamma_{n+1}$, so that $P_n^{-1}(A) \subset \pi^{-1}(\Gamma_{n+1}) = B_{n+1}$. Hence, we have (i) for $n+1$. Note that for any θ

$$(\text{Inte}(A) \setminus (\Delta_0 \cup \dots \cup \Delta_n))_\theta = \text{Inte}(A_\theta) \setminus (\Delta_{0,\theta} \cup \dots \cup \Delta_{n,\theta}).$$

Thus this set is the union of at most $n + 1$ intervals. Hence, since $P_{n+1,\theta}$ is increasing, so is the set

$$U_{n+1,\theta} = P_{n,\theta}^{-1}(\text{Inte}(A_\theta) \setminus (\Delta_{0,\theta} \cup \dots \cup \Delta_{n,\theta})).$$

This gives (iv-bis). Since $U_{n+1,\theta}$ is disjoint from $U_{0,\theta} \cup \dots \cup U_{n,\theta}$, by the induction hypothesis (v).1 for n and the definition of A , we have

$$\text{Leb}(U_{n+1,\theta}) = \alpha_n \text{Leb}(\text{Inte}(A_\theta) \setminus (\Delta_{0,\theta} \cup \dots \cup \Delta_{n,\theta})) = \alpha_n \text{Leb}(A_\theta) = a_{n+1}.$$

This yields (iii) for $n + 1$. Furthermore $(P_{n,\theta})_*(\text{Leb}|_{U_{n+1,\theta}}) = \alpha_n \text{Leb}|_{A_\theta}$. Using the induction hypothesis (v).1 for n , we see that

$$(P_{n,\theta})_*(\text{Leb}|_{(U_{0,\theta} \cup \dots \cup U_{n+1,\theta})^c}) = \alpha_n \text{Leb}|_{A_\theta^c}.$$

As Ω_θ is just a rotation, and q_θ maps $\text{Leb}|_{\bar{A}_\theta^c}$ to $(1 - \frac{a_{n+1}}{\alpha_n})\text{Leb}$, the projection P_{n+1} satisfies property (v).1. By construction, $P_{n+1}(U_{n+1}) = \mathbb{T}^1 \times \{0\}$, and this implies property (v).2 for $n + 1$. \square

Proof of addendum 3.9. Let us fix $\varepsilon > 0$ and $n \in \mathbb{Z}$. For a given θ , the Lebesgue measure of $U_{n,\theta}$ is equal to a_n . Hence, there exists a closed set $V_\theta \subset \mathbb{T}^1$ contained in $U_{n,\theta}$ whose Lebesgue measure is larger than $(1 - \varepsilon)a_n$. Since U_n is open, for any θ' in a closed neighborhood W_θ of $\theta \in \mathbb{T}^1$ we still have $V_\theta \subset U_{n,\theta'}$. Consider $\theta_1, \dots, \theta_s$ such that \mathbb{T}^1 is covered by the $W_{\theta_1}, \dots, W_{\theta_s}$. The proposition now holds with the set

$$V_n = \bigcup_{i=1}^s W_{\theta_i} \times V_{\theta_i}.$$

\square

3.2 The measure ν

From now on, we assume that $\Gamma_n = R^n(\Gamma)$, where Γ is a curve having flat intersection with all its iterates. Furthermore, we consider a measure μ as defined by (3.3) and a map π as provided by proposition 3.3.

As mentioned at the beginning of this section, we construct the homeomorphism f via a measure ν on \mathbb{T}^2 . We will require that ν satisfies the following properties:

- (ν 1) For all $\theta \in \mathbb{T}^1$ the measure ν_θ is continuous (has no atom) and has full support.
- (ν 2) The mapping $\theta \mapsto \nu_\theta$ is continuous.
- (ν 3) $\pi_*\nu = R_*\mu$.
- (ν 4) In case R is a diffeomorphism which preserves the Lebesgue measure, ν has a continuous and positive density with respect to the Lebesgue measure.

Note that $R_*\mu = \beta \cdot R_*\text{Leb} + \sum_{n \in \mathbb{Z}} a_n \delta_{\Gamma_{n+1}}$.

3.2.1 Construction of ν : the continuous case

In order to construct the measure ν , we first note that, if $\epsilon > 0$, U_n and V_n are chosen as in proposition 3.7, then due to the Lemma of Urysohn there exist continuous functions $g_n : \mathbb{T}^2 \rightarrow [0, 1]$ such that

- (g1) $g_n^{-1}(0, 1] = U_n$;
(g2) $(1 - \epsilon)a_n \leq b_n(\theta) \leq a_n$ where $b_n(\theta) := \int_{U_{n,\theta}} g_n(\theta, x) dx$.

(The lower bound in (g2) can be ensured just by requiring $g_n|_{V_n} = 1$.) Now let

$$g(\theta, x) := \sum_{n \in \mathbb{Z}} \frac{a_n}{b_{n+1}(\theta)} \cdot g_{n+1}(\theta, x).$$

This positive function is in $\mathcal{L}_{\text{Leb}}^1(\mathbb{T}^2)$, since the $\mathcal{L}_{\text{Leb}}^1$ -norm of the n -th term in the sum is exactly a_n .⁴ For the same reason, for any θ , $x \mapsto g(\theta, x)$ is in $\mathcal{L}_{\text{Leb}}^1(\mathbb{T}^1)$ and has a norm bounded by a_n . For each n the mapping

$$\mathbb{T}^1 \rightarrow \mathcal{L}_{\text{Leb}}^1(\mathbb{T}^1) \quad , \quad \theta \mapsto g_n(\theta, \cdot)$$

is continuous, the same is true for $\theta \mapsto g(\theta, \cdot)$ (as the uniform limit of a sequence of continuous functions from \mathbb{T}^1 to $\mathcal{L}_{\text{Leb}}^1(\mathbb{T}^1)$). Consequently, if we let $\nu^1 := g\text{Leb}$, then $\theta \mapsto \nu_\theta^1$ is continuous with respect to the topology of weak convergence. Due to (g1) the function $x \mapsto g(\theta, x)$ is strictly positive on $\bigcup_{n \in \mathbb{Z}} U_{n,\theta}$, and due to (iii) in proposition 3.7 this set is dense in B_θ (remember that $B = \pi^{-1}(\Xi)$). Thus ν_θ^1 has full support in B_θ . By construction, the measure ν^1 projects to $\sum_{n \in \mathbb{Z}} a_n \delta_{\Gamma_{n+1}}$.

Further, as π is injective on $B^c = \pi^{-1}(\Xi^c)$ and $(R_*\text{Leb})(\Xi) = 0$, the measure ν^2 defined by $\nu^2(A) := \beta(R_*\text{Leb}) \circ \pi(A)$ is well-defined and obviously projects to $\beta(R_*\text{Leb})$. In addition, $\theta \mapsto \nu_\theta^2$ is continuous, and the fibre measures ν_θ^2 are continuous and have full support on $\text{int}(B_\theta^c)$. Altogether, this implies that

$$\nu := \nu^1 + \nu^2$$

satisfies $(\nu 1) - (\nu 3)$.

3.2.2 Construction of ν : the differentiable case

As above we suppose that ϵ, U_n and V_n are chosen as in propositions 3.7 and 3.9. Further, we assume that R is a diffeomorphism which preserves Lebesgue measure. Note that π projects $\text{Leb}|_{B^c}$ to βLeb , and $\text{Leb}|_{U_n}$ to $a_n \delta_{\Gamma_n}$. Since $R_*\mu = \beta\text{Leb} + \sum_{n \in \mathbb{Z}} a_n \delta_{\Gamma_{n+1}}$, we will construct the measure $\nu = h\text{Leb}$ by defining a continuous density h which satisfies $h|_{B^c} \equiv 1$ and

$$\int_{U_{n+1,\theta}} h(\theta, x) dx = a_n \quad \forall \theta \in \mathbb{T}^1, n \in \mathbb{Z}. \quad (3.5)$$

This is done by

$$h(\theta, x) := 1 - \sum_{n \in \mathbb{Z}} (a_{n+1} - a_n) \cdot \frac{g_{n+1}(\theta, x)}{b_{n+1}(\theta)}. \quad (3.6)$$

⁴Note that the function g is not continuous since $\frac{a_n}{b_{n+1}(\theta)}$ does not tend to 0.

It is obvious by construction that h satisfies (3.5) and therefore projects to $R_*\mu$.

It remains to show that, for a suitable choice of a_n in (3.3) and ϵ in proposition 3.7, the function h is continuous and strictly positive, such that $(\nu 4)$ (and thus also $(\nu 1)$ and $(\nu 2)$) hold. In order to see this, recall that $\|g_n\|_{C^0} = 1$. Further

$$\left| \frac{a_{n+1} - a_n}{b_{n+1}(\theta)} \right| \leq \frac{|a_{n+1} - a_n|}{(1 - \epsilon)a_n}, \quad (3.7)$$

and for a suitable choice of the sequence a_n (fixing any $\epsilon \in (0, 1)$ one can choose $a_n = (|n| + k)^{-2}$ for sufficiently large k), we have $\sum_{n \in \mathbb{Z}} a_n < 1$, the right side of (3.7) is strictly smaller than 1 and converges to 0 as $n \rightarrow \infty$. Since all the g_n have disjoint support, this implies that h will be continuous and strictly positive as required. Further, if k is large enough we also have $\sum_{n \in \mathbb{Z}} a_n < 1$.

3.2.3 $C^{1+\alpha}$ -Examples

Without going too much into detail, we remark that at least in the case where R leaves the Lebesgue measure invariant, as in the case of an irrational translation of the torus, it is possible to obtain examples with $C^{1+\alpha}$ fibre maps for any $\alpha \in (0, 1/2)$. For this, it suffices that the density h constructed above is α -Hölder continuous with respect to θ .

Although this is not explicitly stated in lemma 3.7, it is obvious from the construction of the sets U_n in the proof of this lemma that the functions $(\theta, x) \mapsto d(x, U_{n,\theta}^c)$ are continuous. In addition, due to (iv) in proposition 3.7, the sets $U_{n,\theta}$ consist of at most $2|n| + 1$ connected components on each fibre. Therefore, the functions

$$g_n(\theta, x) := \min \left\{ 1, \left(\frac{4|n| + 2}{\epsilon a_n} d(x, U_{n,\theta}^c) \right)^\alpha \right\}$$

satisfy (g1) and (g2). In addition, they are α -Hölder continuous with respect to θ with Hölder-constant $((4|n| + 2)/(\epsilon a_n))^\alpha$. In (3.6), the functions g_{n+1} are multiplied with a factor $\leq |a_{n+1} - a_n|/(1 - \epsilon)a_n$, such that the resulting product has α -Hölder constant $\leq \frac{((4|n| + 2)/\epsilon)^\alpha (a_{n+1} - a_n)}{(1 - \epsilon)a_n^{1+\alpha}}$. Hence, if we choose $(a_n)_{n \in \mathbb{Z}}$ such that

$$\sup_{n \in \mathbb{Z}} \frac{|n|^\alpha |a_{n+1} - a_n|}{a_n^{\alpha+1}} < \infty, \quad (3.8)$$

then the resulting sum in (3.6) is α -Hölder continuous (since the functions g_n all have disjoint support). However, (3.8) is true for any sequence $a_n = |n + k|^{-s}$ with $k \geq 1$ and $s \in (1, 1/\alpha - 1)$.

Of course, one should expect that the construction works for all $\alpha \in (0, 1)$. For this, one would need to show that the sets U_n in proposition 3.7 can be chosen such that the number of connected components is bounded by a constant independent of n . Since this would make the proof much more complicated, we refrained from doing so.

3.3 The homeomorphism f

In this section, we consider a projection π as provided by proposition 3.3 and a measure ν which satisfies $(\nu 1)$ – $(\nu 4)$, as constructed in the previous sections. We

denote by $\gamma_0, \gamma_1 : \mathbb{T}^1 \rightarrow \mathbb{T}^1$ the maps whose respective graphs are Γ_0 and Γ_1 , and define the two (discontinuous) maps φ_0^-, φ_1^- by letting $\varphi_i^-(\theta) = \text{Inf}(\pi^{-1}(\Gamma_i))$ (this has a meaning since \mathbb{T}^1 is an oriented circle).

Definition of f and verification of $\pi \circ f = R \circ \pi$. Due to (ν_1) , the map

$$\eta_\theta : \mathbb{T}^1 \rightarrow \mathbb{T}^1 \quad , \quad x \mapsto \nu_\theta[\varphi_1^-(\theta), x] \pmod{1}$$

is a homeomorphism. Using this fact, we let

$$f_\theta(x) := \eta_{\theta+\omega}^{-1}(\text{Leb}[\varphi_0^-(\theta), x]) \quad (3.9)$$

In other words, we simply define f by requiring that it maps φ_0^- to φ_1^- and sends the Lebesgue measure to ν , so that

$$\nu_{\theta+\omega}[\varphi_1^-(\theta + \omega), f_\theta(x)] = \text{Leb}[\varphi_0^-(\theta), x] . \quad (3.10)$$

For arbitrary $x_1, x_2 \in \mathbb{T}^1$, we obtain

$$\nu_{\theta+\omega}[f_\theta(x_1), f_\theta(x_2)] = \text{Leb}[x_1, x_2] . \quad (3.11)$$

As all fibre maps f_θ are circle homeomorphisms, the map f is bijective.

We remark that in the case where the fibre measures ν_θ have continuous (α -Hölder-continuous) densities, as in sections 3.2.2 and 3.2.3, it follows immediately from the definition that the fibre maps f_θ are \mathcal{C}^1 ($\mathcal{C}^{1+\alpha}$).

In order to verify the relation $\pi \circ f = R \circ \pi$, it suffices to check that for every $(\theta, x) \in \mathbb{T}^2$ there holds

$$\pi^{-1}(R(z)) = f(\pi^{-1}(z)) . \quad (3.12)$$

In order to do so, suppose $\pi_\theta^{-1}\{x\} = [\xi_0^-, \xi_0^+]$ and $\pi_{\theta+\omega}^{-1}\{R_\theta(x)\} = [\xi_1^-, \xi_1^+]$. We claim that

$$\nu_{\theta+\omega}[\varphi_1^-(\theta + \omega), f_\theta(\xi_0^\pm)] = \nu_{\theta+\omega}[\varphi_1^-(\theta + \omega), \xi_1^\pm] . \quad (3.13)$$

Since the measure $\nu_{\theta+\omega}$ has full support, this implies (3.12). For the left endpoints, we have

$$\begin{aligned} & \nu_{\theta+\omega}[\varphi_1^-(\theta + \omega), f_\theta(\xi_0^-)] \stackrel{(3.10)}{=} \\ & = \text{Leb}[\varphi_0^-(\theta), \xi_0^-] = \mu_\theta[\gamma_0(\theta), x] \\ & = (R_*\mu)_{\theta+\omega}[\gamma_1(\theta + \omega), R_\theta(x)] \stackrel{(\nu_3)}{=} \nu_{\theta+\omega}[\varphi_1^-(\theta + \omega), \xi_1^-] . \end{aligned}$$

The argument for the right endpoints is similar, which proves (3.13).

Continuity of f . We first show that f is continuous in (θ, x) if $(\theta, x) \notin B$. Recall that this means that $\pi(\theta, x)$ is not contained in $\Xi = \bigcup_{n \in \mathbb{Z}} R^n(\Gamma)$, and consequently the same is true for the point $z = R \circ \pi(\theta, x)$. Therefore $z = R \circ \pi(\theta, x)$ has a unique preimage under π . As $R \circ \pi = \pi \circ f$, this preimage is $f(\theta, x)$. Since π is continuous and due to compactness, it follows that for any point z' which is sufficiently close to z , the preimage $\pi^{-1}(\{z'\})$ is contained in

a small neighbourhood of $f(\theta, x)$. Now suppose (θ', x') is close to (θ, x) . Then $z' := R \circ \pi(\theta', x')$ is close to z by continuity. As $f(\theta', x')$ must be contained in $\pi^{-1}(\{z'\})$, this shows that $f(\theta', x')$ is close to $f(\theta, x)$. Hence f is continuous in (θ, x) .

Now let $(\theta, x) \in \mathbb{T}^2$ be arbitrary. The set B^c intersects every fibre, since this is obviously true for its image Ξ^c under π . Consequently, there exists a continuity point (θ, x_0) of f on the same fibre. Suppose that (θ', x') is a point close to (θ, x) . Then (3.11) implies that $\nu_{\theta'+\omega}([f_{\theta'}(x_0), f_{\theta'}(x')])$ is close to $\nu_{\theta+\omega}([f_{\theta}(x_0), f_{\theta}(x)])$. As f is continuous in (θ, x_0) , the points $f_{\theta'}(x_0)$ and $f_{\theta}(x_0)$ are close. Therefore, the fact that $\nu_{\theta+\omega}$ has full support and the continuity of $\theta \mapsto \nu_{\theta}$ imply that $f_{\theta'}(x')$ has to be close to $f_{\theta}(x)$. Thus, f is continuous on all of \mathbb{T}^2 .

3.4 Proof of addendum 3.2

Suppose that R is transitive, Γ has the additional property provided by addendum 2.3 and Ξ is dense. We have to show that given any two open sets $U, V \subseteq \mathbb{T}^2$ there exists some $n \in \mathbb{N}$ such that $f^n(U) \cap V \neq \emptyset$.

First, we claim that the interior of $B = \pi^{-1}(\Xi)$ is dense. For suppose that there exists an open set $W \subseteq B^c$. For any point $z \in W$, since $\pi^{-1}\{z\}$ is a singleton, compactness and continuity of π imply that there exists some neighbourhood \tilde{W} of $\pi(z)$, such that $\pi^{-1}(\tilde{W}) \subset W$. However, as \tilde{W} is clearly disjoint from Ξ , this contradicts the assumptions.

Thus, by reducing both sets further if necessary, we can assume that U and V are two small rectangles which are included in B . In fact, we can even restrict to the case where $U \subseteq B_k$ and $V \subseteq B_l$ for some $k, l \in \mathbb{Z}$, and that both $I := p_1(U)$ and $J := p_1(V)$ are intervals. However, due to the choice of Γ there exists some $n \in \mathbb{N}$ such that $R^n(\Gamma_{k|I})$ crosses $\Gamma_{l|J}$ over $I + n\omega \cap J$. As all the maps π_{θ} are order-preserving this implies that $f^n(U)$ has to cross V over $I' := (I + n\omega) \cap J$ (more precisely : if O is chosen as in definition 2.1, then $\pi^{-1}(I' \times O) \setminus V$ consists of two connected components, and $f^n(U)$ intersects both of them). However, this is only possible if $f^n(U)$ intersects V .

Remark 3.10. *If R is transitive, it is always possible to construct a curve Γ , such that Ξ is dense, as required by addendum 3.2 . For this, it suffices to choose each Γ_n in the construction of Γ in section 2, such that it contains a point with dense orbit. This is possible due to remark 2.9 . It follows that for each $n \in \mathbb{N}$ there exists an integer N_n , such that the first N_n iterates of Γ_n are $1/n$ -dense in \mathbb{T}^2 . If all subsequent perturbations are chosen small enough, then for all $k \geq n$ the first N_n iterates of Γ_k will be $2/n$ -dense. In the limit, this gives the required property.*

4 The minimal set

In this section, we collect some general results which concern the properties and structure of the minimal sets of a transitive but non minimal qpf circle homeomorphism. It is known that in the absence of invariant strips such a set must have a complicated structure:

Proposition 4.1 ([7, theorem 4.5 and lemma 4.6]). *Suppose $f \in \mathcal{F}$ has no invariant strips and is transitive, but not minimal. Then any minimal invariant set M which is a strict subset of \mathbb{T}^2 has the following two properties:*

- (a) *Every connected component C of M is a vertical segment, i.e. $\#p_1(C) = 1$.*
- (b) *For every open set $U \subseteq \mathbb{T}^2$, the set $p_1(U \cap M)$ is either empty or it contains an interval.*

4.1 Uniqueness and structure

We start by giving two criteria for the uniqueness of the minimal set. (For a previous partial result on quasiperiodic $\mathrm{SL}(2, \mathbb{R})$ -cocycles, see [5, section 4.17].)

Proposition 4.2. *Suppose $f \in \mathcal{F}$ has no invariant strip, or is transitive. Then there is only one f -invariant minimal set.*

Proof. We choose an orientation on the circle, so that the segment (x, y) is well defined whenever $x, y \in \mathbb{T}^1$. Suppose that there exist two minimal sets $K \neq K'$. Then $K \cap K' = \emptyset$. We define the set U_1 as the union of all vertical segments $\{\theta\} \times (x, y)$ which are disjoint from $K \cup K'$ and satisfy $(\theta, x) \in K$ and $(\theta, y) \in K'$. Similarly we define U_2 as the union of all vertical segments $\{\theta\} \times (x, y)$ which are disjoint from $K \cup K'$ and satisfy $(\theta, x) \in K'$ and $(\theta, y) \in K$. These two sets are clearly disjoint and intersect each fibre $\{\theta\} \times \mathbb{T}^1$. As all fibre maps f_θ are order preserving, they are also invariant. Let us prove that they have non-empty interior. The mappings $\theta \mapsto K_\theta$ and $\theta \mapsto K'_\theta$ are semi-continuous, hence their sets of continuity points are two dense G_δ -sets; in particular they have a common point θ_0 . It is easy to see that all the points (θ_0, x) in U_1 belong to the interior of U_1 , similar for U_2 . Thus $\mathrm{Int}(U_1)$ and $\mathrm{Int}(U_2)$ are two disjoint non-empty open invariant sets, and f is not transitive.

It remains to consider the case when f has no invariant strip and is not transitive. By theorem 1.1, f is semi-conjugate to an irrational torus rotation. Therefore, due to a result by Furstenberg (see [12, theorem 4.1]), f is uniquely ergodic, hence it has a unique minimal set. \square

Concerning the number of connected components in each fibre, we have the following:

Proposition 4.3. *Let f be a qpf circle homeomorphism, and K a minimal set for f . Let $c(\theta) \in \mathbb{N} \cup \{\infty\}$ be the number of connected components of K_θ . Let $c(K) = \inf_{\theta \in \mathbb{T}^1} c(\theta)$. Then $c(\theta) = c(K)$ on a dense G_δ subset of the circle.*

Proof. Let Θ_1 be the set of continuity points of the semi-continuous map $\theta \mapsto K_\theta$: this is a dense G_δ subset of the circle. The restriction of the map $\theta \mapsto c(\theta)$ to Θ_1 is again semi-continuous, so it admits a continuity point θ_0 . By continuity there exists an open neighbourhood U of θ_0 such that c is constant on $U \cap \Theta_1$. Since c is invariant under the circle rotation $\theta \mapsto \theta + \omega$, it is constant on a dense open subset Θ_2 of Θ_1 , and Θ_2 is again a dense G_δ subset of the circle. Let c_0 be the value of c on Θ_2 , and let us prove that $c_0 = c(K)$. For this consider a fibre $\theta_0 \in \Theta_2$. Since θ_0 is a continuity point of $\theta \mapsto K_\theta$, there exists a neighbourhood U of θ_0 on which $c(\theta) \geq c_0$. Using the invariance of c under the circle rotation, we see that this inequality holds on the whole circle. \square

This result raises the following question: *for the unique minimal set K of a non minimal qpf circle map with no invariant strip, can $c(K)$ be finite ?*

4.2 Examples

Proposition 4.4.

1. There exists a transitive non minimal qpf circle homeomorphism whose minimal set K is a Cantor set and its intersection with any fibre $\{\theta\} \times \mathbb{T}^1$ is uncountable; in particular $c(K) = +\infty$.
2. There exists a transitive non minimal qpf circle homeomorphism whose minimal set contains a vertical segment.

Proof of item 1. Choose a minimal qpf circle homeomorphism R , and let Γ be a continuous graph constructed by proposition 2.2 and addendum 2.3. Let f be a qpf circle homeomorphism given by proposition 3.1, which in particular satisfies $\pi \circ f = R \circ \pi$ and is transitive but not minimal. Let $\Xi = \bigcup_{n \in \mathbb{Z}} R^n(\Gamma)$. Then we claim that the unique minimal set K for f is the set $\overline{\pi^{-1}(\Xi^c)}$: indeed, since R is minimal $\pi(K) = \mathbb{T}^2$, thus K contains $\pi^{-1}(\Xi^c)$ by (iii) of proposition 3.1.

To construct the first example we suppose that R has no invariant strip and that for each θ , the set Ξ_θ is dense in the circle: in particular this is always true if R is a irrational rotation. It follows from items (ii) and (iii) of proposition 3.1 together with remark 3.8(b) that the set $\text{Int}(\pi^{-1}(\Xi))$ intersects each fibre in an open dense set. In particular, K_θ has empty interior. Since Ξ_θ^c is uncountable, so is K_θ . Since R has no invariant strip, neither has f . Hence every connected component of K is included in a fibre (proposition 4.1); consequently K is totally disconnected. Since K is a minimal infinite set, it is perfect, so K is a Cantor set. \square

Sketch of proof of item 2. To construct the second example, we start with an irrational rotation R_0 and choose a curve Γ_0 as provided by proposition 2.2 and its addendum. We consider a point (θ_0, x_0) whose orbit by R_0 is disjoint from Γ_0 . Applying Rees construction in [13, 15] allows to build a minimal fibred homeomorphism R that is semi-conjugate to R_0 by a semi-conjugacy Φ such that $\Phi^{-1}(\theta, x)$ is a nontrivial vertical segment if (θ, x) belongs to the orbit \mathcal{O} of (θ_0, x_0) under R_0 and a single point otherwise. Since $\Phi^{-1}(\mathcal{O})$ has empty interior and R_0 is minimal, R is minimal. Now we let $\Gamma = \Phi^{-1}(\Gamma_0)$; Γ is a continuous graph whose iterates are disjoint from the non-trivial vertical segment $I = \Phi^{-1}(\theta_0, x_0)$. Furthermore, one can check that R and Γ still satisfy the conclusions of proposition 2.2 and its addendum. We now apply proposition 3.1. Thus we get a map f whose minimal set K contains the non-trivial vertical segment $\pi^{-1}(I)$. \square

4.3 Ergodic measures

In this subsection, we briefly want to discuss, in a rather informal way, the consequences of our construction in sections 2 and 3 for the structure of the invariant measures of the system. First, we recall an old result by Furstenberg [12], which may be seen as a measure-theoretic counterpart to theorem 1.1. In order to state it, we note that p_1 maps any ergodic invariant probability

measure μ of a qpf circle homeomorphism f to the Lebesgue measure on \mathbb{T}^1 , since this is the only invariant probability measure for the underlying irrational circle rotation. Consequently, μ can be decomposed as

$$\mu(A) = \int_{\mathbb{T}^1} \mu_\theta(A_\theta) d\theta ,$$

where the probability measures μ_θ are the conditional measures of μ with respect to the σ -algebra $p_1^{-1}(\mathcal{B}(\mathbb{T}^1))$.

Theorem 4.5 (Furstenberg, [12, theorem 4.1]). *Either f is uniquely ergodic and for almost every θ the measure μ_θ is continuous, or there exists $n \in \mathbb{N}$ such that for almost every θ the measure μ_θ is the equidistribution on n points.*

In the second case, one obtains a measurable invariant set $\Phi := \{(\theta, x) \mid \mu_\theta\{x\} > 0\}$, which contains exactly n points in every fibre.⁵ Since such a set can always be represented as the graph of a measurable n -valued function φ , one speaks of an *invariant graph*. Conversely, every invariant graph determines an ergodic invariant measure μ_Φ , given by

$$\mu_\Phi(A) = \int_{\mathbb{T}^1} \frac{\#(A_\theta \cap \Phi_\theta)}{n} d\theta .$$

(We remark that the requirement that this measure is ergodic is part of the definition of an invariant graph.) For a more detailed discussion of these concepts, see [7, section 2].

The important fact in our context is that the two alternative cases of theorem 4.5 are preserved by topological extension: If f is semi-conjugate to R via a fibre-respecting semi-conjugacy π , then π projects f -invariant graphs to R -invariant graphs. Conversely, the preimage of any R -invariant graph under π intersects every fibre in exactly n connected components, and the endpoints of the latter constitute invariant graphs for the topological extension f .

In order to describe how our construction affects the invariant measures, we place ourselves in the situation of proposition 3.1 and consider the two cases in theorem 4.5 .

1) When the original transformation R has a unique invariant measure with continuous fibre measures, then nothing much happens. The topological extension f will still have a unique invariant measure with continuous fibre measures. The only effect is that the new invariant measure does not have full topological support in \mathbb{T}^2 (since f is not minimal).

The question of what happens with an invariant graph Φ when passing from R to f in proposition 3.1 (and hence in theorem 1.3) mainly depends on the value of $\mu_\Phi(\Gamma)$. Thus, we have to distinguish to subcases.

2a) When $\mu_\Phi(\Gamma) = 0$, nothing changes either. The preimage of Φ under π constitutes an invariant graph for the new map f , and the two systems (R, μ_Φ) and $(f, \mu_{\pi^{-1}\Phi})$ are isomorphic in the measure-theoretic sense.

2b) The more interesting case is the one where $\mu_\Phi(\Gamma) > 0$. In this case, since μ_Φ is ergodic and $\Xi = \bigcup_{n \in \mathbb{Z}} R^n(\Gamma)$ is invariant, we have $\mu_\Phi(\Xi) = 1$.

⁵*A priori*, this only holds on almost every fibre. However, by modifying μ_θ on a set of measure zero, one can always assume without loss of generality that it holds on every fibre.

Consequently, the preimage $\pi^{-1}(\theta, x)$ of almost every point in Φ is a vertical segment. The endpoints of these segments constitute two distinct invariant graphs for f , such that the invariant measure μ_Φ has been (at least) doubled. *A more difficult question, which we have to leave open here, is the one whether there exist further invariant graphs in the preimage of Φ .*

Without going into detail, we remark that both conditions $\mu_\Phi(\Gamma) = 0$ and $\mu_\Phi(\Gamma) > 0$ can be ensured by adapting the construction of the curve Γ in the proof of proposition 2.2. In the former case, the crucial fact is that for any $\epsilon > 0$, it is always possible to render any segment of a given curve disjoint from Φ on a set of fibres of measure arbitrarily close to 1 by an ϵ -perturbation. Performing an infinite sequence of smaller and smaller perturbations, this shows that in any arbitrarily small box there exist continuous curve segments which intersect Φ only on a set of fibres of measure zero. Using this fact appropriately in each step of the construction, namely when the modifications are constructed in the proof of perturbation lemma 2.7 (compare also remark 2.9), allows to ensure that the limit curve obtained in proposition 2.2 satisfies $\mu_\Phi(\Gamma) = 0$.

Conversely, in order to ensure $\mu_\Phi(\Gamma) > 0$, it suffices to start the construction in section 2 with a curve Γ_0 which intersects Φ on a set of positive measure. This is always possible, since there exists compact sets $K \subseteq \mathbb{T}^1$ of measure arbitrarily close to 1 with the property that the restriction of φ to K is continuous (where φ is the measurable function $\mathbb{T}^1 \rightarrow \mathbb{T}^1$ with graph Φ as above). If the modifications in each step of the construction are then performed only over sets of sufficiently small measure, then the resulting limit curve Γ will still intersect Φ on a set of positive measure. It is even possible to ensure this condition for any finite number of invariant graphs at the same time.

Finally, we want to mention that it is possible to repeat this “doubling procedure” as many times as wanted, producing a sequence of topological extensions f_1, f_2, \dots of R with more and more invariant graphs in the preimage of an initial R -invariant graph. Furthermore, the C^0 -distance between both f_n and f_{n+1} and the corresponding semi-conjugacies can be made arbitrarily small in each step. Then the f_n converge to a limit f_∞ , which is a topological extension of R with an infinite number of invariant graphs that project down to Φ .

4.4 The linear case

Finally, we restrict ourselves to qpf linear circle homeomorphisms. We identify the 2-torus \mathbb{T}^2 with $\mathbb{T}^1 \times \mathbb{P}^1(\mathbb{R})$ and consider the projective action of $\mathrm{SL}(2, \mathbb{R})$ on $\mathbb{P}^1(\mathbb{R})$. Then a *qpf linear circle homeomorphism* is a homeomorphism of the 2-torus, isotopic to the identity, of the form $(\theta, x) \mapsto (\theta + \omega, f_\theta(x))$ where $f_\theta \in \mathrm{SL}(2, \mathbb{R})$.

Proposition 4.6. *Let f be a qpf linear circle homeomorphism. Assume f is transitive but not minimal. Let K be the unique minimal invariant set for f . Then K_θ contains at most two points for every θ in a dense G_δ subset of the circle. In particular we have $c(K) = 1$ or 2 in proposition 4.3.*

In case there is no invariant strip, the preceding result can be improved further. Let us recall that we know no example of a qpf linear circle homeomorphism with no invariant strip which is not minimal (so that the following

statement could turn out to be void).

Proposition 4.7. *Let f be a qpf linear circle homeomorphism. Assume f is not minimal and has no invariant strip. Let K be the unique minimal invariant set for f . Then K_θ is a singleton for every θ in a dense G_δ subset of the circle. In particular we have $c(K) = 1$ in proposition 4.3.*

In the case where f has invariant strips, one has the following result:

Proposition 4.8 ([16]). *Let f be a qpf linear circle homeomorphism. Assume f has invariant strips and let K be a minimal invariant set for f . Then there are two (non exclusive) possibilities:*

1. K is a continuous (p, q) -invariant graph, or
2. for every θ in a dense G_δ subset of the circle the cardinality of the set K_θ is 1 or 2. In particular we have $c(K) = 1$ or 2 in proposition 4.3.

This result is in [16] formulated for continuous time systems, but the translation to the discrete time case and vice versa is plain sailing. We provide a new proof of it. The authors in [16] also show that any minimal set belongs to one of five different cases, but for one of the possibilities (which they call “Denjoy extension”) they leave open whether it can be realised or not. The proposition 4.6 above excludes the existence of such Denjoy extensions.

Proposition 1.6 now follows from propositions 4.7 and 4.8: strict minimal set K for a qpf linear circle homeomorphism f , if f has no invariant strip then for a generic θ the cardinality of K_θ is 1 (proposition 4.7); if f has invariant strips, then the cardinality of K_θ is generically 1 or 2, or K is a (p, q) -invariant graph (proposition 4.8). Note that if the cardinality of K_θ is generically greater than 2 then we are in the last case with $pq > 2$, and then it is easy to see that f is actually conjugate to a rotation.

Proof of proposition 4.6. Let f and K be as in the proposition. Let Θ_1 be the set of $\theta_0 \in \mathbb{T}^1$ such that θ_0 is a continuity point of $\theta \mapsto K_\theta$, and such that for a dense set of $x_0 \in \mathbb{T}^1$ the positive orbit of (θ_0, x_0) is dense in \mathbb{T}^2 . Given the transitivity of f , the set of points whose positive orbit is dense is a dense G_δ subset of the 2-torus. Thus Θ_1 contains a dense G_δ subset of the circle.⁶

We now suppose that there exists some $\theta_0 \in \Theta_1$ such that K_{θ_0} is not a single point. We choose a connected component (a, b) of $\mathbb{T}^1 \setminus K_{\theta_0}$; note that $a \neq b$.

Claim 4.9. *Every map $A \in \text{SL}(2, \mathbb{R})$ that fixes a and b globally fixes K_{θ_0} .*

In order to prove the claim, we introduce the maps $f_\theta^n \in \text{SL}(2, \mathbb{R})$ defined by $f_\theta^n(\theta, x) = (\theta + n\omega, f_\theta^n(x))$. By definition of Θ_1 there exists a point $z_0 = (\theta_0, x_0) \in \{\theta_0\} \times (a, b)$ with a dense positive orbit. Let $z = (\theta_0, A(x_0))$, and consider an increasing sequence (n_i) such that $f_{\theta_0}^{n_i}(z_0)$ converges to z . By definition of Θ_1 , the sequence $f_{\theta_0}^{n_i}(K_{\theta_0})$ converges to K_{θ_0} . This implies that $f_{\theta_0}^{n_i}(a)$ converges to a and $f_{\theta_0}^{n_i}(b)$ converges to b . The map $g \mapsto (g(a), g(z_0), g(b))$ is a homeomorphism between $\text{SL}(2, \mathbb{R})$ and the space of cyclically ordered triples.

⁶Recall that for any dense G_δ subset E of \mathbb{T}^2 , there exists a dense G_δ subset of points $\theta \in \mathbb{T}^1$ for which E_θ is a dense G_δ subset in \mathbb{T}^1 .

Thus the sequence $f_{\theta_0}^{n_i}$ converges to A , and $A(K_{\theta_0}) = K_{\theta_0}$, and the claim is proved.

Since $\mathrm{SL}(2, \mathbb{R})$ acts transitively on positively ordered triplets, this claim implies that K_{θ_0} either contains $[b, a]$ or is disjoint from (b, a) . In the first case K_{θ_0} would contain a point with dense positive orbit and K would equal \mathbb{T}^2 , a contradiction. Thus we have proved that for every $\theta_0 \in \Theta_1$ the set K_{θ_0} contains either one or two points. \square

Proof of proposition 4.7. First of all, suppose that f is semi-conjugate to an irrational rotation. Then it follows from the classification for the dynamics of qpf linear circle homeomorphisms in [10] that the semi-conjugacy has linear fibre maps (i.e. $\pi_\theta \in \mathrm{SL}(2, \mathbb{R}) \forall \theta \in \mathbb{T}^1$), which further implies that the unique f -invariant measure is equivalent to the Lebesgue measure. Hence f is minimal. Due to theorem 1.1, this means that we can assume, without loss of generality, that f is ρ -unbounded and transitive.

We argue by contradiction. Applying proposition 4.6, we see that K_θ contains exactly two points for a dense G_δ subset Θ_2 of the circle. We can also assume that every point in Θ_2 is a continuity point of $\theta \mapsto K_\theta$.

We fix some $\theta_0 \in \Theta_2$. Up to a linear fibred conjugacy, we can assume that $K_{\theta_0} = \{1/8, 5/8\}$. By continuity we can choose an open interval I containing θ_0 such that for each $\theta \in I$, K_θ does not meet $[1/4, 1/2]$ neither $[3/4, 1]$, and meets both $(0, 1/4)$ and $(1/2, 3/4)$.

We choose a lift F of f to $\mathbb{T}^1 \times \mathbb{R}$, and consider the lift \tilde{K} of K . By the choice of I , for every $z \in \tilde{K} \cap (I \times \mathbb{R})$ there exists $\tau(z) \in \frac{1}{2}\mathbb{Z}$ such that $z \in I \times (\tau(z), \tau(z) + 1/4)$. Now let θ, n be such that θ and $\theta + n\omega$ belong to $I \cap \Theta_2$. The definitions of I and Θ_2 entails that, for a fixed value of θ , the number $\tau(F^n(z)) - \tau(z)$ is constant on \tilde{K}_θ . We denote this number by $d_n(\theta)$.

Claim 4.10. *There exist an interval $J \subset I$ and an integer n such that $J + n\omega \subset I$ and the function d_n is not constant on $J \cap \Theta_2$.*

Let us prove the claim. Let J be any compact non-trivial interval inside I . There exists $n_0 > 0$ such that every interval J' with the same length as J has an iterate $J' + m\omega$ inside I with $0 \leq m \leq n_0$. Since f is not ρ -bounded, the vertical diameter of the sequence of iterates $F^n(J \times \{0\})$ is not bounded. Thus there exists $N > 0$ such that for every $m = 0, \dots, n_0$ the vertical diameter of $F^{N+m}(J \times \{0\})$ is greater than 1. By the choice of n_0 we can choose such an m with $J + (N + m)\omega \subset I$. Let $n = N + m$, then d_n cannot be constant on $J \cap \Theta_2$ since this would imply that the vertical diameter of $F^n(J \times \{0\})$ is smaller than $3/4$. This proves the claim.

For every half integer k let us consider the set $J_k = \overline{d_n^{-1}\{k\}}$. The interval J is contained in the union of finitely many J_k 's, thus the previous claim implies that there exist $k \neq k'$ and $\theta_1 \in J \cap J_k \cap J_{k'}$. As θ_1 is a limit point of $d_n^{-1}(k)$, there exist $x_1 \in (0, 1/4)$ and $x_2 \in (1/2, 3/4)$ such that $F_{\theta_1}^n(x_1) \in (k, k + 1/4)$ and $F_{\theta_1}^n(x_2) \in (k + 1/2, k + 3/4)$. Similarly there exist $x'_1 \in (0, 1/4)$ and $x'_2 \in (1/2, 3/4)$ such that $F_{\theta_1}^n(x'_1) \in (k', k' + 1/4)$ and $F_{\theta_1}^n(x'_2) \in (k' + 1/2, k' + 3/4)$. To fix ideas we assume that $x_1 < x'_1$. Since $F_{\theta_1}^n$ is order-preserving this implies that $k' > k$, and thus we also have $x_2 < x'_2$. Since $x'_1 < x_2$ we see that actually

$k' = k + 1/2$. Thus we have

$$F_{\theta_1}^n((x_1x'_1)) \supset \left[k + \frac{1}{4}, k + \frac{1}{2} \right] \text{ and } F_{\theta_1}^n((x_2x'_2)) \supset \left[k + \frac{3}{4}, k + 1 \right].$$

We consider the projective circle homeomorphism $\varphi : x \mapsto f_{\theta_1}^n(x) - 1/4 - k$. Both intervals $[0, 1/4]$ and $[1/2, 3/4]$ are contained in the interior of their image by φ , and both intervals $[1/4, 1/2]$ and $[3/4, 1]$ are contained in the interior of their pre-image: thus φ is not the identity but has a fixed point inside the interior of each of these four intervals. This is absurd since φ is a projective map. \square

Proof of proposition 4.8. Let K be a minimal set. The following properties are well known.

Claim 4.11. *There exists a compact invariant set $A \subseteq \mathbb{T}^2$ and $n \geq 1$ such that*

- $K \subseteq A$;
- for all $\theta \in \mathbb{T}^1$ the set A_θ consists of exactly n connected components;
- K_θ coincides with A_θ and contains exactly n points whenever θ is a continuity point of the mapping $\theta \mapsto K_\theta$.

In particular, for all θ in a dense G_δ subset of \mathbb{T}^1 , the cardinality of K_θ is equal to n .

Briefly spoken, this is due to the general fact that if f has invariant strips, then every minimal set is contained in an invariant strip, which can furthermore be chosen minimal with respect to the inclusion amongst all invariant strips. Such a minimal invariant strip automatically has the required properties of the set A above. In order to give precise references, we argue as follows: Recall that, by assumption, f has an invariant strip. It follows from [7, Lemma 3.9] that

$$\rho(f) = \frac{k}{q}\omega + \frac{l}{p} \pmod{1}$$

for suitable integers $k, q, l, p \in \mathbb{Z}$, $q, p \neq 0$ (which are further specified in [7]). If we go over to a suitable iterate, consider a lift of f to the q -fold cover $(\mathbb{R}/q\mathbb{Z}) \times \mathbb{T}^1$ and perform a conjugacy of the form $(\theta, x) \mapsto (\theta, x - m\theta)$ with suitable $m \in \mathbb{Z}$, then we can assume that $\rho(f) = 0$. Therefore, we can choose a lift $F : \mathbb{T}^1 \times \mathbb{R} \rightarrow \mathbb{T}^1 \times \mathbb{R}$ with $\rho(F) = 0$. Let \tilde{K} be the ω -limit set of any point $(\theta, x) \in \mathbb{T}^1 \times \mathbb{R}$ which lifts a point of K . Since F -orbits are bounded due to the bounded deviations, \tilde{K} is a minimal F -invariant set, which projects down to K . It follows from the results in [18], for example corollary 4.4, that the set

$$\tilde{A} := \{(\theta, x) \mid \inf \tilde{K}_\theta \leq x \leq \sup \tilde{K}_\theta\}$$

is reduced to a point (and in particular coincides with \tilde{K}) on all fibres which are continuity points of $\theta \mapsto \tilde{K}_\theta$. The facts that $\tilde{K} \subseteq \tilde{A}$ and that for all $\theta \in \mathbb{T}^1$ the set \tilde{A}_θ consists of exactly one connected component follows directly from the definition. By projecting \tilde{A} to the torus and redoing the transformations described above, we obtain the required set A . This proves the claim.

Now suppose $n > 2$. We have to show that K is the graph of a continuous n -valued curve in this case. If the cardinality of K_θ equals n for any $\theta \in \mathbb{T}^1$, then this follows easily. Hence, suppose for a contradiction that there exists some $\theta_0 \in \mathbb{T}^1$ with $\#K_{\theta_0} \geq n + 1$. Further, fix some $\theta_1 \in \mathbb{T}^1$ which is a continuity point of $\theta \mapsto K_\theta$. Note that this implies $\#K_{\theta_1} = c(K) = n$, and let $K_{\theta_1} = \{x_1, \dots, x_n\}$. Let $A \supseteq K$ be as above and suppose $A_{\theta_0}^1, \dots, A_{\theta_0}^n$ are the n connected components of A_{θ_0} . Let n_j be an increasing sequence of integers, such that $\theta_0 + n_j\omega \rightarrow \theta_1$. By going over to a subsequence and relabelling if necessary, we can assume that each of the sequences $f_{\theta_0}^{n_j}(A_{\theta_0}^i)$ converges to the point x_i . (Note that the continuity of $\theta \mapsto K_\theta$ in θ_1 implies that the limit points of different connected components of A_{θ_0} are distinct.) Let us choose arbitrary points $y_1 \in A_{\theta_0}^1, y_2 \in A_{\theta_0}^2, y_3 \in A_{\theta_0}^3$. As all $f_{\theta_0}^{n_j}$ are the projective actions of $\text{SL}(2, \mathbb{R})$ -matrices, this implies that the $f_{\theta_0}^{n_j}$ themselves converge to the linear circle homeomorphism g which maps (y_1, y_2, y_3) on (x_1, x_2, x_3) . However, this leads to a contradiction: at least one of the intervals $A_{\theta_0}^i$ must be non-degenerate (since $K_{\theta_0} \subseteq A_{\theta_0}$ and $\#K_{\theta_0} \geq n + 1$), but in the limit it is contracted by g to a single point x_i . □

References

- [1] T. Jäger and J. Stark. Towards a classification for quasiperiodically forced circle homeomorphisms. *Journal of the LMS*, 73(3):727–744, 2006.
- [2] S. Aubry and G. André. Analyticity breaking and Anderson localization in incommensurate lattices. *Proc. 8th Int. Coll. on Group Theoretical Methods in Physics, Kiryat Anavim, Israel 1979 (Bristol: Hilger)*. *Ann. Isr. Phys. Soc.*, 3:133–164, 1980.
- [3] A. Haro and J. Puig. Strange non-chaotic attractors in harper maps. *Chaos*, 16, 2006.
- [4] Mingzhou Ding, Celso Grebogi, and Edward Ott. Evolution of attractors in quasiperiodically forced systems: From quasiperiodic to strange nonchaotic to chaotic. *Physical Review A*, 39(5):2593–2598, 1989.
- [5] Michael R. Herman. Une méthode pour minorer les exposants de Lyapunov et quelques exemples montrant le caractère local d’un théorème d’Arnold et de Moser sur le tore de dimension 2. *Commentarii Mathematici Helvetici*, 58:453–502, 1983.
- [6] L. G. Shnirelman. An example of a transformation of the plane. *Proc. Don Polytechnic Inst. (Novochekassk)*, 14 (Science section, Fis-math. part), 1930.
- [7] T. Jäger and G. Keller. The Denjoy type-of argument for quasiperiodically forced circle diffeomorphisms. *Ergodic Theory and Dynamical Systems*, 26(2):447–465, 2006.
- [8] A. Katok and B. Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press, 1997.
- [9] K. Bjerklöv. Positive Lyapunov exponent and minimality for a class of one-dimensional quasi-periodic Schrödinger equations. *Ergodic Theory and Dynamical Systems*, 25:1015–1045, 2005.

- [10] P. Thieullen. Ergodic reduction of random products of two-by-two matrices. *Journal d'Analyse Mathématique*, 73:19–64, 1997.
- [11] A. Avila and R. Krikorian. Reducibility or non-uniform hyperbolicity for quasiperiodic Schrödinger cocycles. *Annals of Mathematics (2)*, 164(3):910–941, 2006.
- [12] H. Furstenberg. Strict ergodicity and transformation of the torus. *American Journal of Mathematics*, 83:573–601, 1961.
- [13] M. Rees. A point distal transformation of the torus. *Israel J. Math.* 32:201–208, 1979.
- [14] M. Rees. A minimal positive entropy homeomorphism of the 2-torus. *Journal of the LMS (2)*, 23:537–550, 1981.
- [15] F. Béguin, S. Crovisier and F. Le Roux. Construction of curious minimal uniquely ergodic homeomorphisms on manifolds: the Denjoy-Rees technique. *Ann. Sci. École Norm. Sup. (4)*, 40:191–350, 2007.
- [16] K. Bjerklöv and R. Johnson. Minimal subsets of projective flows. Preprint, 2007.
- [17] K. Bjerklöv and T. Jäger. Rotation numbers for quasiperiodically forced circle maps - mode-locking vs strict monotonicity. Preprint, 2006, arXiv:math/0607598.
- [18] J. Stark. Transitive sets for quasiperiodically forced monotone maps. *Dynamical Systems*, 18(4):351–364, 2003.