

TD 3 - QUILLEN FUNCTORS, DERIVED FUNCTORS AND (A BIT OF) HOMOTOPY COLIMITS

Exercise 1 (Two-out-of-six). Let \mathcal{W} be a collection of arrows of \mathcal{C} . We say \mathcal{W} satisfies 2-out-of-3 if for every couple of composable arrows f, g , any two out of the three $f, g, g \circ f$ being in \mathcal{W} implies the third is. We say \mathcal{W} satisfies 2-out-of-6 if for every triple of composable arrows f, g, h such that $g \circ f \in \mathcal{W}$ and $h \circ g \in \mathcal{W}$, then the four other maps $f, g, h, h \circ g \circ f \in \mathcal{W}$.

1. Let \mathcal{W} be a collection of arrows of \mathcal{C} satisfying 2-out-of-6. Show \mathcal{W} satisfies 2-out-of-3.
2. (a) Show the collection of isomorphisms of a category always satisfies 2-out-of-3. Does it also satisfy 2-out-of-6?
 (b) Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor and let $\mathcal{I}_{\mathcal{C}}$ be the collection of isomorphisms of \mathcal{C} . Show $F(\mathcal{I}_{\mathcal{C}})$ satisfies 2-out-of-6.
 (c) Find a collection of arrows satisfying 2-out-of-3 but not 2-out-of-6.

Exercise 2. A model structure on \mathbf{Cat} (Charles Rezk)

Denote \mathbf{Cat} the category of small categories. We assume that it is complete and cocomplete. We let \mathcal{W} denote equivalences of categories and \mathcal{C} denote functors that are injective on objects. We let \mathcal{F} denote functors $F : \mathcal{A} \rightarrow \mathcal{B}$ such that for every isomorphism $g : F(a) \rightarrow b$ of \mathcal{B} , there is a map $f : a \rightarrow a'$ with $g = F(f)$; such functors are called isofibrations.

1. Denote $*$ the category with one object and no non-trivial arrows, and I the category with two objects $0, 1$ and exactly one isomorphism in each direction. Let $i : * \rightarrow I$ be the inclusion at 0 . Show that $\mathcal{F} = RLP(\{i\})$.
2. Show that \mathcal{W} verifies 2-out-of-3, and that \mathcal{W}, \mathcal{C} and \mathcal{F} are stable under retracts.
3. (a) Show that every functor F of $\mathcal{C} \cap \mathcal{W}$ has a left inverse G which is also a quasi-inverse and such that the natural transformation $FG \simeq \text{id}$ is equal to the identity on the image of F .
 (b) Deduce that $\mathcal{F} \subset RLP(\mathcal{C} \cap \mathcal{W})$.
4. Show that $\mathcal{C} \subset LLP(\mathcal{F} \cap \mathcal{W})$.
5. Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be a functor. Denote $\text{Path}(F) := \mathcal{A} \times_{\mathcal{B}} \mathcal{B}^I$ where the map $\mathcal{B}^I \rightarrow \mathcal{B}$ is the source map, and $\text{Cyl}(F) := \mathcal{A} \times (I \coprod_{\mathcal{A}} \mathcal{B})$. Show that F factors through $\text{Path}(F)$ and $\text{Cyl}(F)$; deduce that $(\mathcal{C}, \mathcal{F}, \mathcal{W})$ is a model structure on \mathbf{Cat} . What are the fibrant objects, the cofibrant objects?

ON DERIVED FUNCTORS

Exercise 3 (Derived functors in homological algebra vs model categories). The goal of this exercise is to understand why the model-categorical notion of derived functor generalizes what you have seen in homological algebra. Let $\mathcal{A} \xrightarrow{F} \mathcal{B}$ be an additive functor between abelian categories. We suppose that F is right exact.

1. Consider the projective model structure on $Ch_{\geq 0}(\mathcal{A})$. Show that F sends quasi-isos between cofibrant objects to quasi-isos in $Ch_{\geq 0}\mathcal{B}$. Deduce that it has a total left derived functor in the model categorical sense.
2. Show that $\mathbb{L}F(V) \simeq F(P)$ where P is a projective resolution of V .
3. What is the link between the homological-algebraic derived functors $L^i F(V)$ and $\mathbb{L}F(V)$?

4. Apply this to prove the existence and identify the total derived functors of $\text{Hom}_R(-, M)$:

$$\text{Ch}_{\geq 0}(A)^{\text{op}} \rightarrow \text{Ch}_{\geq 0}(A).$$

Identify them with the derived functors $\text{Ext}^j(-, -)$ from the homological algebra course. Distinguish between the cases of projective and injective structures, and explain how this affects the computations.

5. Take $\mathcal{A} = R\text{Mod}$ for R a commutative ring. Do the functors $- \otimes - : \text{Ch}_{\geq 0}(\mathcal{A}) \times \text{Ch}_{\geq 0}(R) \rightarrow \text{Ch}_{\geq 0}(\mathcal{A})$ and $\text{Hom}(-, -) : \text{Ch}_{\geq 0}(\mathcal{A})^{\text{op}} \times \text{Ch}_{\geq 0}(\mathcal{A}) \rightarrow \text{Ch}_{\geq 0}(\mathcal{A})$ have total derived functors?

Exercise 4 (Composition of Derived Functors). Let $F : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ and $G : \mathcal{C}_2 \rightarrow \mathcal{C}_3$ be functors and let \mathcal{W}_i be a class of morphisms in \mathcal{C}_i .

1. Assuming all the relevant total left derived functors exist, use their universal properties to construct a natural transformation $\mathbb{L}G \circ \mathbb{L}F \rightarrow \mathbb{L}(G \circ F)$.
2. Suppose now that $\mathcal{C}_1, \mathcal{C}_2$ and \mathcal{C}_3 are model categories and that F and G are left Quillen functors. Show that $G \circ F$ is a left Quillen functor.
3. Show that the arrow of 1) induces a natural equivalence $\mathbb{L}(G \circ F) \simeq (\mathbb{L}G) \circ (\mathbb{L}F)$.
4. Let (L, R) be an adjoint pair. Show that L is left Quillen if and only if R is right Quillen.
5. Suppose the restriction of a functor F to cofibrant objects preserves acyclic cofibrations, show that F is left derivable. (Hint: Ken Brown's lemma).

Exercise 5 (Slice categories II, by Victor Saunier). Let $(\mathcal{C}, \mathcal{F}, \mathcal{W})$ be a model structure on \mathcal{A} . Let $f : X \rightarrow Y$ be a morphism. Recall that we defined in the last exercise sheet a model structure on every slice category \mathcal{A}/X .

1. Show that the functor $f_! : \mathcal{A}/X \rightarrow \mathcal{A}/Y$ which postcomposes by f admits a right adjoint f^* and describe it.
2. Show that the pair $(f_!, f^*)$ is a Quillen pair of adjoints.
3. Suppose \mathcal{A} is right proper, i.e. weak equivalences are stable under pullback by fibrations, and that $f \in \mathcal{W}$. Show that the pair $(f_!, f^*)$ is a Quillen equivalence.
4. (Rezk) Suppose that for every weak equivalence f , the pair $(f_!, f^*)$ is a Quillen equivalence. Show that \mathcal{A} is right proper.

ON HOMOTOPY (CO)LIMITS

Exercise 6 (Homotopy colimits). In this exercise, we first deal with generalities on homotopy pushouts and then specialize to chain complexes with the projective model structure. Let \mathcal{C} be a model category and let I be the category given by the diagram-shape

$$\begin{array}{ccc} b & \longrightarrow & c \\ \downarrow & & \\ a & & \end{array}$$

1. Let $f : X \rightarrow Y$ be a natural transformation of diagrams $X, Y \in \text{Fun}(I, \mathcal{C})$. Show that f has the left lifting property with respect to all projective acyclic fibrations if and only if the the natural maps

$$X_a \bigsqcup_{X_b} Y_b \rightarrow Y_a, \quad X_b \rightarrow Y_b, \quad X_c \bigsqcup_{X_b} Y_b \rightarrow Y_c$$

are cofibrations in \mathcal{C} . (Here we mean the usual pushouts in \mathcal{C} .)

Deduce that a diagram $Y : I \rightarrow \mathcal{C}$ is cofibrant if and only if Y_b is cofibrant in \mathcal{C} and the maps $Y_a \rightarrow Y_b$ and $Y_a \rightarrow Y_c$ are cofibrations. Moreover, show that $X \rightarrow Y$ has the left lifting property with respect to projective fibrations if and only if the above three maps are acyclic cofibrations.

2. Show that the category of diagrams $\text{Fun}(I, \mathcal{C})$ admits the projective model structure (without using the result seen in class that such a structure exists since I is very small).
3. Show that the colimit functor $\text{colim} : \text{Fun}(I, \mathcal{C}) \rightarrow \mathcal{C}$ is a left Quillen functor.
4. Assume that \mathcal{C} is left proper (i.e. weak equivalences are stable under pushouts along cofibrations). Show that any pushout diagram

$$\begin{array}{ccc} B & \xrightarrow{f} & C \\ \downarrow & & \downarrow \\ A & \longrightarrow & A \bigsqcup_B C \end{array}$$

where $f : B \rightarrow C$ a cofibration, is also a homotopy pushout diagram.

5. **Case of Topological spaces.** Assume now that $\mathcal{C} = \mathbf{Top}$.

- (a) Using that \mathbf{Top} is proper (as seen in exercise 3. from the sheet on Quillen model structure), show that there is a canonical isomorphism

$$\mathbb{L} \text{colim}(X \leftarrow A \rightarrow Y) \cong X \bigsqcup_A^h Y = X \bigsqcup_{A \times \{0\}} \text{Cyl}(A \rightarrow Y)$$

in $\text{Ho}(\mathbf{Top})$ between the homotopy pushout computed by the projective model structure and the formula given by the mapping cylinder.

- (b) Give a formula for computing the homotopy colimit of a tower $(X_0 \rightarrow X_1 \rightarrow X_2 \rightarrow \dots)$ as well as the homotopy limit of a tower $(\dots \rightarrow Y_2 \rightarrow Y_1 \rightarrow Y_0)$.
6. **Case of chain complexes.** Assume now that \mathcal{C} is the model category of chain complexes over a ring R .

- (a) Show that \mathcal{C} is left proper.
- (b) Let $g : A \rightarrow B$ be a map of chain complexes. Recall that the *mapping cone* of g , denoted $C(g)$, is the chain complex given in level n by $B_n \oplus A_{n-1}$ and whose differential $B_{n+1} \oplus A_n \rightarrow B_n \oplus A_{n-1}$ is given $(b, a) \mapsto (\partial_B(b) + g(a), -\partial_A(a))$. Let I denote the chain complex given by $R \oplus R$ in degree 0 and R in degree 1 with differential given by $\partial_R : R \rightarrow R \oplus R$ given by $r \mapsto (-r, r)$. We define the *mapping cylinder* of g , denoted $\text{Cyl}(g)$, as the pushout in chain complexes of

$$\begin{array}{ccc} A & \xrightarrow{g} & B \\ \downarrow i_0 & & \downarrow \\ I \otimes A & \longrightarrow & \text{Cyl}(g) \end{array}$$

where the vertical arrow $A \rightarrow I \otimes A$ is induced by the inclusion $i_0 : R \rightarrow I$ corresponding to the inclusion of the second factor $R \hookrightarrow R \oplus R$ in degree 0. The differential on $I \otimes A$ is given by $r \otimes a \mapsto \partial_R(r) \otimes a + (-1)^{\deg(r)} r \otimes \partial_A(a)$. Show that the mapping cone of g is the pushout of

$$\begin{array}{ccc} I \otimes A & \longrightarrow & \text{Cyl}(g) \\ \downarrow & & \downarrow \\ C(\text{Id}_A) & \longrightarrow & C(g). \end{array}$$

- (c) Let Δ^1 be the category with two objects and one non trivial morphism in between them. Show that the construction of the mapping cone defines a functor $C : \text{Fun}(\Delta^1, \text{Ch}(R)) \rightarrow \text{Ch}(R)$ sending natural transformations objectwise given by quasi-isomorphisms to quasi-isomorphisms.
- (d) Let $Y := (0 \leftarrow A \xrightarrow{g} B)$ be a diagram in \mathcal{C} . Show that there exists a diagram of the form $Y' := (0 \leftarrow A' \xrightarrow{g'} B')$ with g' a cofibration and A' and B' cofibrant, together with a natural transformation $u : Y' \rightarrow Y$ which is objectwise a weak equivalence. Notice that by the previous question the induced map $C(g') \rightarrow C(g)$ is a weak equivalence.
- (e) Let $Y := (0 \leftarrow A \xrightarrow{g} B)$ be a diagram in \mathcal{C} with A and B cofibrant and g a cofibration. Show that $A \rightarrow I \otimes A$ is a weak equivalence and show that we can construct a zigzag of diagrams $Y \leftarrow Y' \rightarrow Y''$ of the form

$$\begin{array}{ccccc}
0 & \longleftarrow & A & \xrightarrow{g} & B \\
\uparrow & & \uparrow & & \uparrow \\
C(A) & \longleftarrow & A & \xrightarrow{g} & B \\
\downarrow & & \downarrow & & \downarrow \\
C(A) & \longleftarrow & I \otimes A & \xrightarrow{g} & \text{Cyl}(g)
\end{array}$$

where each vertical arrow is a weak equivalence and the map $I \otimes A \rightarrow \text{Cyl}(g)$ is a cofibration.

- (f) Let $Y := (0 \leftarrow A \xrightarrow{g} B)$ be any diagram. Conclude that the mapping cone $C(g)$ is a model for the homotopy colimit of the diagram Y .