



WORKSHEET 3

CW COMPLEXES

Exercise 1 (Examples of CW complex).

- (1) For $n \in \mathbb{N}$, is \mathbb{R}^n a CW complex? It is a finite CW complex?
- (2) Show that the product $X \times Y$ of two CW complexes is again a CW complex.
- (3) Let (X, A) be a relative CW complex. Show that X/A is also a CW complex.
- (4) Show that the n -times iterated (unpointed) suspension $\Sigma^n X$ of a CW complex is again a CW complex, where

$$\Sigma X := \frac{X \times I}{X \times \partial I}.$$

- (5) Let $f : X \rightarrow Y$ be a cellular map between CW complexes. Show that the factorisation of f

$$X \xrightarrow{\quad} \text{Cyl}(f) \xrightarrow{\sim} Y$$

is the composite of two cellular maps.



Exercise 2 (C in CW complex).

- (1) Show that every CW complex X is in set-theoretical bijection with

$$X \cong \bigsqcup_{n \in \mathbb{N}, j \in J_n} D^n,$$

where $\mathring{D}^n = D^n \setminus \partial D^n$ is the interior of D^n , with $\mathring{D}^0 = D^0 = \{*\}$ by convention.

- (2) Show that any compact subspace of a CW complex intersects only a finite number of cells.



Exercise 3 (Infinite CW complexes).

- (1) Show that the n -dimension sphere S^n , for any $n \in \mathbb{N}$, admits a CW complex structure with two k -dimensional cells for any $0 \leq k \leq n$.

By passing to the colimit

$$S^0 \subset S^1 \subset S^2 \subset \dots \subset S^\infty = \bigcup_{n \geq 0} S^n,$$

this defines a new CW complex called the *infinite dimensional sphere* and denoted by S^∞ .

S^∞ and $\Sigma^\infty X$

- (2) Let X be a CW complex which is the union (colimit) of an increasing sequence of CW subcomplexes

$$X_1 \subset X_2 \subset X_3 \subset \dots \subset X = \bigcup_{n \geq 1} X_n$$


such that each inclusion $X_n \hookrightarrow X_{n+1}$ is nullhomotopic. Show that X is contractible.

- (3) Show that S^∞ is contractible.



Exercise 4 (Extension Lemma).

- (1) Let (X, A) be a relative CW complex and a map $f : A \rightarrow Y$ with Y path connected. Show that f extends to a map $F : X \rightarrow Y$



if $\pi_{n-1}(Y) = 0$ for any n such that $X \setminus A$ has cells of dimension n .

- (2) Let (X, A) be a CW pair such that A is contractible. Show that X retracts to A .



Exercise 5 (Cellular approximation).

- (1) Show that every map $f : X \rightarrow Y$ between CW complexes is homotopic to a cellular map.
 HINT: We will assume the following result, see [Hatcher, Lemma 4.10]. Let $f : (X, A) \rightarrow (Z, Y)$ be a map of CW-complexes such that the dimension of A is less than n , $X \setminus A$ is made up of only one n -dimensional cell, the dimension of Y is less than m , $Z \setminus Y$ is made up of only one cell of dimension $N \geq m$. The map f is homotopic relative to A to a map $g : (X, A) \rightarrow (Z, Y)$ such that $g(X \setminus A)$ misses a point of $Z \setminus Y$.
- (2) Prove that $\pi_k(S^n) = 0$ for $k < n$.
- (3) Show that every map $f : (X, A) \rightarrow (Y, B)$ between CW pairs is homotopic to a cellular map.
- (4) Let (X, A) be a CW pair such that $X \setminus A$ has only cells in dimension greater than n . Show that (X, A) is n -connected, that is $\pi_k(X, A) = 0$, for $k \leq n$.
- (5) Let X be a CW complex. Show that the CW pair $(X, X^{(n-1)})$ is n -connected and that the inclusion $X^{(n-1)} \hookrightarrow X$ induces isomorphisms

$$\pi_k(X^{(n-1)}) \cong \pi_k(X),$$

for $k < n$, and a surjection $\pi_n(X^{(n-1)}) \twoheadrightarrow \pi_n(X)$.




Exercise 6 (Postnikov tower).

- (1) Let X be a CW complex. For $n \geq 1$, show that there exists a CW complex X_n for which $X \subset X_n$ is a CW subcomplex and which satisfies

$$\begin{cases} \pi_k(X_n) \cong \pi_k(X), & \text{for } k \leq n, \\ \pi_k(X_n) \cong 0, & \text{for } k > n. \end{cases}$$

- (2) Show that there exists maps $X_{n+1} \rightarrow X_n$ which make the following diagram commutative



✉ Bruno Vallette: vallette@math.univ-paris13.fr .

🌐 Web page: www.math.univ-paris13.fr/~vallette/ .